

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 August 2003 (28.08.2003)

PCT

(10) International Publication Number
WO 03/070771 A1

(51) International Patent Classification⁷: C07K 16/00,
A61K 39/395

(21) International Application Number: PCT/US03/01858

(22) International Filing Date: 22 January 2003 (22.01.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10/056,790 23 January 2002 (23.01.2002) US

(71) Applicant: EXELIXIS, INC. [US/US]; P.O. Box 511, 170
Harbor Way, South San Francisco, CA 94083-0511 (US).

Hartmut; Waisenhausgasse 48, 50676 Cologne (DE).
ZEVNIK, Branko; Gezelinallee 46, Leverkusen 51375
(DE). SCHOOR, Michael; Am Scherfenbrand 168, 51375
Leverkusen (DE). REARDON, Brian, John; 444 N. El
Camino Real #101, San Mateo, CA 94401 (US).

(74) Agents: SHAYESTEH, Laleh et al.; Exelixis, Inc., P.O.
Box 511, 170 Harbor Way, South San Francisco, CA
94083-0511 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU,
ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI,
SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(72) Inventors: FRANCIS-LANG, Helen; 1782 Pacific
Avenue, Apt. 2, San Francisco, CA 94109 (US). FRIED-
MAN, Lori; 113 Arundel Road, San Carlos, CA 94070
(US). BELVIN, Marcia; 921 Santa Fe Avenue, Albany,
CA 94706 (US). PLOWMAN, Gregory, D.; 35 Winding
Way, San Carlos, CA 94070 (US). LARSON, Jeffrey,
S.; 1220 El Camino Real #305, Burlingame, CA 94010
(US). CHEN, Changyou; 6465 Crestwood Drive, Castro
Valley, CA 94552 (US). ROBERSTON, Stephanie,
A.; 255 Fowler Avenue, San Francisco, CA 94127 (US).
LIOUBIN, Mario, N.; 3014 Los Prados, #A310, San
Mateo, CA 94403 (US). SHI, Wen; 1324 8th Avenue,
#4, San Francisco, CA 94112 (US). CHAN, Jocelyn;
1049 Cortland Avenue, San Francisco, CA 94110 (US).
FUNKE, Roel, P.; 668, Sierra Point Road, Brisbane,
CA 95005 (US). LI, Danxi; 90 Behr Avenue, #302, San
Francisco, CA 94131 (US). KAUSELMANN, Gunther;
Muenstereifeler Str. 39, 50937 Koeln (DE). TINTRUP,

(54) Title: RRP SEQUENCES AND KNOCKOUT MICE AND USES THEREOF

(57) Abstract: RRP genes are identified as modulators of the p53 or p21 pathway, and thus are therapeutic targets for disorders associated with defective p53 or p21 function. Methods for identifying modulators of p53 or p21, comprising screening for agents that modulate the activity of RRP are provided. Modulating agents identified using the methods of the invention can be used to specifically inhibit growth of tumor cells that overexpress an RRP protein. mRRPI knockout mice are also provided.



WO 03/070771 A1

RRP SEQUENCES AND KNOCKOUT MICE AND USES THEREOF

REFERENCE TO RELATED APPLICATIONS

This application claims priority to US utility patent application 10/056,790 filed
5 1/23/2002, which is a Continuation In Part of US utility patent application 09/908,419
filed 7/18/2001, which claims priority to U.S. provisional patent applications 60/219,289,
filed 7/19/2000, 60/277,487, filed 3/21/2001, 60/277,471, filed 3/21/2001, 60/304,863,
filed 7/12/2001, 60/296,076 filed 6/5/2001, 60/305,017 filed 7/12/2001, 60/328,605 filed
10/10/2001, and 60/328,491 filed 10/10/2001. The contents of the prior applications are
10 hereby incorporated in their entirety.

BACKGROUND OF THE INVENTION

Signal transduction pathways are made up of growth factors, their receptors,
upstream regulators of the growth factors, and downstream intracellular kinase networks.
15 These pathways regulate many cellular processes, including proliferation, and appear to
play a key role in oncogenesis.

The epidermal growth factor receptor (EGFR) and its pathway members are among
the most widely explored signaling pathways. Signaling through this pathway elicits
diverse biological responses whose manifestations can include mitogenesis or apoptosis,
20 enhanced cell motility, protein secretion, and differentiation or dedifferentiation. Up-
regulated EGFR signaling has been implicated in organ morphogenesis, maintenance and
repair, and is correlated with invasion and metastasis of many types of tumors.
Upregulated EGF and EGFR in various tumor types leads to increased expression of
p21/Waf1/Cip1, leading to an arrest in the G1 phase of the cell cycle (Reddy KB et al.,
25 (1999) Int. J. Cancer 15:301-306).

Signal transduction pathways, such as the EGFR pathway, are evolutionarily
conserved among species as distant as the worm *Caenorhabditis elegans*, the fruit fly
Drosophila melanogaster, and vertebrates (Duffy JB, and Perrimon N, Curr. Opin. Cell
Biol. (1996) 8:231-238). In fact, ligands for the *Drosophila* EGFR (DER), known as Spitz
30 (Rutledge B, et al, Genes Dev. (1992) 6:1503-1517) and Gurken (Neuman-Silberberg FS,
and Schupbach T, Cell (1993) 75:165-174), are both similar to TGF α (transforming
growth factor alpha), the ligand for the vertebrate EGFR (Massague J, J Biol Chem.
(1990) 265:21393-21396). The rhomboid gene, which encodes a transmembrane protein,
is another upstream member of this pathway (Bier E., et al., Genes Dev. (1990) 4:190-

203). In *Drosophila*, rhomboid protein is a transmembrane serine protease, cleaves Spitz, and transforms Spitz from a membrane-bound to a secreted form, and thus triggers and upregulates the DER signaling pathway (Wasserman JD et al, Genes Devel (2000) 14:1651-1663; Lee JR, et al., (2001) Cell 107:161-171; Urban S, et al., (2001) Cell 107: 5 173-182). DNA sequences related to rhomboid have been identified in *C. elegans* (Wasserman JD, and Freeman M, Trends Cell Biol (1997) 7:431-436), and in mammals (Pascall JC, and Brown KD, FEBS letters (1998) 429:337-340; human: GI#3287191 and GI#7020534, among others; and rat: GI#3297936;), suggesting that rhomboid function may be evolutionarily conserved. Modulation of EGF-receptor related activity by 10 synthetic peptides or humanized monoclonal antibodies inhibit tumor growth (Baselga J, et al., (1998) Cancer Research 58:2825-2831; Park B-W, et al., (2000) Nature Biotechnology 18:194-197).

The p53 gene is mutated in over 50 different types of human cancers, including familial and spontaneous cancers, and is believed to be the most commonly mutated gene 15 in human cancer (Zambetti and Levine, FASEB (1993) 7:855-865; Hollstein, *et al.*, Nucleic Acids Res. (1994) 22:3551-3555). Greater than 90% of mutations in the p53 gene are missense mutations that alter a single amino acid that inactivates p53 function. Aberrant forms of human p53 are associated with poor prognosis, more aggressive tumors, metastasis, and short survival rates (Mitsudomi et al., Clin Cancer Res 2000 Oct; 20 6(10):4055-63; Koshland, Science (1993) 262:1953).

The human p53 protein normally functions as a central integrator of signals including DNA damage, hypoxia, nucleotide deprivation, and oncogene activation (Prives, Cell (1998) 95:5-8). In response to these signals, p53 protein levels are greatly increased with the result that the accumulated p53 activates cell cycle arrest (through activation of 25 p21/Waf1/Cip1) or apoptosis depending on the nature and strength of these signals. Indeed, multiple lines of experimental evidence have pointed to a key role for p53 as a tumor suppressor (Levine, Cell (1997) 88:323-331). For example, homozygous p53 "knockout" mice are developmentally normal but exhibit nearly 100% incidence of neoplasia in the first year of life (Donehower *et al.*, Nature (1992) 356:215-221).

30 The biochemical mechanisms and pathways through which p53 functions in normal and cancerous cells are not fully understood, but one clearly important aspect of p53 function is its activity as a gene-specific transcriptional activator. Among the genes with known p53-response elements are several with well-characterized roles in either

regulation of the cell cycle or apoptosis, including GADD45, p21/Waf1/Cip1, cyclin G, Bax, IGF-BP3, and MDM2 (Levine, Cell (1997) 88:323-331).

Modulating signal transduction pathway activity involved in tumor growth and development is essential in understanding the development of many cancers, and eventually, for the treatment of cancer. The ability to manipulate the genomes of model organisms such as *Drosophila* provides a powerful means to analyze signal transduction pathways that, due to significant evolutionary conservation, have direct relevance to more complex vertebrate organisms. Due to a high level of gene and pathway conservation, the strong similarity of cellular processes, and the functional conservation of genes between these model organisms and mammals, identification of the involvement of novel genes in particular pathways and their functions in such model organisms can directly contribute to the understanding of the correlative pathways and methods of modulating them in mammals (see, for example, Mechler BM et al., 1985 EMBO J 4:1551-1557; Gateff E. 1982 Adv. Cancer Res. 37: 33-74; Watson KL., et al., 1994 J Cell Sci. 18: 19-33; Miklos GL, and Rubin GM. 1996 Cell 86:521-529; Wassarman DA, et al., 1995 Curr Opin Gen Dev 5: 44-50; and Booth DR. 1999 Cancer Metastasis Rev. 18: 261-284). For example, a genetic screen can be carried out in an invertebrate model organism having underexpression (e.g. knockout) or overexpression of a gene (referred to as a "genetic entry point") that yields a visible phenotype. Additional genes are mutated in a random or targeted manner. When a gene mutation changes the original phenotype caused by the mutation in the genetic entry, the gene is identified as a "modifier" involved in the same or overlapping pathway as the genetic entry point. When the genetic entry point is an ortholog of a human gene implicated in a disease pathway, such as p53 or p21, modifier genes can be identified that may be attractive candidate targets for novel therapeutics.

All references cited herein, including sequence information in referenced Genbank identifier numbers and website references, are incorporated herein in their entireties.

SUMMARY OF THE INVENTION

We have discovered genes that modify the p53 or p21 pathway in *Drosophila*, and identified their mammalian orthologs, hereinafter referred to as Rhomboid Related Proteins (RRP), and more specifically, RRP1 – RRP8, and mouse RRP1 (mRRP1). The invention provides isolated nucleic acid molecules that comprise nucleic acid sequences encoding RRP protein as well as fragments and derivatives thereof. Vectors and host cells comprising the RRP nucleic acid molecules are also described.

The invention provides methods for utilizing these p53 or p21 modifier genes and polypeptides to identify RRP modulating agents, which are candidate therapeutic agents that can be used in the treatment of disorders associated with defective p53 or p21 function.

5 In one embodiment, candidate p53 or p21 modulating agents are tested with an assay system comprising a RRP polypeptide or nucleic acid. Candidate agents that produce a change in the activity of the assay system relative to controls are identified as candidate p53 or p21 modulating agents. The assay system may be cell-based or cell-free. Candidate modulating agents include small molecule modulators, antibodies, and nucleic
10 acid modulators. In one specific embodiment, a small molecule modulator is identified using a protease assay. In specific embodiments, the screening assay system is selected from an apoptosis assay, a cell proliferation assay, an angiogenesis assay, and a hypoxic induction assay.

 In another embodiment, candidate p53 or p21 pathway modulating agents are
15 further tested using a second assay system that detects changes in the p53 or p21 pathway, such as angiogenic, apoptotic, or cell proliferation changes produced by the originally identified candidate agent or an agent derived from the original agent. The second assay system may use cultured cells or non-human animals. In specific embodiments, the secondary assay system uses non-human animals, including animals predetermined to
20 have a disease or disorder implicating the p53 or p21 pathway, such as an angiogenic, apoptotic, or cell proliferation disorder (e.g. cancer).

 The invention further provides methods for modulating the p53 or p21 pathway in a mammalian cell by contacting the mammalian cell with an agent that specifically binds a RRP polypeptide or nucleic acid. The agent may be a small molecule modulator, a nucleic
25 acid modulator, or an antibody and may be administered to a mammalian animal predetermined to have a pathology associated the p53 or p21 pathway.

 Modulating agents identified using the methods of the invention can be used to specifically inhibit growth of tumor cells that overexpress an RRP protein.

 The invention also provides transgenic knockout mice harboring disrupted RRP
30 genes. The disruption may be heterozygous, leading to decreased expression of RRP, or homozygous, leading to lack of expression of the RRP gene. Cells from the mice as well as cells harboring disrupted RRP genes are also provided. Methods of producing antibody to RRP using the mice of the invention are also provided.

Targeting vectors to produce transgenic knockout mice are also provided. Preferably, a targeting vector is provided that allows sequential deletion of vector sequences from the same cell in the generation of the knockout mice.

5 **BRIEF DESCRIPTION OF THE DRAWINGS**

FIG.1 depicts the targeting vector, pEasyfloxFRTGK12 mRRP1.

FIG.2 shows the mRRP1 mouse targeting vector, the mRRP1 wild type (WT) genomic region and the locations to which the vector is targeted, and the mRRP1 target allele after homologous recombination (HR). Locations of the 5' external probe, the
10 internal probe, and the 3' external probe are underlined. The expected sizes of the genomic fragments after digestion with the indicated restriction enzymes and Southern hybridization are also indicated.

FIG.3 depicts the target allele after HR and after Cre induced deletion. The expected sizes of the genomic fragments after digestion with restriction enzymes and
15 Southern hybridization are indicated for the target allele in each of the following states: WT, HR, and Cre deleted. FIG.4 depicts the target allele after HR, after FLP induced deletion, and after Cre induced deletion. The expected sizes of the genomic fragments after digestion with the indicated restriction enzymes and Southern hybridization are indicated for the target allele in each of the following states: WT, HR, FLP deleted, and
20 Cre deleted.

DETAILED DESCRIPTION OF THE INVENTION

Genetic screens were designed to identify modifiers of the p53 or p21 pathways in *Drosophila*. Genetic modifier screens were carried out in which p53 (Ollmann M, et al.,
25 Cell 2000 101: 91-101) or p21 (Bourne HR, et al., Nature (1990) 348(6297):125-132; Marshall CJ, Trends Genet (1991) 7(3):91-95) were overexpressed. *Drosophila* rhomboid genes were identified as modifiers of the p53 or p21 pathways. Accordingly, vertebrate orthologs of these modifiers, hereinafter referred to as RRP genes (i.e., nucleic acids and polypeptides), are attractive drug targets for the treatment of pathologies associated with a
30 defective p53 or p21 signaling pathways, such as cancer. Further, gene targeting in mice is an ideal method to investigate the function of a distinct protein in wild type and disease states. In order to study the RRP1 function in mammals we generated the genomic sequence of the RRP1 region in mice, deduced its cDNA and protein sequence, and then produced targeted RRP1 knockout (KO) mice.

In vitro and in vivo methods of assessing RRP function as provided herein, and modulation of the RRP or their respective binding partners is useful for understanding the association of the p53 or p21 pathways and their members in normal and disease conditions and for developing diagnostics and therapeutic modalities for p53 or p21 related pathologies. RRP-modulating agents that act by inhibiting or enhancing RRP expression, directly or indirectly, for example, by affecting an RRP function such as enzymatic (e.g., catalytic) or binding activity, can be identified using methods provided herein. RRP-modulating agents include RRP related proteins (e.g. dominant negative mutants, and biotherapeutics); RRP-specific antibodies; RRP-specific antisense oligomers; and chemical agents that specifically bind RRP or compete with RRP binding target. The invention provides methods of identifying and making RRP modulating agents, and their use in diagnosis, therapy and pharmaceutical development.

Preferred RRP-modulating agents specifically bind to RRP polypeptides and enhance or inhibit RRP function. Other preferred RRP-modulating agents are antisense oligomers and RNAi that repress RRP gene expression or product activity by, for example, binding to and inhibiting the respective nucleic acid (i.e. DNA or mRNA). RRP-specific modulating agents may be evaluated by any convenient *in vitro* or *in vivo* assay for molecular interaction with an RRP polypeptide or nucleic acid.

The method of this invention is useful in the therapy of malignant or benign tumors of mammals that overexpress RRP gene products.

Nucleic acids and polypeptides of the invention

Sequences related to RRP nucleic acids (RRP1: SEQ ID NO:1, RRP2: SEQ ID NO:3, RRP3: SEQ ID NO:5, RRP4: SEQ ID NO:7, RRP5: SEQ ID NO:9, RRP6: SEQ ID NO:11, RRP7: SEQ ID NO:13, and RRP8: SEQ ID NO:15) and polypeptides (RRP1: SEQ ID NO:2, RRP2: SEQ ID NO: 4, RRP3: SEQ ID NO:6, RRP4: SEQ ID NO:8, RRP5: SEQ ID NO:10, RRP6: SEQ ID NO:12, RRP7: SEQ ID NO:14, and RRP8: SEQ ID NO:16) are available in the public databases (for RRP1: cDNA: GI#3287190, SEQ ID NO:18; proteins GI#3287191, SEQ ID NO:36; for RRP2: cDNAs: GI#s: 12762689 (SEQ ID NO:19), 12096415 (SEQ ID NO:20), 6657080 (SEQ ID NO:21), 7947756 (SEQ ID NO:22), 3117010 (SEQ ID NO:23), 9339870 (SEQ ID NO:24), 7152496 (SEQ ID NO:25), 11317512 (SEQ ID NO:26), 11317513 (SEQ ID NO:27), and 11317511 (SEQ ID NO:28); proteins: GI#s:8923409 (SEQ ID NO:37), 12719522 (SEQ ID NO:38), and 11421817 (SEQ ID NO:39); for RRP3: cDNA: GI#10199673 (SEQ ID NO:29) and

GI#2003992 (SEQ ID NO:30); for RRP4: cDNA GI#11066249 (SEQ ID NO:31) and protein GI#11066250 (SEQ ID NO:40); for RRP5: cDNA GI#11967982 (SEQ ID NO:32) and protein GI#11967983 (SEQ ID NO:41); for RRP6: cDNA GI#10438685 (SEQ ID NO:33) and protein GI#10438686 (SEQ ID NO:42); for RRP7: cDNA GI#10190733 (SEQ ID NO:34) and protein GI#10190734 (SEQ ID NO:43); and for RRP8: cDNA GI#11072100 (SEQ ID NO:35) and protein GI#11072101 (SEQ ID NO:44)). Sequences of human and rat RRP1 were used to deduce the mouse RRP1 (mRRP1) cDNA (SEQ ID NO:45), polypeptide (SEQ ID NO:46), and genomic (SEQ ID NO:47) sequences, as described in Example VII. The mRRP1 cDNA sequence shares 69% identity with human RRP1 and 88% identity with rat partial RRP1 for nucleotides 884-1340 of the mouse RRP1. The mRRP1 protein shares 80% identity with human RRP1 and 88% identity with rat partial RRP1 for amino acids 297-448 of mRRP1.

RRPs are a family of integral membrane proteins that contain five or more transmembrane domains and three strongly conserved histidine residues in the putative transmembrane regions. In a preferred embodiment, the invention provides RRP proteins which comprise or consist of an amino acid sequence of SEQ ID NOs:4, 6, or 46, or fragments or derivatives thereof.

The term "RRP polypeptide" refers to a full-length RRP protein or a functionally active fragment or derivative thereof. A "functionally active" RRP fragment or derivative exhibits one or more functional activities associated with a full-length, wild-type RRP protein, such as antigenic or immunogenic activity, enzymatic activity, ability to bind natural cellular substrates, etc. The functional activity of RRP proteins, derivatives and fragments can be assayed by various methods known to one skilled in the art (Current Protocols in Protein Science (1998) Coligan *et al.*, eds., John Wiley & Sons, Inc., Somerset, New Jersey) and as further discussed below. For purposes herein, functionally active fragments also include those fragments that comprise one or more structural domains of an RRP, such as a protease or rhomboid domain or a binding domain. Catalytic and other domains can be identified using the PFAM program (Bateman A., *et al.*, Nucleic Acids Res, 1999, 27:260-2; website at pfam.wustl.edu). Methods for obtaining RRP polypeptides are also further described below. Preferred fragments are functionally active, domain-containing fragments sharing at least 80% sequence identity or similarity, preferably at least 85%, more preferably at least 90%, and most preferably at least 95% sequence identity or similarity with a contiguous stretch of at least 25 amino acids, preferably at least 50 amino acids, more preferably at least 100 amino acids, and in

some cases, the entire length of any one of SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, 16, and 46. In further preferred embodiments, the fragment comprises the entire rhomboid domain (PFAM 01694).

RRP protein derivatives typically share a certain degree of sequence identity or
5 sequence similarity with SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, 16, or 46 or a fragment thereof. RRP derivatives can be produced by various methods known in the art. The manipulations which result in their production can occur at the gene or protein level. For example, a cloned RRP gene sequence can be cleaved at appropriate sites with restriction endonuclease(s) (Wells *et al.*, Philos. Trans. R. Soc. London SerA (1986) 317:415),
10 followed by further enzymatic modification if desired, isolated, and ligated *in vitro*, and expressed to produce the desired derivative. Alternatively, an RRP gene can be mutated *in vitro* or *in vivo*, to create and/or destroy translation, initiation, and/or termination sequences, or to create variations in coding regions and/or to form new restriction endonuclease sites or destroy preexisting ones, to facilitate further *in vitro* modification.
15 A variety of mutagenesis techniques are known in the art such as chemical mutagenesis, *in vitro* site-directed mutagenesis (Carter *et al.*, Nucl. Acids Res. (1986) 13:4331), use of TAB[®] linkers (available from Pharmacia and Upjohn, Kalamazoo, MI), *etc.*

At the protein level, manipulations include post translational modification, *e.g.* glycosylation, acetylation, phosphorylation, amidation, derivatization by known
20 protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, *etc.* Any of numerous chemical modifications may be carried out by known technique (*e.g.* specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH₄, acetylation, formylation, oxidation, reduction, metabolic synthesis in the presence of tunicamycin, *etc.*). Derivative proteins can also be
25 chemically synthesized by use of a peptide synthesizer, for example to introduce nonclassical amino acids or chemical amino acid analogs as substitutions or additions into the RRP protein sequence.

Chimeric or fusion proteins can be made comprising an RRP protein or fragment thereof (preferably comprising one or more structural or functional domains of the RRP
30 protein) joined at its amino- or carboxy-terminus via a peptide bond to an amino acid sequence of a different protein. Chimeric proteins can be produced by any known method, including: recombinant expression of a nucleic acid encoding the protein (comprising a RRP-coding sequence joined in-frame to a coding sequence for a different protein); ligating the appropriate nucleic acid sequences encoding the desired amino acid sequences

to each other in the proper coding frame, and expressing the chimeric product; and protein synthetic techniques, *e.g.* by use of a peptide synthesizer.

The subject RRP polypeptides also encompass minor deletion mutants, including N-, and/or C-terminal truncations. Such deletion mutants are readily screened for RRP
5 competitive or dominant negative activity.

The term "RRP nucleic acid" refers to a DNA or RNA molecule that encodes a RRP polypeptide. In preferred embodiments, the nucleic acid encodes a polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, 16, and 46. In some embodiments, the nucleic acid comprises a sequence selected from the group
10 consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15, and 45. In a specific embodiment, the invention provides an isolated nucleic acid which encodes a human RRP3 as shown in SEQ ID NO:5, and also an isolated nucleic acid that encodes a mouse RRP (mRRP1) as shown in SEQ ID NO:45.

The invention includes a fragment of a nucleic acid, such as a fragment that
15 encodes a binding domain of one of the full-length sequences of the invention. Fragments of an RRP nucleic acid sequence can be used for a variety of purposes. As an example, interfering RNA (RNAi) fragments, particularly double-stranded (ds) RNAi, can be used to generate loss-of-function phenotypes; which can, in turn, be used, among other uses, to determine gene function. Certain "antisense" fragments, *i.e.* that are reverse complements
20 of portions of the coding and/or untranslated regions (*e.g.* 5' UTR) have utility in inhibiting the function of RRP proteins. The fragments are of length sufficient to specifically hybridize with the corresponding RRP sequence. The fragments consist of or comprise at least 12, preferably at least 24, more preferably at least 36, and more preferably at least 96 contiguous nucleotides of RRP. When the fragments are flanked by
25 other nucleic acid sequences, the total length of the combined nucleic acid sequence is less than 15 kb, preferably less than 10 kb or less than 5kb, more preferably less than 2 kb, and in some cases, preferably less than 500 bases.

In other specific embodiments, preferred fragments of SEQ ID NO:5 encode extracellular or intracellular domains which are located at approximately nucleotides 248-
30 598, 665-796, 862-870, 934-943, 1006-1138, 1201-1225, and 1289-1336. Additional preferred fragments of SEQ ID NO:45 encode extracellular or intracellular domains which are located at approximately nucleotides 1-714, 774-912, 972-984, 1044-1089, 1149-1212, 1272-1305, 1365-1408. Preferred fragments may also include a binding domain or an RRP motif (*e.g.* PFAM 01694). These domains may be useful to locate the function

and/or binding partners of a protein. For example, a nucleic acid that encodes an extracellular or intracellular domain of a protein may be used to screen for binding partners related to the protein.

The subject nucleic acid sequences may consist solely of the RRP nucleic acid or fragments thereof. Alternatively, the subject nucleic acid sequences and fragments thereof may be joined to other components such as labels, peptides, agents that facilitate transport across cell membranes, hybridization-triggered cleavage agents or intercalating agents. The subject nucleic acid sequences and fragments thereof may also be joined to other nucleic acid sequences (i.e. they may comprise part of larger sequences) and are of synthetic/non-natural sequences and/or are isolated and/or are purified, i.e. unaccompanied by at least some of the material with which it is associated in its natural state. Preferably, the isolated nucleic acids constitute at least about 0.5%, and more preferably at least about 5% by weight of the total nucleic acid present in a given fraction, and are preferably recombinant, meaning that they comprise a non-natural sequence or a natural sequence joined to nucleotide(s) other than that which it is joined to on a natural chromosome.

The subject nucleic acids find a wide variety of applications including use as translatable transcripts, hybridization probes, PCR primers, diagnostic nucleic acids, etc.; use in detecting the presence of RRP genes and gene transcripts and in detecting or amplifying nucleic acids encoding additional RRP homologs and structural analogs. In diagnosis, RRP hybridization probes find use in identifying wild-type and mutant RRP alleles in clinical and laboratory samples. Mutant alleles are used to generate allele-specific oligonucleotide (ASO) probes for high-throughput clinical diagnoses. In therapy, therapeutic RRP nucleic acids are used to modulate cellular expression or intracellular concentration or availability of active RRP. In a preferred embodiment, the mouse RRP1 sequence is used to produce a targeting vector for production of mice that are deficient in RRP1 in a heterozygous or homozygous (i.e., knockout) manner.

In one preferred embodiment, the derivative nucleic acid encodes a polypeptide comprising a RRP3 amino acid sequence of SEQ ID NO:6, an mRRP1 amino acid sequence of SEQ ID NO:46, or a fragment or derivative thereof. A derivative RRP3 nucleic acid sequence, or fragment thereof, may comprise 100% sequence identity with SEQ ID NO:5 or 45, but be a derivative thereof in the sense that it has one or more modifications at the base or sugar moiety, or phosphate backbone. Examples of modifications are well known in the art (Bailey, Ullmann's Encyclopedia of Industrial

Chemistry (1998), 6th ed. Wiley and Sons). Such derivatives may be used to provide modified stability or any other desired property.

Preferably, the RRP polypeptide nucleic acid, fragment, ortholog, or derivative thereof has at least 70% sequence identity, preferably at least 80%, more preferably 85%,
5 still more preferably 90%, and most preferably at least 95% sequence identity with RRP. Normally, orthologs in different species retain the same function, due to presence of one or more protein motifs and/or 3-dimensional structures. As used herein, "percent (%) sequence identity" with respect to a subject sequence, or a specified portion of a subject
10 derivative sequence identical with the nucleotides or amino acids in the subject sequence (or specified portion thereof), after aligning the sequences and introducing gaps, if necessary to achieve the maximum percent sequence identity, as generated by the program WU-BLAST-2.0a19 (Altschul *et al.*, J. Mol. Biol. (1997) 215:403-410; website at blast.wustl.edu/blast/README.html) with all the search parameters set to default values.
15 The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched. A % identity value is determined by the number of matching identical nucleotides or amino acids divided by the sequence length for which the percent identity is being reported. "Percent
20 (%) amino acid sequence similarity" is determined by doing the same calculation as for determining % amino acid sequence identity, but including conservative amino acid substitutions in addition to identical amino acids in the computation.

A conservative amino acid substitution is one in which an amino acid is substituted for another amino acid having similar properties such that the folding or activity of the
25 protein is not significantly affected. Aromatic amino acids that can be substituted for each other are phenylalanine, tryptophan, and tyrosine; interchangeable hydrophobic amino acids are leucine, isoleucine, methionine, and valine; interchangeable polar amino acids are glutamine and asparagine; interchangeable basic amino acids are arginine, lysine and histidine; interchangeable acidic amino acids are aspartic acid and glutamic acid; and
30 interchangeable small amino acids are alanine, serine, threonine, cysteine and glycine.

Alternatively, an alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman (Smith and Waterman, 1981, *Advances in Applied Mathematics* 2:482-489; Smith and Waterman, 1981, *J. of Molec.Biol.*, 147:195-197; Nicholas *et al.*, 1998, "A Tutorial on Searching Sequence Databases and Sequence

Scoring Methods" (website at www.psc.edu) and references cited therein.; W.R. Pearson, 1991, *Genomics* 11:635-650). This algorithm can be applied to amino acid sequences by using the scoring matrix developed by Dayhoff (Dayhoff: *Atlas of Protein Sequences and Structure*, M. O. Dayhoff ed., 5 suppl. 3:353-358, National Biomedical Research Foundation, Washington, D.C., USA), and normalized by Gribskov (Gribskov 1986 *Nucl. Acids Res.* 14(6):6745-6763). The Smith-Waterman algorithm is used to search databases for sequences similar to a query sequence. Smith-Waterman uses dynamic programming to determine how an optimal alignment between the query sequence and a database sequence can be produced. This alignment is obtained by determining what transformations the query sequence would need to undergo to match the database sequence. Transformations include substituting one character for another and inserting or deleting a string of characters. A score is assigned for each character-to-character comparison--positive scores for exact matches and some substitutions, negative scores for other substitutions and insertions/deletions. The first character in an insertion or deletion gap is scored with a gap open penalty and subsequent characters are scored with a gap extension penalty. Scores are obtained from statistically-derived scoring matrices. The combination of transformations that results in the highest score is used to generate an alignment between the query sequence and database sequence. Smith-Waterman algorithm may be employed where default parameters are used for scoring (for example, gap open penalty of 12, gap extension penalty of two). From the data generated the "Match" value reflects "sequence identity."

Derivative nucleic acid molecules of the subject nucleic acid molecules include sequences that hybridize to the nucleic acid sequence of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, or 45. The stringency of hybridization can be controlled by temperature, ionic strength, pH, and the presence of denaturing agents such as formamide during hybridization and washing. Conditions routinely used are set out in readily available procedure texts (*e.g.*, *Current Protocol in Molecular Biology*, Vol. 1, Chap. 2.10, John Wiley & Sons, Publishers (1994); Sambrook *et al.*, *Molecular Cloning*, Cold Spring Harbor (1989)). In some embodiments, a nucleic acid molecule of the invention is capable of hybridizing to a nucleic acid molecule containing the nucleotide sequence of any one of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, or 45 under stringent hybridization conditions that comprise: prehybridization of filters containing nucleic acid for 8 hours to overnight at 65°C in a solution comprising 6X single strength citrate (SSC) (1X SSC is 0.15 M NaCl, 0.015 M Na citrate; pH 7.0), 5X Denhardt's solution, 0.05% sodium pyrophosphate and

100 µg/ml herring sperm DNA; hybridization for 18-20 hours at 65° C in a solution containing 6X SSC, 1X Denhardt's solution, 100 µg/ml yeast tRNA and 0.05% sodium pyrophosphate; and washing of filters at 65° C for 1h in a solution containing 0.2X SSC and 0.1% SDS (sodium dodecyl sulfate).

- 5 In other embodiments, moderately stringent hybridization conditions are used that comprise: pretreatment of filters containing nucleic acid for 6 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatured salmon sperm DNA; hybridization for 18-20h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl
10 (pH7.5), 5mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 µg/ml salmon sperm DNA, and 10% (wt/vol) dextran sulfate; followed by washing twice for 1 hour at 55° C in a solution containing 2X SSC and 0.1% SDS.

- Alternatively, low stringency conditions can be used that comprise: incubation for 8 hours to overnight at 37° C in a solution comprising 20% formamide, 5 x SSC, 50 mM
15 sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured sheared salmon sperm DNA; hybridization in the same buffer for 18 to 20 hours; and washing of filters in 1 x SSC at about 37° C for 1 hour.

20 **Isolation, Production, Expression, and Mis-expression of RRP Nucleic Acids and Polypeptides**

- RRP nucleic acids and polypeptides, useful for identifying and testing agents that modulate RRP function and for other applications related to the involvement of RRP in the p53 or p21 pathways. RRP nucleic acids and derivatives and orthologs thereof may be obtained using any available method. For instance, techniques for isolating cDNA or
25 genomic DNA sequences of interest by screening DNA libraries or by using polymerase chain reaction (PCR) are well known in the art. In general, the particular use for the protein will dictate the particulars of expression, production, and purification methods. For instance, production of proteins for use in screening for modulating agents may require methods that preserve specific biological activities of these proteins, whereas
30 production of proteins for antibody generation may require structural integrity of particular epitopes. Expression of proteins to be purified for screening or antibody production may require the addition of specific tags (*e.g.*, generation of fusion proteins). Overexpression of an RRP protein for assays used to assess RRP function, such as involvement in cell cycle regulation or hypoxic response, may require expression in eukaryotic cell lines

capable of these cellular activities. Techniques for the expression, production, and purification of proteins are well known in the art; any suitable means therefore may be used (e.g., Higgins SJ and Hames BD (eds.) *Protein Expression: A Practical Approach*, Oxford University Press Inc., New York 1999; Stanbury PF et al., *Principles of Fermentation Technology*, 2nd edition, Elsevier Science, New York, 1995; Doonan S (ed.) *Protein Purification Protocols*, Humana Press, New Jersey, 1996; Coligan JE et al, *Current Protocols in Protein Science* (eds.), 1999, John Wiley & Sons, New York; U.S. Pat. No. 6,165,992). In particular embodiments, recombinant RRP is expressed in a cell line known to have defective p53 or p21 function (e.g. for p53: SAOS-2 osteoblasts, H1299 lung cancer cells, C33A and HT3 cervical cancer cells, HT-29 and DLD-1 colon cancer cells, among others, and for p21: HCT116 colon cancer cells, among others, available from American Type Culture Collection (ATCC), Manassas, VA). The recombinant cells are used in cell-based screening assay systems of the invention, as described further below.

The nucleotide sequence encoding an RRP polypeptide can be inserted into any appropriate expression vector. The necessary transcriptional and translational signals, including promoter/enhancer element, can derive from the native RRP gene and/or its flanking regions or can be heterologous. A variety of host-vector expression systems may be utilized, such as mammalian cell systems infected with virus (e.g. vaccinia virus, adenovirus, *etc.*); insect cell systems infected with virus (e.g. baculovirus); microorganisms such as yeast containing yeast vectors, or bacteria transformed with bacteriophage, plasmid, or cosmid DNA. A host cell strain that modulates the expression of, modifies, and/or specifically processes the gene product may be used.

To detect expression of the RRP gene product, the expression vector can comprise a promoter operably linked to an RRP gene nucleic acid, one or more origins of replication, and, one or more selectable markers (e.g. thymidine kinase activity, resistance to antibiotics, *etc.*). Alternatively, recombinant expression vectors can be identified by assaying for the expression of the RRP gene product based on the physical or functional properties of the RRP protein in *in vitro* assay systems (e.g. immunoassays).

The RRP protein, fragment, or derivative may be optionally expressed as a fusion, or chimeric protein product (i.e. it is joined via a peptide bond to a heterologous protein sequence of a different protein), for example to facilitate purification or detection. A chimeric product can be made by ligating the appropriate nucleic acid sequences encoding the desired amino acid sequences to each other using standard methods and expressing the

chimeric product. A chimeric product may also be made by protein synthetic techniques, *e.g.* by use of a peptide synthesizer (Hunkapiller *et al.*, Nature (1984) 310:105-111).

Once a recombinant cell that expresses the RRP gene sequence is identified, the gene product can be isolated and purified using standard methods (*e.g.* ion exchange, affinity, and gel exclusion chromatography; centrifugation; differential solubility; electrophoresis, *cite purification reference*). Alternatively, native RRP proteins can be purified from natural sources, by standard methods (*e.g.* immunoaffinity purification). Once a protein is obtained, it may be quantified and its activity measured by appropriate methods, such as immunoassay, bioassay, or other measurements of physical properties, such as crystallography.

The methods of this invention may also use cells that have been engineered for altered expression (mis-expression) of RRP or other genes associated with the p53 or p21 pathway. As used herein, mis-expression encompasses ectopic expression, over-expression, under-expression, and non-expression (*e.g.* by gene knock-out or blocking expression that would otherwise normally occur).

Genetically modified animals

Animal models that have been genetically modified to alter RRP expression may be used in *in vivo* assays to test for activity of a candidate p53 or p21 modulating agent, or to further assess the role of RRP in a p53 or p21 pathway process such as apoptosis or cell proliferation. Preferably, the altered RRP expression results in a detectable phenotype, such as decreased or increased levels of cell proliferation, angiogenesis, or apoptosis compared to control animals having normal RRP expression. The genetically modified animal may additionally have altered p53 or p21 expression (*e.g.* p53 or p21 knockout). Preferred genetically modified animals are mammals such as primates, rodents (preferably mice), cows, horses, goats, sheep, pigs, dogs and cats. Preferred non-mammalian species include zebrafish, *C. elegans*, and *Drosophila*. Preferred genetically modified animals are transgenic animals having a heterologous nucleic acid sequence present as an extrachromosomal element in a portion of its cells, *i.e.* mosaic animals (see, for example, techniques described by Jakobovits, 1994, Curr. Biol. 4:761-763.) or stably integrated into its germ line DNA (*i.e.*, in the genomic sequence of most or all of its cells). Heterologous nucleic acid is introduced into the germ line of such transgenic animals by genetic manipulation of, for example, embryos or embryonic stem cells of the host animal.

Methods of making transgenic animals are well-known in the art (for transgenic mice see Brinster *et al.*, Proc. Nat. Acad. Sci. USA 82: 4438-4442 (1985), U.S. Pat. Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Pat. No. 4,873,191 by Wagner *et al.*, U.S. Pat. No. 6,127,598, by German *et al.*, and Hogan, B., Manipulating the Mouse Embryo, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1986); for particle bombardment see U.S. Pat. No., 4,945,050, by Sandford *et al.*; for transgenic *Drosophila* see Rubin and Spradling, Science (1982) 218:348-53 and U.S. Pat. No. 4,670,388; for transgenic insects see Berghammer A.J. *et al.*, A Universal Marker for Transgenic Insects (1999) Nature 402:370-371; for transgenic Zebrafish see Lin S., Transgenic Zebrafish, Methods Mol Biol. (2000);136:375-3830); for microinjection procedures for fish, amphibian eggs and birds see Houdebine and Chourrout, Experientia (1991) 47:897-905; for transgenic rats see Hammer *et al.*, Cell (1990) 63:1099-1112; and for culturing of embryonic stem (ES) cells and the subsequent production of transgenic animals by the introduction of DNA into ES cells using methods such as electroporation, calcium phosphate/DNA precipitation and direct injection see, e.g., Teratocarcinomas and Embryonic Stem Cells, A Practical Approach, E. J. Robertson, ed., IRL Press (1987)). Clones of the nonhuman transgenic animals can be produced according to available methods (see Wilmut, I. *et al.* (1997) Nature 385:810-813; and PCT International Publication Nos. WO 97/07668 and WO 97/07669).

In one embodiment, the transgenic animal is a "knock-out" animal having a heterozygous or homozygous alteration in the sequence of an endogenous RRP gene that results in a decrease of RRP function, preferably such that RRP expression is undetectable or insignificant. Knock-out animals are typically generated by homologous recombination with a vector comprising a transgene having at least a portion of the gene to be knocked out. Typically a deletion, addition or substitution has been introduced into the transgene to functionally disrupt it. The transgene can be a human gene (e.g., from a human genomic clone) but more preferably is an ortholog of the human gene derived from the transgenic host species. For example, a mouse RRP gene is used to construct a homologous recombination vector suitable for altering an endogenous RRP gene in the mouse genome as shown in Example VIII. Detailed methodologies for homologous recombination in mice are available (see Capecchi, Science (1989) 244:1288-1292; Joyner *et al.*, Nature (1989) 338:153-156). Procedures for the production of non-rodent transgenic mammals and other animals are also available (Houdebine and Chourrout, *supra*; Pursel *et al.*, Science (1989) 244:1281-1288; Simms *et al.*, Bio/Technology (1988)

6:179-183). In a preferred embodiment, knock-out animals, such as mice harboring a knockout of a specific gene, may be used to produce antibodies against the human counterpart of the gene that has been knocked out (Claesson MH et al. (1994) Scand J Immunol 40:257-264; Declerck PJ, et al., (1995) J Biol Chem 270:8397-8400).

5 In another embodiment, the transgenic animal is a "knock-in" animal having an alteration in its genome that results in altered expression (e.g., increased (including ectopic) or decreased expression) of the RRP gene, e.g., by introduction of additional copies of RRP, or by operatively inserting a regulatory sequence that provides for altered expression of an endogenous copy of the RRP gene. Such regulatory sequences include
10 inducible, tissue-specific, and constitutive promoters and enhancer elements. The knock-in can be homozygous or heterozygous.

Transgenic nonhuman animals can also be produced that contain selected systems allowing for regulated expression of the transgene. One example of such a system that may be produced is the cre/loxP recombinase system of bacteriophage P1 (Lakso *et al.*,
15 PNAS (1992) 89:6232-6236; U.S. Pat. No. 4,959,317). If a cre/loxP recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other
20 containing a transgene encoding a recombinase. Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae* (O'Gorman et al. (1991) Science 251:1351-1355; U.S. Pat. No. 5,654,182). In a preferred embodiment, both Cre-LoxP and Flp-Frt are used in the same system to regulate expression of the transgene, and for sequential deletion of vector sequences in the same cell, such as shown
25 in Example VIII.

The genetically modified animals can be used in genetic studies to further elucidate the p53 or p21 pathway, as animal models of disease and disorders implicating defective p53 or p21 function, and for *in vivo* testing of candidate therapeutic agents, such as those identified in the screens described below. Gene targeting in mice is an ideal method to
30 investigate the function of a distinct protein in wild type and disease states. Further, animal models deficient in rhomboid RRP sequence and function are desirable tools for modulating the EGFR signaling pathway, for testing the effect of candidate compounds against RRP, and for production of antibodies against human RRP, among others. The candidate therapeutic agents are administered to a genetically modified animal having

altered RRP function and phenotypic changes are compared with appropriate control animals such as genetically modified animals that receive placebo treatment, and/or animals with unaltered RRP expression that receive candidate therapeutic agent.

In additional to the above-described genetically modified animals having altered RRP function, animal models having defective p53 or p21 function (and otherwise normal RRP function), can be used in the methods of the present invention. For example, a p53 or p21 knockout mouse can be used to assess, *in vivo*, the activity of a candidate p53 or p21 modulatory agent identified in one of the *in vitro* assays described below. p53 or p21 knockout mice are described in the literature (p53: Jacks et al., Nature 2001;410:1111-1116, 1043-1044; Donehower *et al.*, supra; p21: Umanoff H, et al., Proc Natl Acad Sci U S A 1995 Feb 28;92(5):1709-13).

Modulating Agents

The invention provides methods to identify agents that interact with and/or modulate the function of RRP and/or the p53 or p21 pathway. Such agents are useful in a variety of diagnostic and therapeutic applications associated with the p53 or p21 pathways, as well as in further analysis of the RRP protein and its contribution to the p53 or p21 pathways. Accordingly, the invention also provides methods for modulating the p53 or p21 pathway comprising the step of specifically modulating RRP activity by administering an RRP-interacting or -modulating agent.

In a preferred embodiment, RRP-modulating agents inhibit or enhance RRP activity or otherwise affect normal RRP function, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In a further preferred embodiment, the candidate p53 or p21 pathway- modulating agent specifically modulates the function of the RRP. The phrases "specific modulating agent", "specifically modulates", etc., are used herein to refer to modulating agents that directly bind to the RRP polypeptide or nucleic acid, and preferably inhibit, enhance, or otherwise alter, the function of the RRP. The term also encompasses modulating agents that alter the interaction of the RRP with a binding partner or substrate (e.g. by binding to a binding partner of an RRP, or to a protein/binding partner complex, and inhibiting function).

Preferred RRP-modulating agents include small molecule compounds; RRP-interacting proteins, including antibodies and other biotherapeutics; antisense and RNA inhibitors. The modulating agents may be formulated in pharmaceutical compositions, for example, as compositions that may comprise other active ingredients, as in combination

therapy, and/or suitable carriers or excipients. Techniques for formulation and administration of the compounds may be found in "Remington's Pharmaceutical Sciences" Mack Publishing Co., Easton, PA, 19th edition.

5 **Small molecule modulators**

Small molecules, are often preferred to modulate function of proteins with enzymatic function, and/or containing protein interaction domains. Chemical agents, referred to in the art as "small molecule" compounds are typically organic, non-peptide molecules, having a molecular weight less than 10,000, preferably less than 5,000, more preferably less than 1,000, and most preferably less than 500. This class of modulators includes chemically synthesized molecules, for instance, compounds from combinatorial chemical libraries. Synthetic compounds may be rationally designed or identified based on known or inferred properties of the RRP protein or may be identified by screening compound libraries. Alternative appropriate modulators of this class are natural products, particularly secondary metabolites from organisms such as plants or fungi, which can also be identified by screening compound libraries for RRP-modulating activity. Methods for generating and obtaining compounds are well known in the art (Schreiber SL, Science (2000) 151: 1964-1969; Radmann J and Gunther J, Science (2000) 151:1947-1948).

Small molecule modulators identified from screening assays, as described below, can be used as lead compounds from which candidate clinical compounds may be designed, optimized, and synthesized. Such clinical compounds may have utility in treating pathologies associated with the p53 or p21 pathway. The activity of candidate small molecule modulating agents may be improved several-fold through iterative secondary functional validation, as further described below, structure determination, and candidate modulator modification and testing. Additionally, candidate clinical compounds are generated with specific regard to clinical and pharmacological properties. For example, the reagents may be derivatized and re-screened using *in vitro* and *in vivo* assays to optimize activity and minimize toxicity for pharmaceutical development.

30 **Protein Modulators**

Specific RRP-interacting proteins are useful in a variety of diagnostic and therapeutic applications related to the p53 or p21 pathway and related disorders, as well as in validation assays for other RRP-modulating agents. In a preferred embodiment, RRP-interacting proteins affect normal RRP function, including transcription, protein

expression, protein localization, and cellular or extra-cellular activity. In another embodiment, RRP-interacting proteins are useful in detecting and providing information about the function of RRP proteins, as is relevant to p53 or p21 related disorders, such as cancer (e.g., for diagnostic means).

5 An RRP-interacting protein may be endogenous, i.e. one that naturally interacts genetically or biochemically with an RRP, such as TGF α , EGF, amphiregulin, heregulin, a member of the RRP pathway that modulates RRP expression, localization, and/or activity. RRP-modulators include dominant negative forms of RRP-interacting proteins and of RRP proteins themselves. Yeast two-hybrid and variant screens offer preferred methods for
10 identifying endogenous RRP-interacting proteins (Finley, R. L. et al. (1996) in DNA Cloning-Expression Systems: A Practical Approach, eds. Glover D. & Hames B. D (Oxford University Press, Oxford, England), pp. 169-203; Fashema SF et al., Gene (2000) 250:1-14; Drees BL Curr Opin Chem Biol (1999) 3:64-70; Vidal M and Legrain P Nucleic Acids Res (1999) 27:919-29; and U.S. Pat. No. 5,928,868). Mass spectrometry is an
15 alternative preferred method for the elucidation of protein complexes (reviewed in, e.g., Pandley A and Mann M, Nature (2000) 405:837-846; Yates JR 3rd, Trends Genet (2000) 16:5-8).

 An RRP-interacting protein may be an exogenous protein, such as an RRP-specific antibody or a T-cell antigen receptor (see, e.g., Harlow and Lane (1988) Antibodies, A
20 Laboratory Manual, Cold Spring Harbor Laboratory). RRP antibodies are further discussed below.

 In preferred embodiments, an RRP-interacting protein specifically binds an RRP protein. In alternative preferred embodiments an RRP-modulating agent binds an RRP substrate, binding partner, or cofactor.

25

Antibodies

 In another embodiment, the protein modulator is an RRP specific antibody agonist or antagonist. The antibodies have therapeutic and diagnostic utilities, and can be used in screening assays to identify RRP modulators. For example, uses for antibodies include the
30 detection of an RRP protein in a biological sample and the inhibition of RRP activity, for instance, to block the development of an oncogenic disorder. The antibodies can also be used in dissecting the portions of the RRP pathway responsible for various cellular responses and in the general processing and maturation of the RRP.

Antibodies that specifically bind RRP polypeptides can be generated using known methods. Preferably the antibody is specific to a mammalian ortholog of RRP polypeptide, and more preferably, to human RRP. Antibodies may be polyclonal, monoclonal (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, fragments produced by a FAb expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. Monoclonal antibodies with affinities of 10^8 M^{-1} preferably 10^9 M^{-1} to 10^{10} M^{-1} , or stronger can be made by standard procedures as described (Harlow and Lane, Antibodies: A Laboratory Manual, CSH Laboratory (1988); Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed) Academic Press, New York; and U.S. Pat. Nos. 4,381,292; 4,451,570; and 4,618,577). Antibodies may be generated against crude cell extracts of RRP or substantially purified fragments thereof. If RRP fragments are used, they preferably comprise at least 10, and more preferably, at least 20 contiguous amino acids of an RRP protein. In a particular embodiment, RRP-specific antigens and/or immunogens are coupled to carrier proteins that stimulate the immune response. For example, the subject polypeptides are covalently coupled to the keyhole limpet hemocyanin (KLH) carrier, and the conjugate is emulsified in Freund's complete adjuvant, which enhances the immune response. An appropriate immune system such as a laboratory rabbit or mouse is immunized according to conventional protocols. In a preferred embodiment, due to close similarity of RRP sequences from mice and humans, transgenic mice that are RRP deficient or RRP knockout, such as those generated in the present invention (Example VIII), are used to produce antibodies against human RRP.

The presence of RRP-specific antibodies is assayed by an appropriate assay such as a solid phase enzyme-linked immunosorbant assay (ELISA) using immobilized corresponding RRP polypeptides. Other assays, such as radioimmunoassays or fluorescent assays might also be used.

Chimeric antibodies specific to RRP polypeptides can be made that contain different portions from different animal species. For instance, a human immunoglobulin constant region may be linked to a variable region of a murine mAb, such that the antibody derives its biological activity from the human antibody, and its binding specificity from the murine fragment. Chimeric antibodies are produced by splicing together genes that encode the appropriate regions from each species (Morrison et al., Proc. Natl. Acad. Sci. (1984) 81:6851-6855; Neuberger et al., Nature (1984) 312:604-608; Takeda et al., Nature (1985) 31:452-454). Humanized antibodies, which are a form of

chimeric antibodies, can be generated by grafting complementary-determining regions (CDRs) (Carlos, T. M., J. M. Harlan. 1994. Blood 84:2068-2101) of mouse antibodies into a background of human framework regions and constant regions by recombinant DNA technology (Riechmann LM, et al., 1988 Nature 323: 323-327). Humanized
5 antibodies contain ~10% murine sequences and ~90% human sequences, and thus further reduce or eliminate immunogenicity, while retaining the antibody specificities (Co MS, and Queen C. 1991 Nature 351: 501-501; Morrison SL. 1992 Ann. Rev. Immun. 10:239-265). Humanized antibodies and methods of their production are well-known in the art (U.S. Pat. No. 5,530,101; U.S. Pat. No. 5,585,089; U.S. Pat. No. 5,693,762, and
10 U.S. Pat. No. 6,180,370).

RRP-specific single chain antibodies which are recombinant, single chain polypeptides formed by linking the heavy and light chain fragments of the Fv regions via an amino acid bridge, can be produced by methods known in the art (U.S. Pat. No. 4,946,778; Bird, Science (1988) 242:423-426; Huston et al., Proc. Natl. Acad. Sci. USA
15 (1988) 85:5879-5883; and Ward et al., Nature (1989) 334:544-546).

Other suitable techniques for antibody production involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors (Huse et al., Science (1989) 246:1275-1281). As used herein, T-cell antigen receptors are included within the scope of antibody modulators
20 (Harlow and Lane, 1988, *supra*).

The polypeptides and antibodies of the present invention may be used with or without modification. Frequently, antibodies will be labeled by joining, either covalently or non-covalently, a substance that provides for a detectable signal, or that is toxic to cells that express the targeted protein (Menard S, et al., Int J. Biol Markers (1989) 4:131-134).
25 A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, fluorescent emitting lanthanide metals, chemiluminescent moieties, bioluminescent moieties, magnetic particles, and the like (U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350;
30 3,996,345; 4,277,437; 4,275,149; and 4,366,241). Also, recombinant immunoglobulins may be produced (U.S. Pat. No. 4,816,567). Antibodies to cytoplasmic polypeptides may be delivered and reach their targets by conjugation with membrane-penetrating toxin proteins (U.S. Pat. No. 6,086,900).

When used therapeutically in a patient, the antibodies of the subject invention are typically administered parenterally, when possible at the target site, or intravenously. The therapeutically effective dose and dosage regimen is determined by clinical studies.

Typically, the amount of antibody administered is in the range of about 0.1 mg/kg –to
5 about 10 mg/kg of patient weight. For parenteral administration, the antibodies are formulated in a unit dosage injectable form (e.g., solution, suspension, emulsion) in association with a pharmaceutically acceptable vehicle. Such vehicles are inherently nontoxic and non-therapeutic. Examples are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Nonaqueous vehicles such as fixed oils, ethyl
10 oleate, or liposome carriers may also be used. The vehicle may contain minor amounts of additives, such as buffers and preservatives, which enhance isotonicity and chemical stability or otherwise enhance therapeutic potential. The antibodies' concentrations in such vehicles are typically in the range of about 1 mg/ml–to about 10 mg/ml. Immunotherapeutic methods are further described in the literature (US Pat. No. 5,859,206;
15 WO0073469).

The selection of an appropriate antibody subclass for therapy will depend upon the nature of the tumor antigen. For example, an IgM may be preferred when the antigen is highly specific for the tumor target and rarely occurs on normal cells. However, the IgG subclass may be preferred when the tumor-associated antigen is also expressed in normal
20 tissues, even at much lower levels. The binding of at least two IgG molecules in close proximity is required to activate complement, a serum protein that combines with antibodies to form a defense against cellular antigens. The normal tissues that express smaller amounts of the antigen and bind fewer IgG molecules may thus incur less complement-mediated damage. Furthermore, since IgGs are smaller than IgMs, they may
25 more readily localize to tumor tissue.

Immune responses may assist in the delivery or efficacy of an anti-tumor treatment. There is evidence that complement activation leads to an inflammatory response and macrophage activation (Unanue and Benecerraf, Textbook of Immunology, 2nd Edition, Williams & Wilkins, p. 218 (1984)). Activated macrophages more
30 preferentially destroy tumor cells than normal cells (Fidler and Poste, Springer Semin. Immunopathol. 5, 161 (1982)). Also, the increased vasodilation accompanying inflammation may increase the ability of anti-cancer agents, such as chemotherapeutic drugs or radiolabeled antibodies to localize in tumors. While a significant detriment of standard chemotherapy or radiation treatment is damage to healthy cells, the antigen-

antibody combinations specified by this invention may circumvent many of the problems normally caused by the heterogeneity of tumor cell populations. Additionally, purified antigens (Hakomori, Ann. Rev. Immunol. (1984) 2:103) or the related anti-idiotypic antibodies (Nepom et al., Proc. Natl. Acad. Sci. (1985) 81:2864; Koprowski et al., Proc. Natl. Acad. Sci. (1984) 81:216) which recognize the hypervariance among the same epitopes in different individuals could be used to induce an active immune response in human cancer patients. Such a response includes the formation of antibodies capable of activating human complement and mediating antibody-dependent cell-mediated cytotoxicity and by such mechanisms cause tumor destruction.

Specific biotherapeutics

In a preferred embodiment, an RRP-interacting protein may have biotherapeutic applications. Biotherapeutic agents formulated in pharmaceutically acceptable carriers and dosages may be used to activate or inhibit signal transduction pathways. This modulation may be accomplished by binding a ligand, thus inhibiting the activity of the pathway; or by binding a receptor, either to inhibit activation of, or to activate, the receptor. Alternatively, the biotherapeutic may itself be a ligand capable of activating or inhibiting a receptor. Biotherapeutic agents and methods of producing them are described in detail in U.S. Pat. No. 6,146,628.

Since RRP is a receptor, its ligand(s), antibodies to the ligand(s) or the RRP itself may be used as biotherapeutics to modulate the activity of RRP in the p53 or p21 pathway.

Nucleic Acid Modulators

Other preferred RRP-modulating agents comprise nucleic acid molecules, such as antisense oligomers or double stranded RNA (dsRNA), which generally inhibit RRP activity.

Preferred antisense oligomers that interfere with the function of the RRP nucleic acid such as DNA replication, transcription, translocation of the RRP RNA to the site of protein translation, translation of protein from the RRP RNA, splicing of the RRP RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RRP RNA. Double-stranded RNA inhibition (dsRNAi) is another preferred RRP-modulating agent. For convenience, the term "antisense modulator", as used herein, includes antisense oligomers and dsRNAi.

In one embodiment, the antisense oligomer is an oligonucleotide that is sufficiently complementary to an RRP mRNA to bind to and prevent translation, preferably by binding to the 5' untranslated region. RRP-specific antisense oligonucleotides, preferably range from at least 6 to about 200 nucleotides. In some embodiments the oligonucleotide is preferably at least 10, 15, or 20 nucleotides in length. In other embodiments, the oligonucleotide is preferably less than 50, 40, or 30 nucleotides in length. The oligonucleotide can be DNA or RNA or a chimeric mixture or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone. The oligonucleotide may include other appending groups such as peptides, agents that facilitate transport across the cell membrane, hybridization-triggered cleavage agents, and intercalating agents.

In another embodiment, the antisense oligomer is a phosphothioate morpholino oligomer (PMO). PMOs are assembled from four different morpholino subunits, each of which contain one of four genetic bases (A, C, G, or T) linked to a six-membered morpholine ring. Polymers of these subunits are joined by non-ionic phosphodiarnidate intersubunit linkages. Details of how to make and use PMOs and other antisense oligomers are well known in the art (e.g. see WO99/18193; Probst JC, Antisense Oligodeoxynucleotide and Ribozyme Design, Methods. (2000) 22(3):271-281; Summerton J, and Weller D. 1997 Antisense Nucleic Acid Drug Dev. ;7:187-95, US Pat. No. 5,235,033; and US Pat No. 5,378,841).

Antisense oligomers are commonly used as research reagents, diagnostics, and therapeutics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used to elucidate the function of particular genes (see, for example, U.S. Pat. No. 6,165,790). Antisense oligomers are also used, for example, to distinguish between functions of various members of a biological pathway. Antisense oligomers have been employed as therapeutic moieties in the treatment of disease states in animals and man and have been demonstrated in numerous clinical trials to be safe and effective (Milligan JF, *et al*, Current Concepts in Antisense Drug Design, J Med Chem. (1993) 36:1923-1937; Tonkinson JL *et al.*, Antisense Oligodeoxynucleotides as Clinical Therapeutic Agents, Cancer Invest. (1996) 14:54-65). Accordingly, in one aspect of the invention, an RRP-specific antisense oligomer is used in an assay to further elucidate the role of the RRP in the p53 or p21 pathway, and/or its relationship to other members of the pathway. In another aspect of the invention, an RRP-specific antisense oligomer is used as a therapeutic agent for treatment of p53 or p21-related disease states.

Alternative preferred RRP-modulating agents are double-stranded RNA species mediating RNA interference (RNAi). RNAi is the process of sequence-specific, post-transcriptional gene silencing in animals and plants, initiated by double-stranded RNA (dsRNA) that is homologous in sequence to the silenced gene. Methods relating to the use of RNAi to silence genes in *C. elegans*, *Drosophila*, plants, and humans are known in the art (Fire A, et al., 1998 Nature 391:806-811; Fire, A. Trends Genet. 15, 358-363 (1999); Sharp, P. A. RNA interference 2001. Genes Dev. 15, 485-490 (2001); Hammond, S. M., et al., Nature Rev. Genet. 2, 110-1119 (2001); Tuschl, T. Chem. Biochem. 2, 239-245 (2001); Hamilton, A. et al., Science 286, 950-952 (1999); Hammond, S. M., et al., Nature 404, 293-296 (2000); Zamore, P. D., et al., Cell 101, 25-33 (2000); Bernstein, E., et al., Nature 409, 363-366 (2001); Elbashir, S. M., et al., Genes Dev. 15, 188-200 (2001); WO0129058; WO9932619; Elbashir SM, et al., 2001 Nature 411:494-498).

Assay Systems

The invention provides assay systems for identifying specific modulators of RRP activity. As used herein, an "assay system" encompasses all the components required for performing and analyzing results of an assay that detects and/or measures a particular event. In general, primary assays are used to identify or confirm a modulator's specific biochemical or molecular effect with respect to the RRP nucleic acid or protein. In general, secondary assays further assess the activity of a RRP modulating agent identified by a primary assay and may confirm that the modulating agent affects RRP in a manner relevant to the p53 or p21 pathway. In some cases, RRP modulators will be directly tested in a secondary assay.

In a preferred embodiment, the screening method comprises contacting a suitable assay system comprising an RRP polypeptide with a candidate agent under conditions whereby, but for the presence of the agent, the system provides a reference activity (e.g. protease activity), which is based on the particular molecular event the screening method detects. A statistically significant difference between the agent-biased activity and the reference activity indicates that the candidate agent modulates RRP activity, and hence the p53 or p21 pathway.

Primary Assays

The type of modulator tested generally determines the type of primary assay.

Primary assays for small molecule modulators

For small molecule modulators, screening assays are used to identify candidate modulators. Screening assays may be cell-based or may use a cell-free system that recreates or retains the relevant biochemical reaction of the target protein (reviewed in Sittampalam GS *et al.*, Curr Opin Chem Biol (1997) 1:384-91 and accompanying references). As used herein the term "cell-based" refers to assays using live cells, dead cells, or a particular cellular fraction, such as a membrane, endoplasmic reticulum, or mitochondrial fraction. The term "cell free" encompasses assays using substantially purified protein (either endogenous or recombinantly produced), partially purified cellular extracts, or crude cellular extracts. Screening assays may detect a variety of molecular events, including protein-DNA interactions, protein-protein interactions (*e.g.*, receptor-ligand binding), transcriptional activity (*e.g.*, using a reporter gene), enzymatic activity (*e.g.*, via a property of the substrate), activity of second messengers, immunogenicity and changes in cellular morphology or other cellular characteristics. Appropriate screening assays may use a wide range of detection methods including fluorescent, radioactive, colorimetric, spectrophotometric, and amperometric methods, to provide a read-out for the particular molecular event detected.

In a preferred embodiment, screening assays uses fluorescence technologies, including fluorescence polarization, time-resolved fluorescence, and fluorescence resonance energy transfer. These systems offer means to monitor protein-protein or DNA-protein interactions in which the intensity of the signal emitted from dye-labeled molecules depends upon their interactions with partner molecules (*e.g.*, Selvin PR, Nat Struct Biol (2000) 7:730-4; Fernandes PB, Curr Opin Chem Biol (1998) 2:597-603; Hertzberg RP and Pope AJ, Curr Opin Chem Biol (2000) 4:445-451).

Cell-based screening assays usually require systems for recombinant expression of RRP and any auxiliary proteins demanded by the particular assay. Cell-free assays often use recombinantly produced purified or substantially purified proteins. Appropriate methods for generating recombinant proteins produce sufficient quantities of proteins that retain their relevant biological activities and are of sufficient purity to optimize activity and assure assay reproducibility. Yeast two-hybrid and variant screens, and mass spectrometry provide preferred methods for determining protein-protein interactions and elucidation of protein complexes. In certain applications, when RRP-interacting proteins are used in screens to identify small molecule modulators, the binding specificity of the interacting protein to the RRP protein may be assayed by various known methods such as

substrate processing (e.g. ability of the candidate RRP-specific binding agents to function as negative effectors in RRP-expressing cells), binding equilibrium constants (usually at least about 10^7 M^{-1} , preferably at least about 10^8 M^{-1} , more preferably at least about 10^9 M^{-1}), and immunogenicity (e.g. ability to elicit RRP specific antibody in a heterologous host such as a mouse, rat, goat or rabbit). For enzymes and receptors, binding may be assayed by, respectively, substrate and ligand processing.

The screening assay may measure a candidate agent's ability to specifically bind to or modulate activity of a RRP polypeptide, a fusion protein thereof, or to cells or membranes bearing the polypeptide or fusion protein. The RRP polypeptide can be full length or a fragment thereof that retains functional RRP activity. The RRP polypeptide may be fused to another polypeptide, such as a peptide tag for detection or anchoring, or to another tag. The RRP polypeptide is preferably human or mouse RRP, or is an ortholog or derivative thereof as described above. In a preferred embodiment, the screening assay detects candidate agent-based modulation of RRP interaction with a binding target, such as an endogenous or exogenous protein or other substrate that has RRP-specific binding activity, and can be used to assess normal RRP gene function.

Suitable assay formats that may be adapted to screen for RRP modulators are known in the art. Preferred screening assays are high throughput or ultra high throughput and thus provide automated, cost-effective means of screening compound libraries for lead compounds (Fernandes PB, 1998, *supra*; Sundberg SA, Curr Opin Biotechnol 2000, 11:47-53).

A variety of suitable assay systems may be used to identify candidate RRP and p53 or p21 pathway modulators (e.g. U.S. Pat. No. 6,020,135 (p53 modulation), U.S. Pat. No. 6,114,132 (phosphatase and protease assays)). Specific preferred assays are described in more detail below.

Protease Assays. Proteases are enzymes that cleave protein substrates at specific sites. Exemplary assays detect the alterations in the spectral properties of an artificial substrate that occur upon protease-mediated cleavage. In one example, synthetic caspase substrates containing four amino acid proteolysis recognition sequences, separating two different fluorescent tags are employed; fluorescence resonance energy transfer detects the proximity of these fluorophores, which indicates whether the substrate is cleaved (Mahajan NP *et al.*, Chem Biol (1999) 6:401-409).

Endogenous protease inhibitors may inhibit protease activity. In an example of an assay developed for either proteases or protease inhibitors, a biotinylated substrate is coated on a titer plate and hydrolyzed with the protease; the unhydrolyzed substrate is quantified by reaction with alkaline phosphatase-streptavidin complex and detection of the reaction product. The activity of protease inhibitors correlates with the activity of the alkaline phosphatase indicator enzyme (Gan Z *et al.*, Anal Biochem 1999) 268:151-156).

Apoptosis assays. Assays for apoptosis may be performed by terminal deoxynucleotidyl transferase-mediated digoxigenin-11-dUTP nick end labeling (TUNEL) assay. The TUNEL assay is used to measure nuclear DNA fragmentation characteristic of apoptosis (Lazebnik *et al.*, 1994, Nature 371, 346), by following the incorporation of fluorescein-dUTP (Yonehara *et al.*, 1989, J. Exp. Med. 169, 1747). Apoptosis may further be assayed by acridine orange staining of tissue culture cells (Lucas, R., et al., 1998, Blood 15:4730-41). An apoptosis assay system may comprise a cell that expresses an RRP, and that optionally has defective p53 or p21 function (e.g. p53 or p21 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the apoptosis assay system and changes in induction of apoptosis relative to controls where no test agent is added, identify candidate p53 or p21 modulating agents. In some embodiments of the invention, an apoptosis assay may be used as a secondary assay to test a candidate p53 or p21 modulating agents that is initially identified using a cell-free assay system. An apoptosis assay may also be used to test whether RRP function plays a direct role in apoptosis. For example, an apoptosis assay may be performed on cells that over- or under-express RRP relative to wild type cells. Differences in apoptotic response compared to wild type cells suggests that the RRP plays a direct role in the apoptotic response. Apoptosis assays are described further in US Pat. No. 6,133,437.

Cell proliferation and cell cycle assays. Cell proliferation may be assayed via bromodeoxyuridine (BRDU) incorporation. This assay identifies a cell population undergoing DNA synthesis by incorporation of BRDU into newly-synthesized DNA. Newly-synthesized DNA may then be detected using an anti-BRDU antibody (Hoshino *et al.*, 1986, Int. J. Cancer 38, 369; Campana *et al.*, 1988, J. Immunol. Meth. 107, 79), or by other means.

Cell Proliferation may also be examined using [³H]-thymidine incorporation (Chen, J., 1996, Oncogene 13:1395-403; Jeoung, J., 1995, J. Biol. Chem. 270:18367-73).

This assay allows for quantitative characterization of S-phase DNA syntheses. In this assay, cells synthesizing DNA will incorporate [³H]-thymidine into newly synthesized DNA. Incorporation can then be measured by standard techniques such as by counting of radioisotope in a scintillation counter (e.g., Beckman LS 3800 Liquid Scintillation Counter).

Cell proliferation may also be assayed by colony formation in soft agar (Sambrook et al., Molecular Cloning, Cold Spring Harbor (1989)). For example, cells transformed with RRP are seeded in soft agar plates, and colonies are measured and counted after two weeks incubation.

Involvement of a gene in the cell cycle may be assayed by flow cytometry. Cells transfected with an RRP may be stained with propidium iodide and evaluated in a flow cytometer (available from Becton Dickinson).

Accordingly, a cell proliferation or cell cycle assay system may comprise a cell that expresses an RRP, and that optionally has defective p53 or p21 function (e.g. p53 or p21 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the assay system and changes in cell proliferation or cell cycle relative to controls where no test agent is added, identify candidate p53 or p21 modulating agents. In some embodiments of the invention, the cell proliferation or cell cycle assay may be used as a secondary assay to test a candidate p53 or p21 modulating agents that is initially identified using another assay system such as a cell-free kinase assay system. A cell proliferation assay may also be used to test whether RRP function plays a direct role in cell proliferation or cell cycle. For example, a cell proliferation or cell cycle assay may be performed on cells that over- or under-express RRP relative to wild type cells.

Differences in proliferation or cell cycle compared to wild type cells suggests that the RRP plays a direct role in cell proliferation or cell cycle. A cell proliferation assay may also be used to identify candidate agents that modulate cell proliferation. For example, cells that have decreased expression of RRP or that are RRP knockouts, such as mouse cells generated in the present invention (Example VIII) are treated with candidate agents.

Changes in cell proliferation relative to control cells where no agent is added indicate that the candidate agent modulates cell proliferation.

Angiogenesis. Angiogenesis may be assayed using various human endothelial cell systems, such as umbilical vein, coronary artery, or dermal cells. Suitable assays include Alamar Blue based assays (available from Biosource International) to measure

proliferation; migration assays using fluorescent molecules, such as the use of Becton Dickinson Falcon HTS FluoroBlock cell culture inserts to measure migration of cells through membranes in presence or absence of angiogenesis enhancer or suppressors; and tubule formation assays based on the formation of tubular structures by endothelial cells on Matrigel® (Becton Dickinson). Accordingly, an angiogenesis assay system may comprise a cell that expresses an RRP, and that optionally has defective p53 or p21 function (e.g. p53 or p21 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the angiogenesis assay system and changes in angiogenesis relative to controls where no test agent is added, identify candidate p53 or p21 modulating agents. In some embodiments of the invention, the angiogenesis assay may be used as a secondary assay to test a candidate p53 or p21 modulating agents that is initially identified using another assay system. An angiogenesis assay may also be used to test whether RRP function plays a direct role in cell proliferation. For example, an angiogenesis assay may be performed on cells that over- or under-express RRP relative to wild type cells. Differences in angiogenesis compared to wild type cells suggests that the RRP plays a direct role in angiogenesis.

Hypoxic induction. The alpha subunit of the transcription factor, hypoxia inducible factor-1 (HIF-1), is upregulated in tumor cells following exposure to hypoxia in vitro. Under hypoxic conditions, HIF-1 stimulates the expression of genes known to be important in tumor cell survival, such as those encoding glycolytic enzymes and VEGF. Induction of such genes by hypoxic conditions may be assayed by growing cells transfected with RRP in hypoxic conditions (such as with 0.1% O₂, 5% CO₂, and balance N₂, generated in a Napco 7001 incubator (Precision Scientific)) and normoxic conditions, followed by assessment of gene activity or expression by Taqman®. For example, a hypoxic induction assay system may comprise a cell that expresses an RRP, and that optionally has a mutated p53 or p21 (e.g. p53 or p21 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the hypoxic induction assay system and changes in hypoxic response relative to controls where no test agent is added, identify candidate p53 or p21 modulating agents. In some embodiments of the invention, the hypoxic induction assay may be used as a secondary assay to test a candidate p53 or p21 modulating agents that is initially identified using another assay system. A hypoxic induction assay may also be used to test whether RRP function plays a direct role in the hypoxic response. For example, a hypoxic induction assay may be performed on cells that

over- or under-express RRP relative to wild type cells. Differences in hypoxic response compared to wild type cells suggest that the RRP plays a direct role in hypoxic induction.

Cell adhesion. Cell adhesion assays measure adhesion of cells to purified
5 adhesion proteins, or adhesion of cells to each other, in presence or absence of candidate
modulating agents.

Cell-protein adhesion assays measure the ability of agents to modulate the
adhesion of cells to purified proteins. For example, recombinant proteins are produced,
diluted to 2.5g/mL in PBS, and used to coat the wells of a microtiter plate. The wells used
10 for negative control are not coated. Coated wells are then washed, blocked with 1% BSA,
and washed again. Compounds are diluted to 2× final test concentration and added to the
blocked, coated wells. Cells are then added to the wells, and the unbound cells are washed
off. Retained cells are labeled directly on the plate by adding a membrane-permeable
fluorescent dye, such as calcein-AM, and the signal is quantified in a fluorescent
15 microplate reader.

Cell-cell adhesion assays measure the ability of agents to modulate binding of cell
adhesion proteins with their native ligands. These assays use cells that naturally or
recombinantly express the adhesion protein of choice. In an exemplary assay, cells
expressing the cell adhesion protein are plated in wells of a multiwell plate. Cells
20 expressing the ligand are labeled with a membrane-permeable fluorescent dye, such as
BCECF, and allowed to adhere to the monolayers in the presence of candidate agents.
Unbound cells are washed off, and bound cells are detected using a fluorescence plate
reader.

High-throughput cell adhesion assays have also been described. In one such assay,
25 small molecule ligands and peptides are bound to the surface of microscope slides using a
microarray spotter, intact cells are then contacted with the slides, and unbound cells are
washed off. In this assay, not only the binding specificity of the peptides and modulators
against cell lines are determined, but also the functional cell signaling of attached cells
using immunofluorescence techniques in situ on the microchip is measured (Falsey JR et
30 al., Bioconj Chem. 2001 May-Jun;12(3):346-53).

Certain screening assays may also be used to test antibody and nucleic acid
modulators; for nucleic acid modulators, appropriate assay systems involve RRP mRNA
expression.

Primary assays for antibody modulators

For antibody modulators, appropriate primary assays test is a binding assay that tests the antibody's affinity to and specificity for the RRP protein. Methods for testing antibody affinity and specificity are well known in the art (Harlow and Lane, 1988, 1999, *supra*). The enzyme-linked immunosorbant assay (ELISA) is a preferred method for detecting RRP-specific antibodies; others include FACS assays, radioimmunoassays, and fluorescent assays.

Primary assays for nucleic acid modulators

For nucleic acid modulators, primary assays may test the ability of the nucleic acid modulator to inhibit or enhance RRP gene expression, preferably mRNA expression. In general, expression analysis comprises comparing RRP expression in like populations of cells (*e.g.*, two pools of cells that endogenously or recombinantly express RRP) in the presence and absence of the nucleic acid modulator. Methods for analyzing mRNA and protein expression are well known in the art. For instance, Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR (*e.g.*, using the TaqMan®, PE Applied Biosystems), or microarray analysis may be used to confirm that RRP mRNA expression is reduced in cells treated with the nucleic acid modulator (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann Med 2001, 33:142-147; Blohm DH and Guiseppe-Elie, A Curr Opin Biotechnol 2001, 12:41-47). Protein expression may also be monitored. Proteins are most commonly detected with specific antibodies or antisera directed against either the RRP protein or specific peptides. A variety of means including Western blotting, ELISA, or in situ detection, are available (Harlow E and Lane D, 1988 and 1999, *supra*).

Secondary Assays

Secondary assays may be used to further assess the activity of RRP-modulating agent identified by any of the above methods to confirm that the modulating agent affects RRP in a manner relevant to the p53 or p21 pathway. As used herein, RRP-modulating agents encompass candidate clinical compounds or other agents derived from previously identified modulating agent. Secondary assays can also be used to test the activity of a modulator on a particular genetic or biochemical pathway or to test the specificity of the modulator's interaction with RRP.

Secondary assays generally compare like populations of cells or animals (*e.g.*, two pools of cells or animals that endogenously or recombinantly express RRP) in the presence and absence of the candidate modulator. In general, such assays test whether treatment of cells or animals with a candidate RRP-modulating agent results in changes in the p53 or p21 pathway in comparison to untreated (or mock- or placebo-treated) cells or animals. Certain assays use “sensitized genetic backgrounds”, which, as used herein, describe cells or animals engineered for altered expression of genes in the p53 or p21 or interacting pathways.

10 *Cell-based assays*

Cell based assays may use a variety of mammalian cell lines known to have defective p53 or p21 function (*e.g.* for p53: SAOS-2 osteoblasts, H1299 lung cancer cells, C33A and HT3 cervical cancer cells, HT-29 and DLD-1 colon cancer cells, among others, and for p21: HCT116 colon cancer cells, among others, available from American Type Culture Collection (ATCC), Manassas, VA). Cell based assays may detect endogenous p53 or p21 pathway activity or may rely on recombinant expression of p53 or p21 pathway components. Any of the aforementioned assays may be used in this cell-based format. Candidate modulators are typically added to the cell media but may also be injected into cells or delivered by any other efficacious means.

20

Animal Assays

A variety of non-human animal models of normal or defective p53 or p21 pathway may be used to test candidate RRP modulators. Models for defective p53 or p21 pathway typically use genetically modified animals that have been engineered to mis-express (*e.g.*, over-express or lack expression in) genes involved in the p53 or p21 pathway. Assays generally require systemic delivery of the candidate modulators, such as by oral administration, injection, etc.

In a preferred embodiment, p53 or p21 pathway activity is assessed by monitoring neovascularization and angiogenesis. Animal models with defective and normal p53 or p21 are used to test the candidate modulator’s affect on RRP in Matrigel® assays. Matrigel® is an extract of basement membrane proteins, and is composed primarily of laminin, collagen IV, and heparin sulfate proteoglycan. It is provided as a sterile liquid at 4° C, but rapidly forms a solid gel at 37° C. Liquid Matrigel® is mixed with various angiogenic agents, such as bFGF and VEGF, or with human tumor cells which over-

express the RRP. The mixture is then injected subcutaneously into the female athymic nude mice (Taconic, Germantown, NY) to support an intense vascular response. Mice with Matrigel® pellets may be dosed via oral (PO), intraperitoneal (IP), or intravenous (IV) routes with the candidate modulator. Mice are euthanized 5 - 12 days post-injection, and the Matrigel® pellet is harvested for hemoglobin analysis (Sigma plasma hemoglobin kit). Hemoglobin content of the gel is found to correlate the degree of neovascularization in the gel.

In another preferred embodiment, the effect of the candidate modulator on RRP is assessed via tumorigenicity assays. In one example, xenograft human tumors are implanted subcutaneously (SC) into female athymic nude mice, 6-7 week old, as single cell suspensions either from a pre-existing tumor or from *in vitro* culture. The tumors which express the RRP endogenously are injected in the flank, 1×10^5 to 1×10^7 cells per mouse in a volume of 100 μ L using a 27gauge needle. Mice are then ear tagged and tumors are measured twice weekly. Candidate modulator treatment is initiated on the day the mean tumor weight reaches 100 mg. Candidate modulator is delivered intravenously (IV), subcutaneously (SC), intraperitoneously (IP), or orally (PO) by bolus administration. Depending upon the pharmacokinetics (PK) of each unique candidate modulator, dosing can be performed multiple times per day. The tumor weight is assessed by measuring perpendicular diameters with a caliper and calculated by multiplying the measurements of diameters in two dimensions. At the end of the experiment, the excised tumors may be utilized for biomarker identification or further analyses. For immunohistochemistry staining, xenograft tumors are fixed in 4% paraformaldehyde, 0.1M phosphate, PH 7.2, for 6 hours at 4°C, immersed in 30% sucrose in PBS, and rapidly frozen in isopentane cooled with liquid nitrogen.

Diagnostic and therapeutic uses

Specific RRP-modulating agents are useful in a variety of diagnostic and therapeutic applications where disease or disease prognosis is related to defects in the p53 or p21 pathway, such as angiogenic, apoptotic, or cell proliferation disorders. Accordingly, the invention also provides methods for modulating the p53 or p21 pathway in a cell, preferably a cell pre-determined to have defective p53 or p21 function, comprising the step of administering an agent to the cell that specifically modulates RRP activity.

The discovery that RRP is implicated in p53 or p21 pathway provides for a variety of methods that can be employed for the diagnostic and prognostic evaluation of diseases and disorders involving defects in the p53 or p21 pathway and for the identification of subjects having a predisposition to such diseases and disorders.

5 Various expression analysis methods can be used to diagnose whether RRP expression occurs in a particular sample, including Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR, and microarray analysis. (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann
10 Med 2001, 33:142-147; Blohm and Guiseppi-Elie, Curr Opin Biotechnol 2001, 12:41-47). Tissues having a disease or disorder implicating defective p53 or p21 signaling that express an RRP, are identified as amenable to treatment with an RRP modulating agent. In a preferred application, the p53 or p21 defective tissue overexpresses an RRP relative to normal tissue. For example, a Northern blot analysis of mRNA from tumor and normal
15 cell lines, or from tumor and matching normal tissue samples from the same patient, using full or partial RRP cDNA sequences as probes, can determine whether particular tumors express or overexpress RRP. Alternatively, the TaqMan® is used for quantitative RT-PCR analysis of RRP expression in cell lines, normal tissues and tumor samples (PE Applied Biosystems).

20 Various other diagnostic methods may be performed, for example, utilizing reagents such as the RRP oligonucleotides, and antibodies directed against an RRP, as described above for: (1) the detection of the presence of RRP gene mutations, or the detection of either over- or under-expression of RRP mRNA relative to the non-disorder state; (2) the detection of either an over- or an under-abundance of RRP gene product
25 relative to the non-disorder state; and (3) the detection of perturbations or abnormalities in the signal transduction pathway mediated by RRP.

Thus, in a specific embodiment, the invention is drawn to a method for diagnosing a disease in a patient, the method comprising: a) obtaining a biological sample from the patient; b) contacting the sample with a probe for RRP expression; c) comparing results
30 from step (b) with a control; and d) determining whether step (c) indicates a likelihood of disease. Preferably, the disease is cancer, most preferably a cancer as shown in TABLE 1. The probe may be either DNA or protein, including an antibody.

EXAMPLES

The following experimental section and examples are offered by way of illustration and not by way of limitation.

5 I. Drosophila p53 and p21 screens

The *Drosophila* p53 gene was overexpressed specifically in the wing using the vestigial margin quadrant enhancer. Increasing quantities of *Drosophila* p53 (titrated using different strength transgenic inserts in 1 or 2 copies) caused deterioration of normal wing morphology from mild to strong, with phenotypes including disruption of pattern and polarity of wing hairs, shortening and thickening of wing veins, progressive crumpling of the wing and appearance of dark "death" inclusions in wing blade. In a screen designed to identify enhancers and suppressors of *Drosophila* p53, homozygous females carrying two copies of p53 were crossed to 5663 males carrying random insertions of a piggyBac transposon (Fraser M *et al.*, Virology (1985) 145:356-361). Progeny containing insertions were compared to non-insertion-bearing sibling progeny for enhancement or suppression of the p53 phenotypes. Sequence information surrounding the piggyBac insertion site was used to identify the modifier genes. Modifiers of the wing phenotype were identified as members of the p53 pathway.

An overexpression screen was carried out in *Drosophila* to identify genes that interact with the cyclin dependent kinase inhibitor, p21 (Bourne HR, et al., Nature (1990) 348(6297):125-132; Marshall CJ, Trends Genet (1991) 7(3):91-95). Expression of the p21 gene in the eye causes deterioration of normal eye morphology. Modifiers of the eye phenotype were identified as members of the p21 pathway.

25 II. High-Throughput In Vitro Fluorescence Polarization Assay

Fluorescently-labeled RRP peptide/substrate are added to each well of a 96-well microtiter plate, along with a test agent in a test buffer (10 mM HEPES, 10 mM NaCl, 6 mM magnesium chloride, pH 7.6). Changes in fluorescence polarization, determined by using a Fluorolite FPM-2 Fluorescence Polarization Microtiter System (Dynatech Laboratories, Inc), relative to control values indicates the test compound is a candidate modifier of RRP activity.

III. High-Throughput In Vitro Binding Assay.

³³P-labeled RRP peptide is added in an assay buffer (100 mM KCl, 20 mM HEPES pH 7.6, 1 mM MgCl₂, 1% glycerol, 0.5% NP-40, 50 mM beta-mercaptoethanol, 1 mg/ml BSA, cocktail of protease inhibitors) along with a test agent to the wells of a Neutralite-
5 avidin coated assay plate and incubated at 25°C for 1 hour. Biotinylated substrate is then added to each well and incubated for 1 hour. Reactions are stopped by washing with PBS, and counted in a scintillation counter. Test agents that cause a difference in activity relative to control without test agent are identified as candidate p53 or p21 modulating agents.

10

IV. Immunoprecipitations and Immunoblotting

For coprecipitation of transfected proteins, 3×10^6 appropriate recombinant cells containing the RRP proteins are plated on 10-cm dishes and transfected on the following day with expression constructs. The total amount of DNA is kept constant in each
15 transfection by adding empty vector. After 24 h, cells are collected, washed once with phosphate-buffered saline and lysed for 20 min on ice in 1 ml of lysis buffer containing 50 mM Hepes, pH 7.9, 250 mM NaCl, 20 mM -glycerophosphate, 1 mM sodium orthovanadate, 5 mM p-nitrophenyl phosphate, 2 mM dithiothreitol, protease inhibitors (complete, Roche Molecular Biochemicals), and 1% Nonidet P-40. Cellular debris is
20 removed by centrifugation twice at $15,000 \times g$ for 15 min. The cell lysate is incubated with 25 μ l of M2 beads (Sigma) for 2 h at 4 °C with gentle rocking. After extensive washing with lysis buffer, proteins bound to the beads are solubilized by boiling in SDS sample buffer, fractionated by SDS-polyacrylamide gel electrophoresis, transferred to polyvinylidene difluoride membrane and blotted with the indicated
25 antibodies. The reactive bands are visualized with horseradish peroxidase coupled to the appropriate secondary antibodies and the enhanced chemiluminescence (ECL) Western blotting detection system (Amersham Pharmacia Biotech).

V. Expression analysis

30

All cell lines used in the following experiments are NCI (National Cancer Institute) lines, and are available from ATCC (American Type Culture Collection, Manassas, VA 20110-2209). Normal and tumor tissues were obtained from Impath, UC Davis, Clontech, Stratagene, and Ambion.

TaqMan analysis was used to assess expression levels of the disclosed genes in various samples.

RNA was extracted from each tissue sample using Qiagen (Valencia, CA) RNeasy kits, following manufacturer's protocols, to a final concentration of 50ng/μl. Single
5 stranded cDNA was then synthesized by reverse transcribing the RNA samples using random hexamers and 500ng of total RNA per reaction, following protocol 4304965 of Applied Biosystems (Foster City, CA, website at www.appliedbiosystems.com).

Primers for expression analysis using TaqMan assay (Applied Biosystems, Foster City, CA) were prepared according to the TaqMan protocols, and the following criteria: a)
10 primer pairs were designed to span introns to eliminate genomic contamination, and b) each primer pair produced only one product.

Taqman reactions were carried out following manufacturer's protocols, in 25 μl total volume for 96-well plates and 10 μl total volume for 384-well plates, using 300nM primer and 250 nM probe, and approximately 25ng of cDNA. The standard curve for
15 result analysis was prepared using a universal pool of human cDNA samples, which is a mixture of cDNAs from a wide variety of tissues so that the chance that a target will be present in appreciable amounts is good. The raw data were normalized using 18S rRNA (universally expressed in all tissues and cells).

For each expression analysis, tumor tissue samples were compared with matched
20 normal tissues from the same patient. A gene was considered overexpressed in a tumor when the level of expression of the gene was 2 fold or higher in the tumor compared with its matched normal sample. In cases where normal tissue was not available, a universal pool of cDNA samples was used instead. In these cases, a gene was considered overexpressed in a tumor sample when the difference of expression levels between a
25 tumor sample and the average of all normal samples from the same tissue type was greater than 2 times the standard deviation of all normal samples (i.e., $\text{Tumor} - \text{average}(\text{all normal samples}) > 2 \times \text{STDEV}(\text{all normal samples})$).

Results are shown in Table 1. Data presented in bold indicate that greater than 50% of tested tumor samples of the tissue type indicated in row 1 exhibited over
30 expression of the gene listed in column 1, relative to normal samples. Underlined data indicates that between 25% to 49% of tested tumor samples exhibited over expression. A modulator identified by an assay described herein can be further validated for therapeutic effect by administration to a tumor in which the gene is overexpressed. A decrease in tumor growth confirms therapeutic utility of the modulator. Prior to treating a patient with

the modulator, the likelihood that the patient will respond to treatment can be diagnosed by obtaining a tumor sample from the patient, and assaying for expression of the gene targeted by the modulator. The expression data for the gene(s) can also be used as a diagnostic marker for disease progression. The assay can be performed by expression analysis as described above, by antibody directed to the gene target, or by any other available detection method.

TABLE I

	Breast	Breast	Colon	Lung	Ovary	Ovary	Kidney
Gene Name	Matched	Total	Matched	Matched	Matched	Total	Matched
RRP1	0 / 3	0 / 5	<u>11 / 26</u>	8 / 14	<u>1 / 4</u>	<u>3 / 14</u>	2 / 19
RRP2	<u>1 / 3</u>	<u>1 / 5</u>	3 / 26	7 / 14	2 / 4	9 / 14	12 / 19
RRP3	0 / 3	0 / 5	4 / 26	<u>4 / 14</u>	<u>1 / 4</u>	<u>3 / 14</u>	<u>4 / 19</u>
RRP4	0 / 3	0 / 5	3 / 26	<u>3 / 14</u>	<u>1 / 4</u>	<u>5 / 14</u>	<u>4 / 19</u>
RRP5	0 / 3	0 / 5	<u>6 / 26</u>	1 / 14	<u>1 / 4</u>	1 / 14	<u>4 / 19</u>
RRP6	2 / 3	3 / 5	<u>10 / 26</u>	<u>5 / 14</u>	0 / 4	0 / 14	13 / 19
RRP7	0 / 3	0 / 5	<u>6 / 26</u>	1 / 14	<u>1 / 4</u>	1 / 14	1 / 19
RRP8	<u>1 / 3</u>	<u>1 / 5</u>	<u>6 / 26</u>	<u>5 / 14</u>	0 / 4	0 / 14	3 / 19

VI. Cell Biology and Functional Characterization

RRP Localization. In *Drosophila*, rhomboid is a cell surface protein. However, localization of human RRP's has not happened to date. Furthermore, in humans, it is not known whether each of the termini of the rhomboids is intracellular or extracellular. To answer these questions, RRP1 was subcloned into pcDNA expression vector (Invitrogen) in frame with myc-his tag at the c-terminus, according to manufacturer's protocols. The resulting expression vector was transiently transfected using Lipofectamine Plus reagent (Life technologies) into human embryonic kidney HEK-293 cells. Immunofluorescence staining using anti-myc antibody was then carried out on the cells to localize RRP1. Results of these experiments indicated that RRP1 is expressed at the cell surface. Furthermore, permeabilized (0.1% triton in PBS) and non-permeabilized cells show the same staining pattern, indicating that the c-terminus of the protein is outside, and the N-

terminus, inside. RRP2 – RRP8 are also subcloned and tagged at the c-terminus for localization of each protein, and also to assess the direction of the protein ends in each case.

5 **Involvement of RRP in EGFR signaling pathway.** While *Drosophila* rhomboid is essential in the EGFR pathway, the function of human RRP1 have not been assessed to date. To assess the role of human RRP1 in the EGFR signaling pathway, RRP1 was stably transfected using CaPO4 transfection kit (Clontech) into Hela cells, which have endogenous EGFR activity. Pooled stable cells were then examined in the following
10 assays.

EGFR activation as measured by tyrosine phosphorylation

Hela cells overexpressing RRP1 and parental Hela cells were tested for EGFR tyrosine phosphorylation by immunoblot against anti-phosphotyrosine using anti-phosphotyrosine antibody (Upstate Biotechnology). Cells overexpressing RRP1
15 demonstrated up to two-fold increase in EGFR tyrosine phosphorylation as compared to parental Hela controls.

EGFR expression in cells overexpressing RRP1.

Expression of EGFR was increased when tested by immunoblot using anti-EGFR antibody against direct cell lysates in cells overexpressing RRP1. Furthermore, expression
20 of EGFR was also increased in tumor samples overexpressing RRP1, by TaqMan® analysis. The correlation of both in-vitro and in-vivo results for concordant overexpression of EGFR and RRP1 is quite significant, and provides a tool for assessing the functional relationship of the two proteins in any tumor sample or cell line.

Cell proliferation and Migration.

25 Hela cells overexpressing RRP1 demonstrate a two-fold increase in cell proliferation compared to Hela parental cells as measured by Procheck (Intergen) and AlamarBlue (Biosource International) chemiluminescent assays. In addition, Fluroblok (BD Biosource) analysis showed a slight increase in motility in cells overexpressing RRP1 compared to Hela parental cells.

30 Taken together, these results demonstrate a strong functional conservation of rhomboids across evolution. Furthermore, increased RRP1 expression in tumor samples and cell lines, concurrent increased expression of EGFR, and increased proliferation and motility of cells expressing RRP1 suggest involvement of RRP1 in various cancers. RRP1 expression levels can thus be used to screen for tumors with defective EGFR pathways, to

identify tumors amenable to treatment with the compounds and antibodies identified by the methods of the present invention. Same aforementioned experiments are performed with RRP2 – RRP8 to assess their involvement in the EGFR signaling pathway.

5 **RRP1 binding target.** In *Drosophila*, rhomboid transforms Spitz (homolog of human TGF- α) from a membrane-bound to a secreted form, and thus triggers and upregulates the DER (*Drosophila* EGFR) pathway. No such function has been shown for human rhomboids. HEK293 cells were transiently co-transfected with RRP1 and TM-TGF- α transmembrane-bound TGF), then immunoprecipitated with anti-myc antibody
10 (for RRP1) and immunoblotted with anti-TGF- α . Cells transfected with vector alone, RRP1 alone, or TM-TGF- α alone were used as controls. A band corresponding to the size of TGF- α was recognized by anti-TGF- α antibody, while no bands were detected in the control lanes. This data indicates that RRP1 physically interacts with TGF- α . The same experiments are performed with RRP2 – RRP8 to determine their binding partners.

15

Effect of RRP1 on HB_EGF and ERK phosphorylation. RRP1 was transfected into HEK293 embryonic kidney cells to investigate effects of RRP1 on HB-EGF (an EGFR ligand) and ERK phosphorylation. The RRP1 full-length clone was inserted into the mammalian expression vector pCDNA3.1 (Invitrogen) with N-terminal FLAG,
20 according to manufacturer's protocol. HEK293T cells were seeded into 6-well plates at 600,000 cells per well, grown in DMEM with 10% Fetal Bovine Serum (FBS) and a Penicillin/Streptomycin cocktail. Seeded cells were allowed to grow for 24 hours at 37 °C and 5% CO₂. Cells were transfected with 2 μ g of RRP1 plasmid per well using LipofectAmine PLUS reagent (Life Technologies) according to manufacturer's protocol.
25 Three hours after transfection, FBS concentration was brought up to 10%, total volume was increased to 4 ml with DMEM, and cells were placed back into incubator for 48 hours. Transfected cells were then processed for Western analysis. Cells were lysed in NP40 lysis buffer containing protease and phosphatase inhibitors. Protein concentrations were determined with Pierce BCA Protein Assay Reagent, and 20 μ g of protein was added
30 to each lane on a 10% NUPAGE Bis-Tris Gel (Invitrogen), run and transferred to nitrocellulose paper, following instructions in the NUPAGE Bis-Tris instruction booklet. To block, blots were soaked in 10ml of 0.5% non-fat milk for 2 hours, and washed. Primary antibodies were then added at appropriate dilution (diluted in binding buffer): Mouse α -FLAG: 1:1000; Mouse α -HBEGF: 1:1000 (R&D Systems); Rabbit α -Total

ERK: 1:1000 (Cell Signaling); Rabbit α -Phosphorylated ERK: 1:1000 (Cell Signaling). Blots were rinsed and secondary antibodies were added: Donkey α -rabbit: 1:10000 (Jackson ImmunoResearch, HRP conjugated); Donkey α -mouse: 1:10000 (Jackson ImmunoResearch, HRP conjugated). Blots were then rinsed, incubated 1:1 Supersignal West Pico Stable Peroxide Solution and Luminol /Enhancer solution (Pierce) for 5 minutes according to instructions, and exposed to film. In these experiments, RRP-1 transfected HEK-293T cells showed elevated levels of HB-EGF, and elevated ERK phosphorylation.

In *Drosophila*, RRP1 has been shown to increase EGFR signaling through cleaving and releasing activated EGFR ligand (spitz; described *supra*). Our findings are consistent with those reports. Transient transfection of human RRP1 in HEK293T cells clearly induced ERK phosphorylation, a signaling event down-stream of EGFR activation. In addition, co-transfection of RRP1 and HB-EGF (one of EGFR ligands in human) in HEK293T cells yielded higher HB-EGF protein level than HB-EGF transfection along, suggesting RRP1 may increase the overall level of EGFR ligands in cells. Overall, these experiments suggest that human RRP1 increases EGFR signaling.

Treatment of RRP1transfected cells with EGFR and HB-EGF inhibitors. The aim of this experiment was to test the effect of EGFR and HB-EGF inhibitors on phosphorylated ERK levels in RRP1 transfected cells. HEK293T cells were grown and transfected as described above. At 24 and 44 hours after transfection, the following EGFR or HB-EGF inhibitors were added to wells: for HB_EGF: HB-EGF mouse monoclonal antibody (R&D Sysytems; 10ug/ml), and CRM197 (10ug/ml). For EGFR: Ab225 mouse monoclonal antibody (ATCC; 5ug/ml), and 168393 small molecule inhibitor (Calbiochem; 2.7uM). At 48 hours post transfection, cell lysates were collected and analyzed by western analysis, as described above. Both EGFR inhibitors (Ab225 and 168393) decreased the levels of pERK in RRP1 transfected cells. Anti HBEGF antibody also was able to reduce pERK levels in RRP1 transfected cells. The HBEGF inhibitor CRM197 did not have an affect. This result may be potentially useful in designing an assay to test antibodies/compounds that affect the MAPK signaling cascade. Based on the observed results, we propose the following hypothesis: human RRP1 increases EGFR/Erk signaling through cleavage/shedding of HB-EGF. EGFR signaling is a pivotal factor in tumorigenesis, and HB-EGF is one of the most important growth factors that drive tumor

development through autocrine and other mechanisms. Therefore, RRP1 may play an important role in cancer and other diseases that link to irregularity of EGFR signaling.

Antibodies against human rhomboids. A peptide antibody, SEQ ID NO: 17, and referred to as peptide 1, was made against the N-terminus of RRP1. This antibody
5 displayed strong affinity for denatured RRP1, such as in cells fixed with formalin or proteins on western blots. In addition, staining pattern remained the same when permeabilized cells were co-stained with anti-myc antibodies (for the c-terminus of RRP1) and peptide 1, suggesting that Peptide 1 does recognize RRP1. Furthermore, the cells were significantly immunostained only when they were permeabilized, suggesting that the
10 N-terminus of RRP1 is inside the cells. This result correlates with the previous staining pattern using anti-myc antibody for the C-terminus.

RRP1 protein expression in tumors was examined by Immunohistochemistry (IHC) using peptide 1. Positive staining on epithelial cells of colon adenocarcinoma tissue sections were observed as compared to background staining in preimmune serum,
15 suggesting the antibody is specific for its target, RRP1, in colon cancer cells. Moreover, increased staining of colorectal adenocarcinomas was observed compared to matched normal tissues using peptide 1 antibody. Taken together, these results suggest that peptide 1 antibody is a powerful tool to assess expression levels of RRP1 in tumor samples. Antibodies are also produced against the rest of RRP1s to assess their expression levels in
20 tumors.

VII. Mouse RRP1 (mRRP1) gene structure

A BAC shotgun sequencing approach (for a review, see Green ED. (2001) Nature Review Genetics 2:573-583) was used to determine the intron-exon structure of the
25 murine RRP1 gene. Based upon the genebank entries of a partial rat cDNA and the human gene structure, a PCR primer pair was designed that allowed the amplification of a part of the murine RRP1 gene from mouse genomic DNA. (RRP1s (SEQ ID NO:48): 5'-CACCCACCAGCCGCACTGGTC; RRP8as (SEQ ID NO:49): 5'-GAGCAGCTAGGGTTCAATG, 95°C 120sec, 95°C 45sec, 60°C 45sec, 72°C 45sec,
30 40cycles, 1.5mM MgCl₂). This PCR primer set was then used to screen a commercially available ES-129SvJ BAC library from Incyte Genomics (St. Louis, USA). 2 PCR positive clones were identified (clone addresses 178/O15 and 195/D15). Clone 195/D15 was chosen to construct a shotgun sequencing library according to standard procedures (TOPO Shotgun Subcloning Kit, #K-7000-01, Invitrogen). Upon sequence assembly, a

contiguous 30 Kilobases (kb) genomic sequence was identified that carried the entire mRRP1 gene. Subsequently, the DNA from clone 195/D15 BAC served as a PCR template to amplify mRRP1 genomic regions used for the gene targeting approach. The mRRP1 gene contains 8 exons, which span a genomic region of at least 2.6 kb. The translation start on the sequence stretch defined as exon 1 has not been identified so that additional 5' exons may exist.

VIII. Generation of the mice

Targeting Strategy

We utilized a targeting strategy that allows sequential deletion of vector sequences in the same cells. Chimeras generated by injection of mRRP1 targeted ES-cells can be used to create i) KO mice harboring a completely inactivated mRRP gene by breeding to Cre-deleter mice (described below) as well as ii) mice with functional mRRP1 that can be deleted in a tissue specific or time specific manner by breeding to FLP-deleter mice (described below).

This strategy has the advantage that the NEO selection cassette is removed in both chimeras, thus avoiding potential hypomorphic effect of the selection marker.

Targeting vector

The targeting vector, comprising the nucleotide sequence to be incorporated into the wild-type (WT) genomic sequence, and one or more selectable markers (FIG. 1) was made based on a BAC genomic clone from the mouse strain 129SvJ background obtained from Incyte Genomics. The mRRP1 targeting vector was designed such that exons 2-8 of the mRRP1 gene were flanked with Cre recombinase recognition sequences (loxP sites) after homologous recombination. Exons 2-8 of the murine RRP encode the rhomboid domain which provide protein activity, therefore a deletion of exons 2-8 results in a null allele. Three genomic fragments were amplified by PCR and cloned individually in reversed orientation into the base vector pEasyfloxFRTGK12 (FIG. 1). This base vector has: an ampicillin resistance gene; an origin of replication; a pgtN29 plasmid (New England Biolabs), used as the source of pgk neo; pGT60mcs plasmid (InvivoGen), used as the source for TK reading frame; and LoxP and FRT sites, introduced by synthetic oligonucleotides. The 5' region of homology (fragment A) amplified with oligo 164 (SEQ ID NO:62) and oligo 144 (SEQ ID NO:50) covered a 3.5 kb fragment that was inserted into PacI / PmeI sites upstream the 2nd loxP site of pEasyfloxFRTGK12. A 2.5 kb 3'

fragment encompassing exons 2-8 (fragment B) and amplified with oligos 145 (SEQ ID NO:51) and 146 (SEQ ID NO:52) was cloned into a *AscI* / *FseI* sites downstream of 2nd loxP site and upstream of 2nd FRT site and *PGK-neo* cassette of pEasyfloxFRTGK12. An additional 1.0 kb fragment further 3' (fragment C) and amplified with oligos 147 (SEQ ID NO:53) and 148 (SEQ ID NO:54) was inserted into *NotI*/*SgfI* sites located downstream the 1st FRT and 1st loxP site of pEasyfloxFRTGK12. The loxP- and FRT-flanked neomycin resistance cassette downstream of exons 2-8 of the RRP gene allowed genomic selection for homologous recombination and a *PGK-thymidine kinase* gene cassette (PGK-TK) inserted at the 5' end of the construct was used to select against random integration events.

In order to establish a positive control for the PCR screen on ES cell clones showing homologous recombination events, an additional 1.25 kb fragment was amplified using oligo 147 (SEQ ID NO:53) and oligo 149 (SEQ ID NO:55) and cloned into *NotI* / *SgfI* sites in pEasyfloxFRTGK12. This fragment extended the 3' homology region of the final targeting vector by 250 base pairs, allowing establishment of PCR screening with a vector specific oligo 174 (SEQ ID NO:64) and RRP gene specific oligo 169 (SEQ ID NO:63) located 3' external in relation to the targeting vector.

Generation of targeted ES-cells

Three parental embryonic stem (ES) cell lines **B6-2** (EUROGENETEC, Ref: TG-ES01-02, derived from mouse strain C57BL/6 N TacfBr), **SvEv** (EUROGENETEC, Ref: TG-ES01-01, derived from mouse strain 129S6/SvEvTacfBr), and **E-14** ((Hooper et al., (1987) *Nature (London)* 326, 292-295); derived from mouse strain 129P2/OlaHsd) were employed for targeting of the mRRP1 locus.

All ES cell lines were cultured and electroporated under the conditions described in Torres, R. M. & Kuehn, R. Laboratory Protocols for Conditional Gene Targeting, Oxford, New York, Tokyo: 1997. In brief, for electroporation 1×10^7 cells were mixed with 30 μ g of *SfiI* linearized targeting DNA in a total volume of 800 μ l transfection buffer and electroporated using a Bio-Rad Gene Pulser (0,24 kV, 500 μ F, 0,4 cm electrode distance). Cells were reseeded on 10-cm tissue culture dishes, containing selection resistant mouse embryonic fibroblast feeder cells (prepared from targeted mouse strain IL4 pepneo; Mueller et al., (1992) *Eur. J. Immunol.* 21: 921 -25) at a density of 2.5×10^6 cells per plate. Drug selection (G418 (Gibco/BRL # 10131-019)), 200 mg/ml) was started after 1 day. ES cell clones were negatively selected the 3rd day after electroporation with the

antibiotic Gancyclovir (Cymeven, Roche), 2 μ M). After 8 days, resistant colonies were randomly picked and individually expanded. Correctly targeted ES clones were obtained for all 3 ES cell lines employed in the experiment.

- Successful targeting of the mRRP gene was detected in resistant ES clones by PCR-screening with primer oligo 174 (SEQ ID NO:64) and oligo 169 (SEQ ID NO:63). Corecombination of the distal loxP site was detected by using oligo 200 (SEQ ID NO:65) and oligo 201 (SEQ ID NO:66). PCR-positive clones were expanded and confirmed by EcoRI / SfiI restriction enzyme digestion and Southern hybridization employing the 5' external probe amplified with oligos 152 (SEQ ID NO:56) and 153 (SEQ ID NO:57). Upon homologous recombination, the wild type 10.0 kb band was reduced to 7.5 kb (Figure 2). Corecombination of the 2nd loxP site in homologous recombinants was confirmed by HindIII and 3' external probe amplified with oligos 160 (SEQ ID NO:60) and 162 (SEQ ID NO:61)(Figure 2). Upon corecombination of the 2nd loxP site, the wildtype 4.3 kb band was increased to 5.8 kb (Figure 2). Single integration of the targeting vector was confirmed by HindIII digestion and Southern hybridization using the internal probe amplified with oligos 154 (SEQ ID NO:58) and 155 (SEQ ID NO:59)(Figure 2).

Table 2 provides a summary of these experiments.

- Table 2: Homologous recombination of the RRP-1 targeting construct in various ES cells

ES Cell	strain	# cells x 10 ⁷	G418 res. anal. Clones	HR	2. loxP #	Clone
B6-2	C57BL/6	4	2000 876	3	2	G-D5 E-D9
SvEv	129S6	2	Not counted 95	1	0	
		3	Not counted 295	1	1	D-E7
E-14	129P2	6	Not counted 641	11	3	B-E2 E-F2 C-C8

Generation of mRRP1 chimeras

Chimeras were generated by injection of targeted ES cells into Balb/C host embryos (Hogan, B., Beddington, R., Costantini, F. & Lacy, E. eds. Manipulating the Mouse Embryo, a Laboratory Manual. 2nd ed. Cold Spring Harbor, New York: Cold Spring Harbor, Laboratory Press, 1994). Germline transmission was obtained by breeding chimeric mice to C57BL/6 females, resulting in mRRP1 heterozygous animals on a pure C57BL/6 strain background and hybrid (C57BL/6 x 129SvEv)_{F1} and (C57BL/6 x 129P2/Ola)_{F1} background.

Resulting animals were typed by Southern blot hybridization on tail DNA after digestion with HindIII with the 3' external probe. In heterozygote animals for the target allele, the 4.3kb wild type band was increased to 5.8 kb.

Table 3 provides a summary of these experiments.

Table 3: Summary of chimera production and germline transmission of the target allele.

Clone	blastocyst s transf.	pups born	# of chimeras	GL transmis.	heterozyg . mice
B6-2					
G-D5	246	37	32	10/11	yes
E-D9	372	103	70	6/10	yes
Total	618	140	102	17/21	
SvEv					
D-E7	302	71	40	8/10	yes
E-14					
B-E2	64	17	14	8/10	yes
Grand Total	984	228	156	33 / 41	

Heterozygous animals by in vivo deletion.

Target allele with Cre directed deletion

An in vivo deletion approach was pursued to generate C57BL/6 mice carrying the target alleles with Cre-directed deletion from mRRP1. Using this approach, the Cre transgene directs the removal of the sequences flanked by the LoxP sites, and thus, creates an RRP1 knockout in every cell of the animal (FIG.3).

Heterozygote mice carrying the mRRP1 target allele (mRRP1^{targ/+}) from the B6-2 clones G-D5 and E-D9 (Table 2) were bred with transgenic C57BL/6-Cre-Deleter mice (Schwenk et al, NAR (95) Vol. 23, No 24, 5080-5081). This deleter mice have been crossed back to B6 for 10 generations, to establish the transgene on an congenic C57BL/6

background. This was done to maintain the C57BL/6 background after the in vivo deletion of the mRRP1 targeted allele. Resulting pups were genotyped on tail DNA for the presence of the target allele with Cre-directed deletion of mRRP1 by Southern Blot hybridization and by PCR for the presence of the Cre transgene.

- 5 Tail DNA was digested with HindIII and hybridized with the 3' external probe. In heterozygote animals the 5.8 kb band was reduced to a 1.3 kb band for the target allele. Alternatively animals were typed by PCR.

Target allele with Flp-directed deletion

- 10 An in vivo deletion approach was pursued, to generate C57BL/6 mice carrying the target allele with Flp directed deletion of the Neo marker from RRP1. In this case, the mRRP1 gene is still functional in every cell, but the animals may later be crossed with conditional Cre deleter mice for tissue specific or time-specific deletion of the sequences flanked by LoxP to create conditional mRRP1 knockout mice.

- 15 To avoid any potential hypomorphic effects, the Frt flanked Neo selection marker was removed in vivo by Flp recombination. Heterozygote mice carrying the mRRP1 target allele from the B6-2 clones **G-D5** and **E-D9** (Table 2) were bred with transgenic C57BL/6 mice heterozygous for the CAAGS-FLPe transgene on a pure B6 background (for FLPe description, see Rodriguez et al., (2000) Nat Genet 25:139-140). Resulting
20 pups were genotyped on tail DNA for the presence of the target allele by Southern Blot hybridization and by PCR for the presence of the CAAGS-Flp transgene.

- Tail DNA was digested with HindIII and hybridized the 3' external probe. In heterozygote animals the 4.3 kb target allele band was reduced to a 3.8 kb band after the removal of the Frt-flanked Neo marker (FIG.4). Alternatively animals were typed by
25 PCR.

Homozygous animals

Target allele with Cre directed deletion

- To generate mice homozygous for the mRRP1 knockout allele on the C57BL/6
30 background, heterozygous animals derived from the in vivo deleted clones **G-D5** and **E-D9** were intercrossed. From this cross 17 litters were born, giving rise to 106 pups. 28 of these pups died pre-weaning. Two of these pups were recovered and genotyped by PCR as homozygous complete KO. 49 of the living animals were weaned and 14 animals were genotyped on tail DNA with the 3' external probe after digestion with HindIII. In

homozygous animals the original 4.3 kb wild type band was reduced to a 1.3 kb band. Alternatively animals were typed by PCR.

From the 14 animals 3 typed as homozygous mutant, 7 typed as heterozygous mutant and 4 typed as wildtype. These numbers matched the expected Mendelian distribution. The homozygous animals appear to be normal. No gross difference was detected in comparison to their wildtype littermates.

To test for the fertility of the homozygous mutant animals and to increase further the colony, homozygous mutant animals for the mRRP1 derived from the in vivo deleted clones G-D5 were intercrossed. To date, 14 pups have been born from these intercrosses, and 4 have been weaned, demonstrating the fertility of the homozygous mRRP1 KO mice.

The homozygous mRRP1 KO mice and cells are used to analyze mRRP1 function in mammals. Data from Example VI show that overexpression of human RRP1 in HeLa cells results in enhanced phosphorylation of EGFR. Inactivation of mRRP1 gene function should therefore result in attenuated phosphorylation of EGFR and or MAPK in tissues obtained from mRRP1 KO mice.

Growth factors like EGF, PDGF cause proliferation in fibroblast cell culture. As Drosophila Rhomboid and mammalian RRP are thought to release membrane-bound growth factors by cleavage, RRP function with regard to cell cycle progression and proliferation is analyzed with embryonic and adult fibroblasts isolated from mRRP1 KO mice.

Direct effects of RRP function on tumor development is analyzed by crossing mRRP1 KO mice with APCMin mice, a model for human colon cancer. As the mRRP1 KO strain and the APCmin (Moser AR et al., (1993) Proc Natl Acad Sci U S A 90:8977-81) strain are both on C57BL/6 background, tumor formation can be analyzed on pure genetic backgrounds, and thus side effects resulting from mixed genetic backgrounds can be excluded. A strong reduction of small intestinal tumor formation in mRRP1 ^{-/-}, APCMin ^{+/-} mice versus APCMin ^{+/-} mice is expected, demonstrating that mRRP1 is a powerful target for anti-cancer drug development.

RRP is located at the cell surface and therefore a target for antibody therapy. As the extracellular loops of mouse and human RRP protein are strongly conserved, it is difficult to generate antibodies against human RRP in mice, as mice will recognize the human RRP as self. Therefore, RRP KO mice are used for the production of monoclonal antibodies directed against human RRP protein.

WHAT IS CLAIMED IS:

1. A method of specifically inhibiting growth of tumor cells that overexpress an RRP protein comprising contacting said tumor cells with an amount of an exogenous RRP binding agent that results in inhibition in growth of tumor cells.
5
2. The method of claim 1 wherein said tumor cells are from a tissue selected from the group consisting of breast, ovary, lung, kidney, and colon.
- 10 3. The method of claim 1 wherein said RRP binding agent is an antibody or a fragment of said antibody.
4. The method according to claim 3 wherein said antibody is humanized.
- 15 5. The method according to claim 3 wherein said antibody comprises human constant region residues and rhomboid-specific mouse variable region residues.
6. The method according to claim 3 wherein said antibody is bound to a detectable label.
- 20 7. The method according to claim 3 wherein said antibody is bound to a cytotoxic label.
8. A method of screening for agents that modulate the interaction of an RRP polypeptide with an RRP binding target, comprising:
 - a) expressing a recombinant RRP polypeptide,
 - 25 b) incubating the recombinant RRP polypeptide with an RRP binding target and a candidate RRP modulating agent and
 - c) determining whether said candidate RRP modulating agent modulates the binding of the RRP polypeptide with the RRP binding target.
- 30 9. The method according to claim 8 wherein said binding target is selected from the group consisting of TGF α , EGF, and amphiregulin.
10. The method according to claim 8 wherein said binding target is TGF α .

11. The method according to claim 8 wherein said candidate RRP modulating agent is an antibody.
12. The method according to claim 8 wherein said candidate RRP modulating agent is a
5 small organic molecule.
13. A method of diagnosing a tumor cell as having abnormal p53 or p21 pathway signaling, comprising measuring RRP expression levels in said tumor cell, wherein increased RRP expression levels in the tumor cell compared to a non-tumor cell of the
10 same tissue or organ as the tumor cell, indicates defective p53 or p21 pathway signaling in the tumor cell.
14. A method of identifying a candidate p53 or p21 pathway modulating agent, said method comprising the steps of:
- 15 a) providing an assay system comprising a purified RRP polypeptide or nucleic acid or a functionally active fragment or derivative thereof;
- b) contacting the assay system with a test agent under conditions whereby, but for the presence of the test agent, the system provides a reference activity; and
- 20 c) detecting a test agent-biased activity of the assay system, wherein a difference between the test agent-biased activity and the reference activity identifies the test agent as a candidate p53 or p21 pathway modulating agent.
15. The method of Claim 14 wherein the assay system comprises cultured cells that express the RRP polypeptide.
- 25 16. The method of Claim 15 wherein the cultured cells additionally have defective p53 or p21 function.
17. The method of Claim 14 wherein the assay system includes a screening assay
30 comprising a RRP polypeptide, and the candidate test agent is a small molecule modulator.
18. The method of Claim 17 wherein the assay is a protease assay.

19. The method of Claim 14 wherein the assay system is selected from the group consisting of an apoptosis assay system, a cell proliferation assay system, an angiogenesis assay system, and a hypoxic induction assay system.
- 5 20. The method of Claim 14 wherein the assay system includes a binding assay comprising a RRP polypeptide and the candidate test agent is an antibody.
21. The method of Claim 14 wherein the assay system includes an expression assay comprising a RRP nucleic acid and the candidate test agent is a nucleic acid modulator.
- 10 22. The method of claim 21 wherein the nucleic acid modulator is an antisense oligomer.
23. The method of Claim 21 wherein the nucleic acid modulator is a PMO.
- 15 24. The method of Claim 14 additionally comprising:
d) administering the candidate p53 or p21 pathway modulating agent identified in (c) to a model system comprising cells defective in p53 or p21 function and, detecting a phenotypic change in the model system that indicates that the p53 or p21 function is restored.
- 20 25. The method of Claim 24 wherein the model system is a mouse model with defective p53 or p21 function.
26. A method for modulating a p53 or p21 pathway of a cell comprising contacting a cell
25 defective in p53 or p21 function with a candidate modulator that specifically binds to an RRP polypeptide comprising an amino acid sequence selected from group consisting of SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, 16, and 46 whereby p53 or p21 function is restored.
27. The method of claim 26 wherein the candidate modulator is administered to a
30 vertebrate animal predetermined to have a disease or disorder resulting from a defect in p53 or p21 function.
28. The method of Claim 26 wherein the candidate modulator is selected from the group consisting of an antibody and a small molecule.

29. The method of Claim 14, comprising the additional steps of:
- e) providing a secondary assay system comprising cultured cells or a non-human animal expressing RRP ,
 - 5 f) contacting the secondary assay system with the test agent of (b) or an agent derived therefrom under conditions whereby, but for the presence of the test agent or agent derived therefrom, the system provides a reference activity; and
 - g) detecting an agent-biased activity of the second assay system,
 - h) wherein a difference between the agent-biased activity and the reference activity of
 - 10 the second assay system confirms the test agent or agent derived therefrom as a candidate p53 or p21 pathway modulating agent,
 - i) and wherein the second assay detects an agent-biased change in the p53 or p21 pathway.
- 15 30. The method of Claim 29 wherein the secondary assay system comprises cultured cells.
31. The method of Claim 29 wherein the secondary assay system comprises a non-human animal.
- 20 32. The method of Claim 31 wherein the non-human animal mis-expresses a p53 or p21 pathway gene.
33. A method of modulating p53 or p21 pathway in a mammalian cell comprising contacting the cell with an agent that specifically binds a RRP polypeptide or nucleic acid.
- 25 34. The method of Claim 33 wherein the agent is administered to a mammalian animal predetermined to have a pathology associated with the p53 or p21 pathway.
35. The method of Claim 33 wherein the agent is a small molecule modulator, a nucleic acid modulator, or an antibody.
- 30 36. A method for diagnosing a disease in a patient comprising:
- a) obtaining a biological sample from the patient;
 - b) contacting the sample with a probe for RRP expression;

- c) comparing results from step (b) with a control;
- d) determining whether step (c) indicates a likelihood of disease.

37. The method of claim 36 wherein said disease is cancer.

5

38. The method of claim 37, wherein said cancer is a cancer as shown in Table 1 as having >25% expression level.

39. A purified nucleic acid molecule that encodes a polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:46, or reverse complement thereof.

10

40. The nucleic acid molecule of Claim 39 which is capable of hybridizing to a nucleic acid sequence of SEQ ID NO:45 using high stringency hybridization conditions.

41. A recombinant expression system comprising a DNA or RNA molecule, wherein said expression system is capable of producing an mRRP1 polypeptide comprising the amino acid sequence of SEQ ID NO:46 when said expression system is present in a compatible host cell.

15

42. A host cell comprising the expression system of claim 41.

20

43. A process for producing an mRRP1 protein comprising culturing the host cell of Claim 42 under conditions suitable for expression of said mRRP1 protein and recovering said protein.

25

44. A process for producing a cell which produces an mRRP1 protein comprising the transformation or transfection of a host cell with the expression system of claim 41 such that the host cell, under appropriate culture conditions, produces an mRRP1 protein.

45. A recombinant host cell expressing the protein produced by the method of claim 44.

30

46. A transgenic mouse whose genome comprises a disruption in an endogenous RRP gene wherein said disruption results in decreased expression or a lack of expression of said endogenous RRP gene.

47. The mouse of claim 46, wherein said mouse is homozygous for the disrupted RRP gene.

5 48. A cell isolated from the transgenic mouse of claim 46.

49. A mouse cell whose genome comprises a disruption in an endogenous RRP gene wherein said disruption results in decreased expression or a lack of expression of said endogenous RRP gene.

10

50. A method of selecting an agent that modulates cell proliferation comprising:

- a) providing a first and a second isolated mouse cell wherein the genome of both the first and second isolated mouse cell has been manipulated to comprise a disruption in an endogenous RRP gene wherein said disruption results in decreased expression or a
15 lack of expression of said endogenous RRP gene;
- b) administering an agent to said first isolated mouse cell ; and
- c) determining the amount of proliferation of the first and second cell, wherein a difference in the amount of proliferation of the first cell as compared to the second cell indicates that the agent modulates cell proliferation.

20

51. A method of making an antibody against a human RRP comprising:

- a) injecting RRP or an antigenic fragment thereof to mouse of claim 46; and
- b) recovering said antibody.

25


FIG.1 pEasyfloxFRTGK12 mRRP1

FIG.2 mRRP1 gene targeting






FIG.3

FIG.4

SEQUENCE LISTING

<110> EXELIXIS, INC.

<120> RRP SEQUENCES AND KNOCKOUT MICE AND USES THEREOF

<130> EX02-004C-PC

<150> US 10/056,790

<151> 2002-01-23

<150> US 09/908,419

<151> 2001-07-18

<150> US 60/219,289

<151> 2000-07-19

<150> US 60/277,487

<151> 2001-03-21

<150> US 60/304,863

<151> 2001-07-12

<150> US 60/277,471

<151> 2001-03-21

<150> US60/305,017

<151> 2001-07-12

<150> US 60/296,076

<151> 2001-06-05

<150> US 60/328,605

<151> 2001-10-10

<150> US 60/328,491

<151> 2001-10-10

<160> 70

<170> PatentIn version 3.2

<210> 1

<211> 1559

<212> DNA

<213> Homo sapiens

<400> 1

cttggacctt ggccctcgct ttccaggatg ggtagggtgg aagacggggg aacaactgag	60
gagctggagg actgggaccc aggcaccagt gccctgccag ctcttgggat caagcagggt	120
cccagggaac agacaggcac gggggccctg tcccaaaagt gctgggagcc tgagcctgat	180
gctcccagcc agcctggccc agccctttgg tccagggggtc gggcccgcac tcaggccttg	240
gctggcgggt cctcactgca gcagctggac cccgagaaca caggcttcat cgggtcggac	300
accttactg gcctggtgca cagccatgag ctgcccttg acccggccaa gctggacatg	360
ctggtggccc tggctcagag caacgagcag ggccaggctt gctaccagga gctggtggac	420

```

ctgatcagca gcaagcgctc cagcagtttc aagcgggcca ttgctaacgg acagcgggca 480
ctgccccggg acgggcccgt ggatgagcca ggcctaggtg tctacaagcg gtttgtgctg 540
tacgtggcct acgagatcct gccttgtgag gtggaccgcc gctgggtactt ctaccgtcac 600
cgcagctgcc cccccccgt gttcatggcc tcggtcactc ttgccagat catcgtgttc 660
ctgtgttacg gggcccgcct caacaagtgg gtgctgcaga cctaccacc cgagtacatg 720
aagagcccc ttgtgtacca ccccgggcac cgtgcccgcg cctggcgctt cctcacctac 780
atgttcatgc acgttgggct ggagcagctg gggttcaacg ccctcctgca gctgatgatc 840
ggggtgcccc tggagatggg gcacggcctg ctccgcatca gcctgctcta cctggcaggg 900
gtgctggcag gctccctaac cgtctccatc accgacatgc gggccccggg ggtgggaggg 960
tccggcgggg tctacgccct gtgctcggca cacctggcca acgttgtcat gaactgggct 1020
gggatgagat gtccctacaa gttgctgagg atgggtgctg ccttgggtgtg catgagctcc 1080
gaggtggggc gggccgtgtg gctgcgcttc tcccgcgcg tgccgcctc gggcccacag 1140
cccagcttca tggcgcacct ggcaggcgcg gtgggtgggg tgagcatggg cctgaccatc 1200
ctgcggagct acgaggagcg cctgcgggac cagtgcggct ggtgggtggg gctgctggcc 1260
tacggcacct tcctgctctt cgcgctcttc tggaacgtct tcgcctacga cctgctgggc 1320
gccacatcc cccaccgcc ctgaccggct acctgaggct gcacaggcca gggctcgggc 1380
atgtggtggc cggccaccag gggccttcac gtctgccctt tgtgaacgga cgtctcaggg 1440
ctgctgtgcc ccttgggtgt ggggtggcctc aaaggaggcc ctgtcccagc caccaccccc 1500
ccactcccag gacttgcggg ctgagccttt ttggataatt aataaatatt ttacacagc 1559

```

```

<210> 2
<211> 438
<212> PRT
<213> Homo sapiens

```

```

<400> 2

```

```

Met Gly Arg Val Glu Asp Gly Gly Thr Thr Glu Glu Leu Glu Asp Trp
1           5           10          15

```

```

Asp Pro Gly Thr Ser Ala Leu Pro Ala Pro Gly Ile Lys Gln Gly Pro
20           25          30

```

```

Arg Glu Gln Thr Gly Thr Gly Pro Leu Ser Gln Lys Cys Trp Glu Pro
35           40          45

```

```

Glu Pro Asp Ala Pro Ser Gln Pro Gly Pro Ala Leu Trp Ser Arg Gly
50           55          60

```

Arg Ala Arg Thr Gln Ala Leu Ala Gly Gly Ser Ser Leu Gln Gln Leu
 65 70 75 80
 Asp Pro Glu Asn Thr Gly Phe Ile Gly Ala Asp Thr Phe Thr Gly Leu
 85 90 95
 Val His Ser His Glu Leu Pro Leu Asp Pro Ala Lys Leu Asp Met Leu
 100 105 110
 Val Ala Leu Ala Gln Ser Asn Glu Gln Gly Gln Val Cys Tyr Gln Glu
 115 120 125
 Leu Val Asp Leu Ile Ser Ser Lys Arg Ser Ser Ser Phe Lys Arg Ala
 130 135 140
 Ile Ala Asn Gly Gln Arg Ala Leu Pro Arg Asp Gly Pro Leu Asp Glu
 145 150 155 160
 Pro Gly Leu Gly Val Tyr Lys Arg Phe Val Arg Tyr Val Ala Tyr Glu
 165 170 175
 Ile Leu Pro Cys Glu Val Asp Arg Arg Trp Tyr Phe Tyr Arg His Arg
 180 185 190
 Ser Cys Pro Pro Pro Val Phe Met Ala Ser Val Thr Leu Ala Gln Ile
 195 200 205
 Ile Val Phe Leu Cys Tyr Gly Ala Arg Leu Asn Lys Trp Val Leu Gln
 210 215 220
 Thr Tyr His Pro Glu Tyr Met Lys Ser Pro Leu Val Tyr His Pro Gly
 225 230 235 240
 His Arg Ala Arg Ala Trp Arg Phe Leu Thr Tyr Met Phe Met His Val
 245 250 255
 Gly Leu Glu Gln Leu Gly Phe Asn Ala Leu Leu Gln Leu Met Ile Gly
 260 265 270
 Val Pro Leu Glu Met Val His Gly Leu Leu Arg Ile Ser Leu Leu Tyr
 275 280 285
 Leu Ala Gly Val Leu Ala Gly Ser Leu Thr Val Ser Ile Thr Asp Met
 290 295 300
 Arg Ala Pro Val Val Gly Gly Ser Gly Gly Val Tyr Ala Leu Cys Ser
 305 310 315 320

Ala His Leu Ala Asn Val Val Met Asn Trp Ala Gly Met Arg Cys Pro
 325 330 335

Tyr Lys Leu Leu Arg Met Val Leu Ala Leu Val Cys Met Ser Ser Glu
 340 345 350

Val Gly Arg Ala Val Trp Leu Arg Phe Ser Pro Pro Leu Pro Ala Ser
 355 360 365

Gly Pro Gln Pro Ser Phe Met Ala His Leu Ala Gly Ala Val Val Gly
 370 375 380

Val Ser Met Gly Leu Thr Ile Leu Arg Ser Tyr Glu Glu Arg Leu Arg
 385 390 395 400

Asp Gln Cys Gly Trp Trp Val Val Leu Leu Ala Tyr Gly Thr Phe Leu
 405 410 415

Leu Phe Ala Val Phe Trp Asn Val Phe Ala Tyr Asp Leu Leu Gly Ala
 420 425 430

His Ile Pro Pro Pro Pro
 435

<210> 3
 <211> 1224
 <212> DNA
 <213> Homo sapiens

<400> 3
 gggccattta ggaggttag atcattttga tcatcttcag ctgtcttctc ttcacatata 60
 ggaaaggcct tggaagcag tcgttgcgcc agacagccca gggaagagcg gcagcctgag 120
 gacctagggc cacctgctgt tccctgggat tcatgtcctt ctggggagga gggaggaccc 180
 aggacaatgg ctgctgttca tgatctggag atggagagca tgaatctgaa tatggggaga 240
 gagatgaaag aagagctgga ggaagaggag aaaatgagag aggatggggg aggtaaagat 300
 cgggccaaga gtaaaaaggt ccacaggatt gtctcaaaat ggatgctgcc cgaaaagtcc 360
 cgaggaacat acttgagag agctaactgc ttccgcctc ccgtgttcat catctccatc 420
 agcctggccg agctggcagt gtttatttac tatgctgtgt ggaagcctca gaaacagtgg 480
 atcacgttgg acacaggcat cttggagagt ccctttatct acagtcctga gaagagggag 540
 gaagcctgga ggtttatctc atacatgctg gtacatgctg gagttcagca catcttgggg 600
 aatctttgta tgcagcttgt tttgggtatt cccttggaat tgggtccaca aggcctccgt 660

```

gtggggctgg tgtacctggc aggagtgatt gcaggggtccc ttgccagctc catctttgac    720
ccactcagat atcttggtggg agcttcagga ggagtctatg ctctgatggg aggctatttt    780
atgaatgttc tgggtgaattt tcaagaaatg attcctgcct ttggaatttt cagactgctg    840
atcatcatcc tgataattgt gttggacatg ggatttgctc tctatagaag gttcttttgtt    900
cctgaagatg ggtctccggg gtcttttgca gctcacattg caggtggatt tgctggaatg    960
tccattgggt acacgggtgtt tagctgcttt gataaagcac tgctgaaaga tccaagggtt   1020
tggatagcaa ttgctgcata tttagcttgt gtcttatttg ctgtgttttt caacattttc   1080
ctatctccag caaactgacc tgccoctatt gtaagtcaat taataaaaag agccatctgg   1140
aggaaataaa aaaaaaagga agactctatg aagaaacaga gaagtctcag aaaaggctaa   1200
caattttaga tagagaacaa aggg                                     1224

```

```

<210> 4
<211> 303
<212> PRT
<213> Homo sapiens

```

```

<400> 4

```

```

Met Ala Ala Val His Asp Leu Glu Met Glu Ser Met Asn Leu Asn Met
1              5              10              15

```

```

Gly Arg Glu Met Lys Glu Glu Leu Glu Glu Glu Lys Met Arg Glu
20              25              30

```

```

Asp Gly Gly Gly Lys Asp Arg Ala Lys Ser Lys Lys Val His Arg Ile
35              40              45

```

```

Val Ser Lys Trp Met Leu Pro Glu Lys Ser Arg Gly Thr Tyr Leu Glu
50              55              60

```

```

Arg Ala Asn Cys Phe Pro Pro Pro Val Phe Ile Ile Ser Ile Ser Leu
65              70              75              80

```

```

Ala Glu Leu Ala Val Phe Ile Tyr Tyr Ala Val Trp Lys Pro Gln Lys
85              90              95

```

```

Gln Trp Ile Thr Leu Asp Thr Gly Ile Leu Glu Ser Pro Phe Ile Tyr
100             105             110

```

```

Ser Pro Glu Lys Arg Glu Glu Ala Trp Arg Phe Ile Ser Tyr Met Leu
115             120             125

```

```

Val His Ala Gly Val Gln His Ile Leu Gly Asn Leu Cys Met Gln Leu
130             135             140

```

Val Leu Gly Ile Pro Leu Glu Met Val His Lys Gly Leu Arg Val Gly
145 150 155 160

Leu Val Tyr Leu Ala Gly Val Ile Ala Gly Ser Leu Ala Ser Ser Ile
165 170 175

Phe Asp Pro Leu Arg Tyr Leu Val Gly Ala Ser Gly Gly Val Tyr Ala
180 185 190

Leu Met Gly Gly Tyr Phe Met Asn Val Leu Val Asn Phe Gln Glu Met
195 200 205

Ile Pro Ala Phe Gly Ile Phe Arg Leu Leu Ile Ile Ile Leu Ile Ile
210 215 220

Val Leu Asp Met Gly Phe Ala Leu Tyr Arg Arg Phe Phe Val Pro Glu
225 230 235 240

Asp Gly Ser Pro Val Ser Phe Ala Ala His Ile Ala Gly Gly Phe Ala
245 250 255

Gly Met Ser Ile Gly Tyr Thr Val Phe Ser Cys Phe Asp Lys Ala Leu
260 265 270

Leu Lys Asp Pro Arg Phe Trp Ile Ala Ile Ala Ala Tyr Leu Ala Cys
275 280 285

Val Leu Phe Ala Val Phe Phe Asn Ile Phe Leu Ser Pro Ala Asn
290 295 300

<210> 5
<211> 1376
<212> DNA
<213> Homo sapiens

<400> 5
ctgaactgat gaggttttca tgggaaatca gtgcctgtct gttcatctga acaccacta 60
gttattcatc caacaaatct tgattgagtg ctgataatgc caggctctgt gctaagtacc 120
ggggataagt tgtgataaccg acccgcgagg cgccgcggtc caaggaggag gcaaaagcag 180
acagacatca gtgtgggctg gagggcccaga ggtctggaca gaacagaggg ttccgtgaga 240
acaggccatg gctgagtttg accctgggaa cacaggctac attagcacag gcaagttccg 300
gagtcttctg gagagccaca gctccaagct ggaccgcac aaaagggagg tcctcctggc 360
tcttgccgac agccacgcgg atgggcagat cggtaccag gattttgtca gcctagtgag 420

```

caacaagcgt tccaacagct tccgccaagc catcctgcag ggcaaccgca ggctaagcag 480
caaggccctg ctggaggaga aggggctgag cctctcgcag cgacttatcc gccatgtggc 540
ctatgagacc ctgccccggg aaattgaccg caagtggtag tatgacagct acacctgtg 600
ccccccacc tggttcatga tcacagtcac gctgctggag gttgcctttt tcctctacaa 660
tggggtgtca ctaggtcaat ttgtactgca ggtaactcat ccacgttact tgaagaactc 720
cctggtttac caccacagc tgcgagcaca ggtttggcgc tacctgacat acatcttcat 780
gcatgcaggg atagaacacc tgggactcaa tgtgggtgctg cagctgctgg tgggggtgcc 840
cctggagatg gtgcatggag ccacccgaat tgggcttgct tacgtggccg gtgttgtggc 900
agggtccttg gcagtgtctg tggctgacat gaccgctcca gtcgtgggct cttctggagg 960
ggtgtatgct ctctctctg cccatctggc caacattgtc atgaactggc caggcatgaa 1020
gtgccagttc aagctgctgc ggatggctgt ggcccttacc tgtatgagca tggagtttgg 1080
gcgggccgtg tggtccgct tccaccgctc ggcctatccc ccgtgccctc acccaagctt 1140
tgtggcgcac ttgggtggcg tggccgtggg catcacctg ggcgtgggtg tcctgaggaa 1200
ctacgagcag aggtccagg accagtcact gtggtggatt tttgtggcca tgtacaccgt 1260
cttcgtgctg ttcgtgtct tctggaacat ctttgcttac accctgctgg acttaaagct 1320
gccgcctccc ccctgagggc tggaggccca aggtcgggga ggggagggaa aagcag 1376

```

```

<210> 6
<211> 362
<212> PRT
<213> Homo sapiens

```

```

<400> 6

```

```

Met Ala Glu Phe Asp Pro Gly Asn Thr Gly Tyr Ile Ser Thr Gly Lys
1          5          10          15

```

```

Phe Arg Ser Leu Leu Glu Ser His Ser Ser Lys Leu Asp Pro His Lys
20          25          30

```

```

Arg Glu Val Leu Leu Ala Leu Ala Asp Ser His Ala Asp Gly Gln Ile
35          40          45

```

```

Gly Tyr Gln Asp Phe Val Ser Leu Val Ser Asn Lys Arg Ser Asn Ser
50          55          60

```

```

Phe Arg Gln Ala Ile Leu Gln Gly Asn Arg Arg Leu Ser Ser Lys Ala
65          70          75          80

```

```

Leu Leu Glu Glu Lys Gly Leu Ser Leu Ser Gln Arg Leu Ile Arg His
85          90          95

```

Val Ala Tyr Glu Thr Leu Pro Arg Glu Ile Asp Arg Lys Trp Tyr Tyr
 100 105 110

Asp Ser Tyr Thr Cys Cys Pro Pro Pro Trp Phe Met Ile Thr Val Thr
 115 120 125

Leu Leu Glu Val Ala Phe Phe Leu Tyr Asn Gly Val Ser Leu Gly Gln
 130 135 140

Phe Val Leu Gln Val Thr His Pro Arg Tyr Leu Lys Asn Ser Leu Val
 145 150 155 160

Tyr His Pro Gln Leu Arg Ala Gln Val Trp Arg Tyr Leu Thr Tyr Ile
 165 170 175

Phe Met His Ala Gly Ile Glu His Leu Gly Leu Asn Val Val Leu Gln
 180 185 190

Leu Leu Val Gly Val Pro Leu Glu Met Val His Gly Ala Thr Arg Ile
 195 200 205

Gly Leu Val Tyr Val Ala Gly Val Val Ala Gly Ser Leu Ala Val Ser
 210 215 220

Val Ala Asp Met Thr Ala Pro Val Val Gly Ser Ser Gly Gly Val Tyr
 225 230 235 240

Ala Leu Val Ser Ala His Leu Ala Asn Ile Val Met Asn Trp Ser Gly
 245 250 255

Met Lys Cys Gln Phe Lys Leu Leu Arg Met Ala Val Ala Leu Ile Cys
 260 265 270

Met Ser Met Glu Phe Gly Arg Ala Val Trp Leu Arg Phe His Pro Ser
 275 280 285

Ala Tyr Pro Pro Cys Pro His Pro Ser Phe Val Ala His Leu Gly Gly
 290 295 300

Val Ala Val Gly Ile Thr Leu Gly Val Val Val Leu Arg Asn Tyr Glu
 305 310 315 320

Gln Arg Leu Gln Asp Gln Ser Leu Trp Trp Ile Phe Val Ala Met Tyr
 325 330 335

Thr Val Phe Val Leu Phe Ala Val Phe Trp Asn Ile Phe Ala Tyr Thr
 340 345 350

Leu Leu Asp Leu Lys Leu Pro Pro Pro Pro
 355 360

<210> 7
 <211> 1140
 <212> DNA
 <213> Homo sapiens

<400> 7
 atggcgtggc gaggtctgggc gcagagaggc tggggctgcg gccaggcgtg ggggtgcgtcg 60
 gtgggcggcc gcagctgcga ggagctcact gcggtcctaa ccccgccgca gtcctctgga 120
 cgcaggttta acttctttat tcaacaaaaa tgcggattca gaaaagcacc caggaagggtt 180
 gaacctcgaa gatcagaccc agggacaagt ggtgaagcat acaagagaag tgctttgatt 240
 cctcctgtgg aagaaacagt cttttatcct tctccctatc ctataaggag tctcataaaa 300
 cctttatttt ttactgttgg gtttacaggc tgtgcatttg gatcagctgc tatttggcaa 360
 tatgaatcac tgaatccag ggtccagagt tattttgatg gtataaaagc tgattgggtg 420
 gatagcataa gaccacaaaa agaaggagac ttcagaaagg agattaacaa gtggtggaat 480
 aacctaagtg atggccagcg gactgtgaca ggtattatag ctgcaaatgt ccttgtattc 540
 tgtttatgga gactaccttc tctgcagcgg acaatgatca gatatttcac atcgaatcca 600
 gcctcaaagg tcctttgttc tccaatgttg ctgtcaacat tcagtcactt ctccttattt 660
 cacatggcag caaatatgta tgttttgtgg agcttctctt ccagcatagt gaacattctg 720
 ggtcaagagc agttcatggc agtgtaccta tctgcagggtg ttatttccaa ttttgtcagt 780
 tacctgggta aagttgccac aggaagatat ggaccatcac ttggtgcatc tgggtgccatc 840
 atgacagtcc tcgcagctgt ctgcactaag atcccagaag ggaggcttgc cattattttc 900
 cttccgatgt tcacgttcac agcagggaat gccctgaaag ccattatcgc catggataca 960
 gcaggaatga tcctgggatg gaaatTTTTT gatcatgcgg cacatcttgg gggagctctt 1020
 tttggaatat ggtatgttac ttacgggtcat gaactgattt ggaagaacag ggagccgcta 1080
 gtgaaaatct ggcataaat aaggactaat ggccccaaaa aaggaggtgg ctctaagtaa 1140

<210> 8
 <211> 379
 <212> PRT
 <213> Homo sapiens

<400> 8

Met Ala Trp Arg Gly Trp Ala Gln Arg Gly Trp Gly Cys Gly Gln Ala
 1 5 10 15

Trp Gly Ala Ser Val Gly Gly Arg Ser Cys Glu Glu Leu Thr Ala Val
 20 25 30

Leu Thr Pro Pro Gln Leu Leu Gly Arg Arg Phe Asn Phe Phe Ile Gln
 35 40 45

Gln Lys Cys Gly Phe Arg Lys Ala Pro Arg Lys Val Glu Pro Arg Arg
 50 55 60

Ser Asp Pro Gly Thr Ser Gly Glu Ala Tyr Lys Arg Ser Ala Leu Ile
 65 70 75 80

Pro Pro Val Glu Glu Thr Val Phe Tyr Pro Ser Pro Tyr Pro Ile Arg
 85 90 95

Ser Leu Ile Lys Pro Leu Phe Phe Thr Val Gly Phe Thr Gly Cys Ala
 100 105 110

Phe Gly Ser Ala Ala Ile Trp Gln Tyr Glu Ser Leu Lys Ser Arg Val
 115 120 125

Gln Ser Tyr Phe Asp Gly Ile Lys Ala Asp Trp Leu Asp Ser Ile Arg
 130 135 140

Pro Gln Lys Glu Gly Asp Phe Arg Lys Glu Ile Asn Lys Trp Trp Asn
 145 150 155 160

Asn Leu Ser Asp Gly Gln Arg Thr Val Thr Gly Ile Ile Ala Ala Asn
 165 170 175

Val Leu Val Phe Cys Leu Trp Arg Val Pro Ser Leu Gln Arg Thr Met
 180 185 190

Ile Arg Tyr Phe Thr Ser Asn Pro Ala Ser Lys Val Leu Cys Ser Pro
 195 200 205

Met Leu Leu Ser Thr Phe Ser His Phe Ser Leu Phe His Met Ala Ala
 210 215 220

Asn Met Tyr Val Leu Trp Ser Phe Ser Ser Ser Ile Val Asn Ile Leu
 225 230 235 240

Gly Gln Glu Gln Phe Met Ala Val Tyr Leu Ser Ala Gly Val Ile Ser
 245 250 255

Asn Phe Val Ser Tyr Leu Gly Lys Val Ala Thr Gly Arg Tyr Gly Pro
 260 265 270

Ser Leu Gly Ala Ser Gly Ala Ile Met Thr Val Leu Ala Ala Val Cys
 275 280 285

Thr Lys Ile Pro Glu Gly Arg Leu Ala Ile Ile Phe Leu Pro Met Phe
 290 295 300

Thr Phe Thr Ala Gly Asn Ala Leu Lys Ala Ile Ile Ala Met Asp Thr
 305 310 315 320

Ala Gly Met Ile Leu Gly Trp Lys Phe Phe Asp His Ala Ala His Leu
 325 330 335

Gly Gly Ala Leu Phe Gly Ile Trp Tyr Val Thr Tyr Gly His Glu Leu
 340 345 350

Ile Trp Lys Asn Arg Glu Pro Leu Val Lys Ile Trp His Glu Ile Arg
 355 360 365

Thr Asn Gly Pro Lys Lys Gly Gly Gly Ser Lys
 370 375

<210> 9
 <211> 2964
 <212> DNA
 <213> Homo sapiens

<400> 9
 gactcgggcgc gggcgccctc ccggccagcg gcggcagccc ctctcccccg gcgcccctcag 60
 gacccccccag agaccccccg cggcggcagc ctgccttgct ctgccaggaa ccatgagtga 120
 ggccccgcagg gacagcacga gcagcctgca gcgcaagaag ccaccctggc taaagctgga 180
 cattccctct gcggtgcccc tgacggcaga agagcccagc ttcctgcagc ccctgaggcg 240
 acaggctttc ctgaggagtg tgagtatgcc agccgagaca gccacatct cttcacccca 300
 ccatgagctc cggcggccgg tgctgcaacg ccagacgtcc atcacacaga ccatccgcag 360
 ggggaccgcc gactggtttg gactgagcaa ggacagtgc agcaccaga aatggcagcg 420
 caagagcatc cgctactgca gccagcgcta cggaagctg aagccccagg tcctccggga 480
 gctggacctg cccagccagg acaacgtgtc gctgaccagc accgagacgc cccccact 540
 ctacgtgggg ccatgccagc tgggcatgca gaagatcata gacccctgg cccgtggccg 600
 tgccttcctg gtggcagatg aactgcgga aggcctgagt gcccacaca ctcccgtcac 660
 gccgggtgct gcctccctct gctccttctc cagctcccg ctaggtttcc accggctccc 720

gcggcgggcg	aagcgagagt	cggtggccaa	gatgagcttc	cgggcgggcg	cagcgctgat	780
gaaaggccgc	tccgttaggg	atggcacctt	tcgccgggca	cagcgtcgaa	gcttcactcc	840
agctagcttt	ctggaggagg	acacaactga	tttccccgat	gagctggaca	catccttctt	900
tgcccgggaa	ggtatcctcc	atgaagagct	gtccacatac	ccggatgaag	ttttcgagtc	960
cccatcggag	gcagcgctaa	aggactggga	gaaggcaccg	gagcaggcgg	acctcaccgg	1020
cggggccctg	gaccgcagcg	agcttgagcg	cagccacctg	atgctgccct	tggagcgagg	1080
ctggcggaag	cagaaggagg	gcgccgcagc	cccgcagccc	aagggtcggc	tccgacagga	1140
ggtggtgagc	accgcggggc	cgcgacgggg	ccagcgatc	gcggtgccgg	tgcgcaagct	1200
cttcgcccgg	gagaagcggc	cgtatgggct	gggcatgggtg	ggacggctca	ccaaccgcac	1260
ctaccgcaag	cgcacgaca	gcttcgtcaa	gcgccagatc	gaggacatgg	acgaccacag	1320
gcccttcttc	acctactggc	ttaccttctg	gcactcgctc	gtcgccatcc	tagccgtgtg	1380
catctatggc	atcgcgcccg	tgggcttctc	gcagcatgag	acggtggact	cggtgctgcg	1440
gaaccgcggg	gtctacgaga	acgtcaagta	cgtgcagcag	gagaacttct	ggatcggggc	1500
cagctcggag	gccctcatcc	acctggggcg	caagtttctg	ccctgcatgc	gccaggaccc	1560
gcagggtgcac	agcttcattc	gctcggcgcg	cgagcgcgag	aagcactccg	cctgctgcgt	1620
gcgcaacgac	aggtcgggct	gcgtgcagac	ctcgaggagg	gagtgtctgt	ccacgctggc	1680
agtgtgggtg	aagtggccca	tccatcccag	cgcccagag	cttgcggggc	acaagagaca	1740
gtttggctct	gtctgccacc	aggatcccag	ggtgtgtgat	gagccctcct	ccgaagaccc	1800
tcatgagtgg	ccagaagaca	tcaccaagtg	gccgatctgc	accaaaaaca	gcgctgggaa	1860
ccacaccaac	catccccaca	tggactgtgt	catcacagga	cggccctgct	gcattggcac	1920
caagggcagg	tgtgagatca	cctcccggga	gtactgtgac	ttcatgaggg	gctacttcca	1980
tgaggaggcc	acgctctgct	ctcagggtgca	ctgcatggat	gatgtgtgtg	ggctcctgcc	2040
ttttctcaac	cccgagggtgc	ctgaccagtt	ctaccgcctg	tggctatccc	tcttcctgca	2100
cgcggggatc	ttgcaactgc	tgggtgtccat	ctgcttcocag	atgactgtcc	tgcgggacct	2160
ggagaagctg	gcaggctggc	accgcatagc	catcatctac	ctgctgagtg	gtgtcaccgg	2220
caacctggcc	agtgccatct	tcctgccata	ccgagcagag	gtgggtcctg	ctggctccca	2280
gttcggcatc	ctggcctgcc	tcttcgtgga	gctcttcocag	agctggcaga	tcctggcgcg	2340
gccctggcgt	gccttcttca	agctgctggc	tgtggtgctc	ttcctcttca	cctttgggct	2400
gctgccgtgg	attgacaact	ttgccacat	ctcgggggttc	atcagtggcc	tcttcctctc	2460
cttcgccttc	ttgccctaca	tcagctttgg	caagttcgac	ctgtaccgga	aacgctgcca	2520
gatcatcatc	tttcagggtg	tcttcctggg	cctcctgggt	ggcctgggtg	tcctcttcta	2580

```

cgtctatcct gtccgctgtg agtgggtgtga gttcctcacc tgcattcccct tcaactgacaa 2640
gttctgtgag aagtacgaac tggacgctca gctccactga gctggctgcg ggctccagcg 2700
gccgtgtgct ccagcaggcc agagccagac acgacctccc tgagcctcac aggcttacag 2760
gagtcacctg ctccatgtgg ggactggcct gtttcttgaa cacagacctc tttcttgtgc 2820
cttgttcact tctgttgaac cctcgtact gccgggcatt tattatacta cttcctgtca 2880
taaccttcta acttgtttct tgacgaccac ctcatgtggc caataaatga actgggagcg 2940
ttttaaaaaa aaaaaaaaaa aaaa 2964

```

<210> 10
 <211> 855
 <212> PRT
 <213> Homo sapiens

<400> 10

Met Ser Glu Ala Arg Arg Asp Ser Thr Ser Ser Leu Gln Arg Lys Lys
1 5 10 15

Pro Pro Trp Leu Lys Leu Asp Ile Pro Ser Ala Val Pro Leu Thr Ala
20 25 30

Glu Glu Pro Ser Phe Leu Gln Pro Leu Arg Arg Gln Ala Phe Leu Arg
35 40 45

Ser Val Ser Met Pro Ala Glu Thr Ala His Ile Ser Ser Pro His His
50 55 60

Glu Leu Arg Arg Pro Val Leu Gln Arg Gln Thr Ser Ile Thr Gln Thr
65 70 75 80

Ile Arg Arg Gly Thr Ala Asp Trp Phe Gly Val Ser Lys Asp Ser Asp
85 90 95

Ser Thr Gln Lys Trp Gln Arg Lys Ser Ile Arg His Cys Ser Gln Arg
100 105 110

Tyr Gly Lys Leu Lys Pro Gln Val Leu Arg Glu Leu Asp Leu Pro Ser
115 120 125

Gln Asp Asn Val Ser Leu Thr Ser Thr Glu Thr Pro Pro Pro Leu Tyr
130 135 140

Val Gly Pro Cys Gln Leu Gly Met Gln Lys Ile Ile Asp Pro Leu Ala
145 150 155 160

Arg Gly Arg Ala Phe Arg Val Ala Asp Asp Thr Ala Glu Gly Leu Ser
 165 170 175

Ala Pro His Thr Pro Val Thr Pro Gly Ala Ala Ser Leu Cys Ser Phe
 180 185 190

Ser Ser Ser Arg Ser Gly Phe His Arg Leu Pro Arg Arg Arg Lys Arg
 195 200 205

Glu Ser Val Ala Lys Met Ser Phe Arg Ala Ala Ala Ala Leu Met Lys
 210 215 220

Gly Arg Ser Val Arg Asp Gly Thr Phe Arg Arg Ala Gln Arg Arg Ser
 225 230 235 240

Phe Thr Pro Ala Ser Phe Leu Glu Glu Asp Thr Thr Asp Phe Pro Asp
 245 250 255

Glu Leu Asp Thr Ser Phe Phe Ala Arg Glu Gly Ile Leu His Glu Glu
 260 265 270

Leu Ser Thr Tyr Pro Asp Glu Val Phe Glu Ser Pro Ser Glu Ala Ala
 275 280 285

Leu Lys Asp Trp Glu Lys Ala Pro Glu Gln Ala Asp Leu Thr Gly Gly
 290 295 300

Ala Leu Asp Arg Ser Glu Leu Glu Arg Ser His Leu Met Leu Pro Leu
 305 310 315 320

Glu Arg Gly Trp Arg Lys Gln Lys Glu Gly Ala Ala Ala Pro Gln Pro
 325 330 335

Lys Val Arg Leu Arg Gln Glu Val Val Ser Thr Ala Gly Pro Arg Arg
 340 345 350

Gly Gln Arg Ile Ala Val Pro Val Arg Lys Leu Phe Ala Arg Glu Lys
 355 360 365

Arg Pro Tyr Gly Leu Gly Met Val Gly Arg Leu Thr Asn Arg Thr Tyr
 370 375 380

Arg Lys Arg Ile Asp Ser Phe Val Lys Arg Gln Ile Glu Asp Met Asp
 385 390 395 400

Asp His Arg Pro Phe Phe Thr Tyr Trp Leu Thr Phe Val His Ser Leu
 405 410 415

Val Ala Ile Leu Ala Val Cys Ile Tyr Gly Ile Ala Pro Val Gly Phe
 420 425 430

Ser Gln His Glu Thr Val Asp Ser Val Leu Arg Asn Arg Gly Val Tyr
 435 440 445

Glu Asn Val Lys Tyr Val Gln Gln Glu Asn Phe Trp Ile Gly Pro Ser
 450 455 460

Ser Glu Ala Leu Ile His Leu Gly Ala Lys Phe Ser Pro Cys Met Arg
 465 470 475 480

Gln Asp Pro Gln Val His Ser Phe Ile Arg Ser Ala Arg Glu Arg Glu
 485 490 495

Lys His Ser Ala Cys Cys Val Arg Asn Asp Arg Ser Gly Cys Val Gln
 500 505 510

Thr Ser Glu Glu Glu Cys Ser Ser Thr Leu Ala Val Trp Val Lys Trp
 515 520 525

Pro Ile His Pro Ser Ala Pro Glu Leu Ala Gly His Lys Arg Gln Phe
 530 535 540

Gly Ser Val Cys His Gln Asp Pro Arg Val Cys Asp Glu Pro Ser Ser
 545 550 555 560

Glu Asp Pro His Glu Trp Pro Glu Asp Ile Thr Lys Trp Pro Ile Cys
 565 570 575

Thr Lys Asn Ser Ala Gly Asn His Thr Asn His Pro His Met Asp Cys
 580 585 590

Val Ile Thr Gly Arg Pro Cys Cys Ile Gly Thr Lys Gly Arg Cys Glu
 595 600 605

Ile Thr Ser Arg Glu Tyr Cys Asp Phe Met Arg Gly Tyr Phe His Glu
 610 615 620

Glu Ala Thr Leu Cys Ser Gln Val His Cys Met Asp Asp Val Cys Gly
 625 630 635 640

Leu Leu Pro Phe Leu Asn Pro Glu Val Pro Asp Gln Phe Tyr Arg Leu
 645 650 655

Trp Leu Ser Leu Phe Leu His Ala Gly Ile Leu His Cys Leu Val Ser
 660 665 670

Ile Cys Phe Gln Met Thr Val Leu Arg Asp Leu Glu Lys Leu Ala Gly
 675 680 685

Trp His Arg Ile Ala Ile Ile Tyr Leu Leu Ser Gly Val Thr Gly Asn
 690 695 700

Leu Ala Ser Ala Ile Phe Leu Pro Tyr Arg Ala Glu Val Gly Pro Ala
 705 710 715 720

Gly Ser Gln Phe Gly Ile Leu Ala Cys Leu Phe Val Glu Leu Phe Gln
 725 730 735

Ser Trp Gln Ile Leu Ala Arg Pro Trp Arg Ala Phe Phe Lys Leu Leu
 740 745 750

Ala Val Val Leu Phe Leu Phe Thr Phe Gly Leu Leu Pro Trp Ile Asp
 755 760 765

Asn Phe Ala His Ile Ser Gly Phe Ile Ser Gly Leu Phe Leu Ser Phe
 770 775 780

Ala Phe Leu Pro Tyr Ile Ser Phe Gly Lys Phe Asp Leu Tyr Arg Lys
 785 790 795 800

Arg Cys Gln Ile Ile Ile Phe Gln Val Val Phe Leu Gly Leu Leu Ala
 805 810 815

Gly Leu Val Val Leu Phe Tyr Val Tyr Pro Val Arg Cys Glu Trp Cys
 820 825 830

Glu Phe Leu Thr Cys Ile Pro Phe Thr Asp Lys Phe Cys Glu Lys Tyr
 835 840 845

Glu Leu Asp Ala Gln Leu His
 850 855

<210> 11
 <211> 2643
 <212> DNA
 <213> Homo sapiens

<400> 11
 gctactccca cctgccacgc cgcaagagaa tgtctgtggc ccacatgagc ttgcaagctg 60
 ccgctgccct cctcaagggg cgctcggtgc tggatgccac cggacagcgg tgccgggtgg 120

tcaagcgcag ctttgccttc ccgagcttcc tggaggagga tgtggtcgat ggggcagaca	180
cgtttgactc ctcccttttt agtaaggaag aaatgagctc catgcctgat gatgtctttg	240
agtccccccc actctctgcc agctacttcc gagggatccc aactcagcc tcccctgtct	300
ccccgatgg ggtgcaaadc cctctgaagg agtatggccg agccccagtc cccgggcccc	360
ggcgcggaac gcgcacgcc tocaaggtga agcactttgc ctttgatcgg aagaagcggc	420
actacggcct cggcgtgggt ggcaactggc tgaaccgcag ctaccgccgc agcatcagca	480
gcactgtgca gggcgagctg gagagcttcg acagccaccg gccctacttc acctactggc	540
tgaccttcgt ccatgtcatc atcacgctgc tgggtgatttg cacgtatggc atcgaccccg	600
tgggctttgc ccagcacgtc accaccacgc tgggtgctgc gaacaaaggt gtgtacgaga	660
gcgtgaagta catccagcag gagaacttct gggttggccc cagctcgatt gacctgatcc	720
acctgggggc caagttctca cctgcatcc ggaaggacgg gcagatcgag cagctgggtgc	780
tgcgcgagcg agacctggag cgggactcag gctgctgtgt ccagaatgac cactccggat	840
gcatccagac ccagcggaag gactgctcgg agactttggc cacttttgtc aagtggcagg	900
atgacactgg gcccccatg gacaagtctg atctgggcca gaagcggact tcgggggctg	960
tctgccacca ggacccagg acctgcgagg agccagcctc cagcggtgcc cacatctggc	1020
ccgatgacat cactaagtgg ccgatctgca cagagcaggc caggagcaac cacacaggct	1080
tcctgcacat ggactgcgag atcaagggcc gccctgctg catcggcacc aagggcagct	1140
gtgagatcac caccgggaa tactgtgagt tcatgcacgg ctatttccat gaggaagcaa	1200
cactctgctc ccagggtcac tgcttggaca aggtgtgtgg gctgctgcc ttctcaacc	1260
ctgaggtccc agatcagttc tacaggctct ggctgtctct ctctctacat gctggcgtgg	1320
tgcactgcct cgtgtctgtg gtctttcaaa tgaccatcct gagggacctg gagaagctgg	1380
ccggctggca ccgatcgc atcatcttca tcctcagtgg catcacaggc aacctcgcca	1440
gtgccatctt tctccatac cgggcagagg tgggcccggc cggtcacag ttccggcctc	1500
tcgcctgcct ctctgtggag ctcttcaga gctggccgct gctggagagg ccctggaagg	1560
ccttcctcaa cctctcgcc atcgtgctct tcctgttcat ctgtggcctc ttgccctgga	1620
tcgacaacat cggccacatc ttccgcttcc tcagtggcct gctgctggc ttgccttcc	1680
tgccctacat caccttcggc accagcgaca agtaccgcaa gggggcactc atcctggtgt	1740
cactgctggc ctttgccggc ctcttcgccc cctcgtgct gtggctgtac atctaccca	1800
ttactggcc ctggatcgag cacctcacct gcttccccct caccagccgc ttctgcgaga	1860
agtatgagct ggaccagggt ctgcactgac cgctgggcca cacggctgcc cctcagccct	1920
gctggaacag ggtctgcctg cgagggtgc cctctgcaga gcgtctctg tgtgccagag	1980

```

agccagagac ccaagacagg gcccgggctc tggacctggg tgccccctg ccaggcgagg 2040
ctgactccgc gtgagatggt tggttaaggc ggggtttttc tggggcgtga ggcctgtgag 2100
atcctgaccc aagctcaggc acaccaagg cacctgcctc tctgagtctt gggctctcagt 2160
tcctaataatc cgcctccttg ctgagaccat ctctggggc agggctcttt tcttcccagg 2220
tcctcagcgc tgcctctgct ggtgccttct cccccactac tactggagcg tgccttgct 2280
ggggacgtgg ctgtgccctc agttgcccc agggctgggt gccaccatg ccccttctc 2340
tttctcctcc tacctctgcc ctgtgagccc atccataagg ctctcagatg ggacattgta 2400
ggaaaggctt tggccatggt ctgggggcag agaacaagg gggagacaca agtagacctc 2460
aggtagaacg acaccgggcg gagccacccc agggcctgct cccaggagg gctcgaggcg 2520
catcaggccc gttttttacc agtttatatc acggtcttca tttttaaaag taacgctaac 2580
tttgtacgga cgatgtctca tggattaaat aatattcttt atggcaaaaa aaaaaaaaaa 2640
aaa 2643

```

<210> 12
 <211> 619
 <212> PRT
 <213> Homo sapiens

<400> 12

Met Ser Val Ala His Met Ser Leu Gln Ala Ala Ala Leu Leu Lys
 1 5 10 15

Gly Arg Ser Val Leu Asp Ala Thr Gly Gln Arg Cys Arg Val Val Lys
 20 25 30

Arg Ser Phe Ala Phe Pro Ser Phe Leu Glu Glu Asp Val Val Asp Gly
 35 40 45

Ala Asp Thr Phe Asp Ser Ser Phe Phe Ser Lys Glu Glu Met Ser Ser
 50 55 60

Met Pro Asp Asp Val Phe Glu Ser Pro Pro Leu Ser Ala Ser Tyr Phe
 65 70 75 80

Arg Gly Ile Pro His Ser Ala Ser Pro Val Ser Pro Asp Gly Val Gln
 85 90 95

Ile Pro Leu Lys Glu Tyr Gly Arg Ala Pro Val Pro Gly Pro Arg Arg
 100 105 110

Gly Lys Arg Ile Ala Ser Lys Val Lys His Phe Ala Phe Asp Arg Lys
 115 120 125

Lys Arg His Tyr Gly Leu Gly Val Val Gly Asn Trp Leu Asn Arg Ser
 130 135 140

Tyr Arg Arg Ser Ile Ser Ser Thr Val Gln Arg Gln Leu Glu Ser Phe
 145 150 155 160

Asp Ser His Arg Pro Tyr Phe Thr Tyr Trp Leu Thr Phe Val His Val
 165 170 175

Ile Ile Thr Leu Leu Val Ile Cys Thr Tyr Gly Ile Ala Pro Val Gly
 180 185 190

Phe Ala Gln His Val Thr Thr Gln Leu Val Leu Arg Asn Lys Gly Val
 195 200 205

Tyr Glu Ser Val Lys Tyr Ile Gln Gln Glu Asn Phe Trp Val Gly Pro
 210 215 220

Ser Ser Ile Asp Leu Ile His Leu Gly Ala Lys Phe Ser Pro Cys Ile
 225 230 235 240

Arg Lys Asp Gly Gln Ile Glu Gln Leu Val Leu Arg Glu Arg Asp Leu
 245 250 255

Glu Arg Asp Ser Gly Cys Cys Val Gln Asn Asp His Ser Gly Cys Ile
 260 265 270

Gln Thr Gln Arg Lys Asp Cys Ser Glu Thr Leu Ala Thr Phe Val Lys
 275 280 285

Trp Gln Asp Asp Thr Gly Pro Pro Met Asp Lys Ser Asp Leu Gly Gln
 290 295 300

Lys Arg Thr Ser Gly Ala Val Cys His Gln Asp Pro Arg Thr Cys Glu
 305 310 315 320

Glu Pro Ala Ser Ser Gly Ala His Ile Trp Pro Asp Asp Ile Thr Lys
 325 330 335

Trp Pro Ile Cys Thr Glu Gln Ala Arg Ser Asn His Thr Gly Phe Leu
 340 345 350

His Met Asp Cys Glu Ile Lys Gly Arg Pro Cys Cys Ile Gly Thr Lys
 355 360 365

Gly Ser Cys Glu Ile Thr Thr Arg Glu Tyr Cys Glu Phe Met His Gly
 370 375 380

Tyr Phe His Glu Glu Ala Thr Leu Cys Ser Gln Val His Cys Leu Asp
 385 390 395 400

Lys Val Cys Gly Leu Leu Pro Phe Leu Asn Pro Glu Val Pro Asp Gln
 405 410 415

Phe Tyr Arg Leu Trp Leu Ser Leu Phe Leu His Ala Gly Val Val His
 420 425 430

Cys Leu Val Ser Val Val Phe Gln Met Thr Ile Leu Arg Asp Leu Glu
 435 440 445

Lys Leu Ala Gly Trp His Arg Ile Ala Ile Ile Phe Ile Leu Ser Gly
 450 455 460

Ile Thr Gly Asn Leu Ala Ser Ala Ile Phe Leu Pro Tyr Arg Ala Glu
 465 470 475 480

Val Gly Pro Ala Gly Ser Gln Phe Gly Leu Leu Ala Cys Leu Phe Val
 485 490 495

Glu Leu Phe Gln Ser Trp Pro Leu Leu Glu Arg Pro Trp Lys Ala Phe
 500 505 510

Leu Asn Leu Ser Ala Ile Val Leu Phe Leu Phe Ile Cys Gly Leu Leu
 515 520 525

Pro Trp Ile Asp Asn Ile Ala His Ile Phe Gly Phe Leu Ser Gly Leu
 530 535 540

Leu Leu Ala Phe Ala Phe Leu Pro Tyr Ile Thr Phe Gly Thr Ser Asp
 545 550 555 560

Lys Tyr Arg Lys Arg Ala Leu Ile Leu Val Ser Leu Leu Ala Phe Ala
 565 570 575

Gly Leu Phe Ala Ala Leu Val Leu Trp Leu Tyr Ile Tyr Pro Ile Asn
 580 585 590

Trp Pro Trp Ile Glu His Leu Thr Cys Phe Pro Phe Thr Ser Arg Phe
 595 600 605

Cys Glu Lys Tyr Glu Leu Asp Gln Val Leu His
 610 615

<210> 13
 <211> 1305
 <212> DNA
 <213> Homo sapiens

<400> 13
 atggggcggg gcctctggga ggcgtggcct cgggccggct cctctgctgt tgccaagggg 60
 aactgccgcg aggaggcgga aggagcagag gaccggcagc cggcgctcag gcggggcgcg 120
 ggaacgacgg cggccatggc ggcctcgggg cccgggtgtc gcagctggtg cttgtgtccc 180
 gaggtgccat cggccacctt cttcactgcg ctgctctcgc tgctgggttc cgggcctcgc 240
 ctgttctctg tgcagcagcc cctggcgccc tcgggcctca cgctgaagtc cgaggccctt 300
 cgcaactggc aagtttacag gctggtaacc tacatctttg tctacgagaa tcccatctcc 360
 ctgctctgcg gcgctatcat catctggcgc tttgctggca atttcgagag aaccgtgggc 420
 accgtccgcc actgcttctt caccgtgatc ttggccatct tctcgcctat catcttctctg 480
 tcattcgagg ctgtgtcatc actgtcaaag ctgggggaag tggaggatgc cagaggtttc 540
 accccagtgg cctttgccat gctgggagtc accaccgtcc gttctcggat gaggcgggccc 600
 ctggtgtttg gcatggttgt gccctcagtc ctggttccgt ggctcctgct ggggtgcctcg 660
 tggctcattc cccagacctc tttcctcagt aatgtctgcg ggctgtccat cgggctggcc 720
 tatgtcacc tactgtatt ccatcgacct ctcagagcga gtggcgctga agctcgatca 780
 gaccttcccc ttcagcctga tgaggaggat atccgtgttc aagtacgtct cagggtcttc 840
 agccgagagg agggcagccc agagccggaa actgaaccgc gtgcctggct cctacccac 900
 acagagctgc caccctcacc tgtccccaag ccaccctgtg tcccagacgc agcacgccag 960
 tggtcagaag ctggcctcct ggctcctgc acccccgggc acatgccac cttgcctccg 1020
 taccagcctg cctccggcct gtgctatgtg cagaaccact ttggtccaaa cccacctcc 1080
 tccagtgtct acccagcttc tgcgggcacc tccctgggca tccagcccc cagcctgtg 1140
 aacagccctg gcacggtgta ttctggggcc ttgggacacc aggggctgca ggctccaagg 1200
 agtctccag ggtcccatg ccctgagaga atttctaggg aagtcattct acttggcctt 1260
 ctgaaggtcc tccctaagag tctcctgaca aaagttactt attga 1305

<210> 14
 <211> 434
 <212> PRT
 <213> Homo sapiens

<400> 14

Met Gly Arg Gly Leu Trp Glu Ala Trp Pro Pro Ala Gly Ser Ser Ala
 1 5 10 15

Val Ala Lys Gly Asn Cys Arg Glu Glu Ala Glu Gly Ala Glu Asp Arg
 20 25 30
 Gln Pro Ala Ser Arg Arg Gly Ala Gly Thr Thr Ala Ala Met Ala Ala
 35 40 45
 Ser Gly Pro Gly Cys Arg Ser Trp Cys Leu Cys Pro Glu Val Pro Ser
 50 55 60
 Ala Thr Phe Phe Thr Ala Leu Leu Ser Leu Leu Val Ser Gly Pro Arg
 65 70 75 80
 Leu Phe Leu Leu Gln Gln Pro Leu Ala Pro Ser Gly Leu Thr Leu Lys
 85 90 95
 Ser Glu Ala Leu Arg Asn Trp Gln Val Tyr Arg Leu Val Thr Tyr Ile
 100 105 110
 Phe Val Tyr Glu Asn Pro Ile Ser Leu Leu Cys Gly Ala Ile Ile Ile
 115 120 125
 Trp Arg Phe Ala Gly Asn Phe Glu Arg Thr Val Gly Thr Val Arg His
 130 135 140
 Cys Phe Phe Thr Val Ile Phe Ala Ile Phe Ser Ala Ile Ile Phe Leu
 145 150 155 160
 Ser Phe Glu Ala Val Ser Ser Leu Ser Lys Leu Gly Glu Val Glu Asp
 165 170 175
 Ala Arg Gly Phe Thr Pro Val Ala Phe Ala Met Leu Gly Val Thr Thr
 180 185 190
 Val Arg Ser Arg Met Arg Arg Ala Leu Val Phe Gly Met Val Val Pro
 195 200 205
 Ser Val Leu Val Pro Trp Leu Leu Leu Gly Ala Ser Trp Leu Ile Pro
 210 215 220
 Gln Thr Ser Phe Leu Ser Asn Val Cys Gly Leu Ser Ile Gly Leu Ala
 225 230 235 240
 Tyr Ala His Leu Leu Leu Phe His Arg Pro Leu Arg Ala Ser Gly Ala
 245 250 255

Glu Ala Arg Ser Asp Leu Pro Leu Gln Pro Asp Glu Glu Asp Ile Arg
 260 265 270

Val Gln Val Arg Leu Arg Val Phe Ser Arg Glu Glu Gly Ser Pro Glu
 275 280 285

Pro Glu Thr Glu Pro Gly Ala Trp Leu Leu Pro His Thr Glu Leu Pro
 290 295 300

Pro Ser Pro Val Pro Lys Pro Pro Cys Val Pro Asp Ala Ala Arg Gln
 305 310 315 320

Trp Ser Glu Ala Gly Leu Leu Ala Ser Cys Thr Pro Gly His Met Pro
 325 330 335

Thr Leu Pro Pro Tyr Gln Pro Ala Ser Gly Leu Cys Tyr Val Gln Asn
 340 345 350

His Phe Gly Pro Asn Pro Thr Ser Ser Ser Val Tyr Pro Ala Ser Ala
 355 360 365

Gly Thr Ser Leu Gly Ile Gln Pro Pro Thr Pro Val Asn Ser Pro Gly
 370 375 380

Thr Val Tyr Ser Gly Ala Leu Gly His Gln Gly Leu Gln Ala Pro Arg
 385 390 395 400

Ser Pro Pro Gly Ser Pro Cys Pro Glu Arg Ile Ser Arg Glu Val Ile
 405 410 415

Ser Leu Gly Leu Leu Lys Val Leu Pro Lys Ser Leu Leu Thr Lys Val
 420 425 430

Thr Tyr

<210> 15
 <211> 1738
 <212> DNA
 <213> Homo sapiens

<400> 15
 tgctgactaa tccgcgggcc gcggggaatg ggtggcgcg cagccggtcg gcaggggtca 60
 cgcgcccggg tgtgcgcgga atggattcag ggttctcggg tcgcgcccgg gagctagagt 120
 cctgactccc agccgggggc cccgaccggc ctttcggggg tgccggggcg gctctgcaaa 180
 ggggaaactg aggcccaagg aaattgagta tctctgcaaa gtcaccagct gagttcgaac 240

cagacagatc ctgggacccc ctactctatc cgccaccgga agaattctccg tccttgtctt 300
 tccattctgc cctcccggct tcccagtaag cacacagccc caggaccacc cttgaccgtc 360
 ctcaaccaag cgatgcatgc caggggcccc catggccaac tgtccccagc actgcctctg 420
 gcctcctcag tcctgatgct gctgatgagc accctgtggc tgggtgggggc cggccccggc 480
 ctggtcctgg ccccgagct gttgctggac ccctggcagg tgcaccggct gctgacccat 540
 gccctgggcc acacggccct gccaggcctg ctctgagcc tgctgctcct gccactgtg 600
 ggctggcagc aggagtgcc cctgggcacg ctgagattcc tgcattgctc agccctgtc 660
 gccctggctt ctgggtgct ggcagtgtg ctggcaggcc ttgggtgtg cagtgcagcc 720
 ggcagctgtg gatacatgcc tgtccacctg gccatgctgg ctggggaggg acaccgccct 780
 agacggcccc gtggggcact gccaccgtgg ctgtcgccgt ggctgctgct tgccctgacc 840
 ccactgtca gctctgagcc acccttctctg cagctccttt ggggcctcct tgccggcctg 900
 gcctatgcag ctggggcctt ccggtggctg gaacctcag agcgacggct gcaggtgtg 960
 caggagggcg tcttgtgcag gaccttggcg ggggtgtggc ccctgaggct ccttgccacc 1020
 ccgggtagcc tggcggagct gcctgtcacc catcctgccg gagtgaggcc tcccatccct 1080
 ggaccgcctt atgtggcctc ccctgacctc tgggtccact gggaagactc agccctgccc 1140
 ccaccaagcc tgaggcctgt gcagcccacc tgggagggct cctcagaggc aggcctggac 1200
 tgggtgtggg ccagcttctc ccaggggact ccgatgtggg cggccttga tgagcagatg 1260
 ctgcaggagg gcatccaggc ctgcttctt gacgggccag ccagggaacc ccagagcgca 1320
 ccatggctgt ccaagtcctc tgtctcctct ctgcggtgc agcagctgga gcgcatgggc 1380
 ttccctacgg agcaggcggg ggtggcactg gcagccacag gccgtgtgga gggtgccgtg 1440
 tcaactgttg ttggaggaca agtgggcact gagaccctgg tgacccatgg aaaggtggg 1500
 cctgcccact ccgaggggtc tgggcctccc tagccaggc agagagtggg gcacaggcag 1560
 gcccttgggt gctaagggct gggctgcatg tgggtagccc gagctcctac tctgtctaaa 1620
 gagggccaca gtggggagca ggggcacctc tggaggcagg agaggcccc cagcatgctg 1680
 ccctagtacg tgtttagaat aaaaaccagt ttgtttttca acctggacct ccttgag 1738

<210> 16

<211> 386

<212> PRT

<213> Homo sapiens

<400> 16

Met His Ala Arg Gly Pro His Gly Gln Leu Ser Pro Ala Leu Pro Leu

1

5

10

15

Ala Ser Ser Val Leu Met Leu Leu Met Ser Thr Leu Trp Leu Val Gly
 20 25 30

Ala Gly Pro Gly Leu Val Leu Ala Pro Glu Leu Leu Leu Asp Pro Trp
 35 40 45

Gln Val His Arg Leu Leu Thr His Ala Leu Gly His Thr Ala Leu Pro
 50 55 60

Gly Leu Leu Leu Ser Leu Leu Leu Leu Pro Thr Val Gly Trp Gln Gln
 65 70 75 80

Glu Cys His Leu Gly Thr Leu Arg Phe Leu His Ala Ser Ala Leu Leu
 85 90 95

Ala Leu Ala Ser Gly Leu Leu Ala Val Leu Leu Ala Gly Leu Gly Leu
 100 105 110

Ser Ser Ala Ala Gly Ser Cys Gly Tyr Met Pro Val His Leu Ala Met
 115 120 125

Leu Ala Gly Glu Gly His Arg Pro Arg Arg Pro Arg Gly Ala Leu Pro
 130 135 140

Pro Trp Leu Ser Pro Trp Leu Leu Leu Ala Leu Thr Pro Leu Leu Ser
 145 150 155 160

Ser Glu Pro Pro Phe Leu Gln Leu Leu Cys Gly Leu Leu Ala Gly Leu
 165 170 175

Ala Tyr Ala Ala Gly Ala Phe Arg Trp Leu Glu Pro Ser Glu Arg Arg
 180 185 190

Leu Gln Val Leu Gln Glu Gly Val Leu Cys Arg Thr Leu Ala Gly Cys
 195 200 205

Trp Pro Leu Arg Leu Leu Ala Thr Pro Gly Ser Leu Ala Glu Leu Pro
 210 215 220

Val Thr His Pro Ala Gly Val Arg Pro Pro Ile Pro Gly Pro Pro Tyr
 225 230 235 240

Val Ala Ser Pro Asp Leu Trp Ser His Trp Glu Asp Ser Ala Leu Pro
 245 250 255

Pro Pro Ser Leu Arg Pro Val Gln Pro Thr Trp Glu Gly Ser Ser Glu
 260 265 270

Ala Gly Leu Asp Trp Ala Gly Ala Ser Phe Ser Pro Gly Thr Pro Met
 275 280 285

Trp Ala Ala Leu Asp Glu Gln Met Leu Gln Glu Gly Ile Gln Ala Ser
 290 295 300

Leu Leu Asp Gly Pro Ala Gln Glu Pro Gln Ser Ala Pro Trp Leu Ser
 305 310 315 320

Lys Ser Ser Val Ser Ser Leu Arg Leu Gln Gln Leu Glu Arg Met Gly
 325 330 335

Phe Pro Thr Glu Gln Ala Val Val Ala Leu Ala Ala Thr Gly Arg Val
 340 345 350

Glu Gly Ala Val Ser Leu Leu Val Gly Gly Gln Val Gly Thr Glu Thr
 355 360 365

Leu Val Thr His Gly Lys Gly Gly Pro Ala His Ser Glu Gly Pro Gly
 370 375 380

Pro Pro
 385

<210> 17
 <211> 19
 <212> PRT
 <213> Homo sapiens

<400> 17

Met Gly Arg Val Glu Asp Gly Gly Thr Thr Glu Glu Leu Glu Asp Trp
 1 5 10 15

Asp Pro Gly

<210> 18
 <211> 1559
 <212> DNA
 <213> Homo sapiens

<400> 18
 cttggacctt ggccctcgct ttccaggatg ggtagggtgg aagacggggg aacaactgag 60
 gagctggagg actgggaccc aggcaccagt gccctgccag ctctgggat caagcagggt 120
 occaggggaac agacaggcac gggggccctg tcccaaaagt gctgggagcc tgagcctgat 180
 gctcccagcc agcctggccc agccctttgg tccaggggtc gggcccgcac tcaggccttg 240

gctggcggct cctcactgca gcagctggac cccgagaaca caggcttcat cgggtgcggac 300
 accttactg gcctggtgca cagccatgag ctgcccctgg acccggccaa gctggacatg 360
 ctggtggccc tggctcagag caacgagcag ggccaggtct gctaccagga gctggtggac 420
 ctgatcagca gcaagcgctc cagcagtttc aagcgggcca ttgctaacgg acagcgggca 480
 ctgccccggg acgggccgct ggatgagcca ggccatagtg tctacaagcg gtttgtgcgt 540
 tacgtggcct acgagatcct gccttgtgag gtggaccgcc gctgggtactt ctaccgtcac 600
 cgcagctgcc cccccccggt gttcatggcc tcggtcactc ttgccagat catcgtgttc 660
 ctgtgttacg gggcccgcct caacaagtgg gtgctgcaga cctaccaccc cgagtacatg 720
 aagagcccc ttgtgtacca ccccgggcac cgtgcccgcg cctggcgctt cctcacctac 780
 atgttcatgc acgttgggct ggagcagctg gggttcaacg ccctcctgca gctgatgatc 840
 ggggtgcccc tggagatggg gcacggcctg ctccgcatca gcctgctcta cctggcaggc 900
 gtgctggcag gctccctaac cgtctccatc accgacatgc gggccccggg ggtgggaggc 960
 tccggcgggg tctacgccct gtgctcggca cacctggcca acgttgtcat gaactgggct 1020
 gggatgagat gtcctacaa gttgctgagg atggtgctgg ccttgggtgtg catgagctcc 1080
 gaggtgggcc gggccgtgtg gctgcgcttc tccccgccgc tgccgcctc gggcccacag 1140
 cccagcttca tggcgcacct ggcaggcgcg gtggtggggg tgagcatggg cctgaccatc 1200
 ctgcggagct acgaggagcg cctgcgggac cagtgcggct ggtgggtggt gctgctggcc 1260
 tacggcacct tcctgctctt cgccgtcttc tggaaagtct tcgcctacga cctgctgggc 1320
 gccacatcc cccaccgcc ctgaccggct acctgaggct gcacaggcca gggctcgggc 1380
 atgtggtggc cgcccaccag gggccttcac gtctgccctt tgtgaacgga cgtctcaggg 1440
 ctgctgtgcc ccttgggtgt ggggtggcctc aaaggaggcc ctgtcccagc caccacccc 1500
 ccactcccag gacttgcggt ctgagccttt ttggataatt aataaatatt ttacacagc 1559

<210> 19
 <211> 593
 <212> DNA
 <213> Homo sapiens

<400> 19
 atttactatg ctgtgtggaa gcctcagaaa cagtggatca cgttggacac aggcattctg 60
 gagagtcctt ttatctacag tcctgagaag agggaggaag cctggagggt tatctcatc 120
 atgtgtgttac atgctggagt tcagcacatc ttggggaatc tttgtatgca gcttgttttg 180
 ggtattccct tggaaatggt ccacaaaggc ctccgtgtgg ggctgggtgta cctggcagga 240
 gtgattgcag ggtcccttgc cagctccatc tttgaccac tcagatatct tgtgggagct 300


```

tcaggaggag tctatgctct gatgggaggc tattttatga atgttctggt gaattttcaa 360
gaaatgattc ctgccttttg aattttcaga ctgctgatca tcatcctgat aattgtgttg 420
gacatgggat ttgctctcta tagaagggtc tttgttcctg aagatgggtc tccggtgtct 480
tttgagctc acattgcagg tggatttgct ggaatgtcca ttggctacac ggtgttttagc 540
tgctttgata aagcactgct gaaagatcca aggttatgga tagcaattgc tgc 593

```

```

<210> 20
<211> 708
<212> DNA
<213> Homo sapiens

```

```

<400> 20
aaggaggagc gcctggcaga gcgggatctt cagggcagcg ggcaaccct tggcccagga 60
agcctggaac gcaaggaccg gagggcgtgg gctgggacgc ccctacgttg gtctttcagg 120
gaaaggcctt ggaaagcagt cgttgcgcca gacagcccag ggaagagcgg cagcctgagg 180
acctagggcc acctgctgtt ccctgggatt catgtccttc tggggaggag ggaggacca 240
ggacaatggc tgctgttcat gatctggaga tggagagcat gaatctgaat atggggagag 300
agatgaaaga agagctggag gaagaggaga aaatgagaga ggatggggga ggtaaagatc 360
gggccaagag taaaaaggtc cacaggattg tctcaaatg gatgctgcc gaaaagtccc 420
gaggaacata cttggagaga gctaactgct tcccgcctcc cgtgttcac atctccatca 480
gcctggccga gctggcagtg tttatttact atgctgtgtg gaagcctcag aaacagtgga 540
tcacgttggg cacaggcatc ttggagagtc cctttatcta cagtcctgag aagagggagg 600
aagcctggag gtttatctca tacatgctgg tacatgctgg gtaagcaatg atagttaagc 660
cctggtatca gaggtgatta tatcattgta acctcttaaa acaaaaaa 708

```

```

<210> 21
<211> 616
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (498)..(498)
<223> n is a, c, g, or t

```

```

<400> 21
cttcttccct ttctacatta agaagcccct tccttaaate ctaaggctct tttataggca 60
atctttgcaa ctgatgagtt taatgctgtt ttgtcctcta tataaaattg ttagcctttg 120
ctgagacttc tctgtttctt catagaggct tccctttttt tttttatttc ctccagatgg 180
ctctttttat taattgactt acaatagggg caggtcagtt tgctggagat aggaaaatgt 240

```

tgaaaaaacac agcaaataag acacaagcta aatatgcagc aattgctatc caaaaccttg 300
 gatctttcat cagtgtcttta tcaaagcagc taaacaccgt gtagccaatg gacattccag 360
 caaatccacc tgcaatgtga gctgcaaaag acaccggaga cccatcttca ggaacaaaga 420
 accttctata gagagcaaat cccatgtcca acacaattat caggatgatg atcagcagtc 480
 tgaaaattcc aaaggcanga atcatttctt gaaaattcac cagaacattc ataaaaatagc 540
 ctcccatcag agcatagact cctcctgaag ctcccacaag atatctgagt gggtaaaga 600
 tggagctggc aaggga 616

<210> 22
 <211> 354
 <212> DNA
 <213> Homo sapiens

<400> 22
 tggatcacgt tggacacagg catttggaga gtccctttat ctacagtcct gagaagaggg 60
 aggaagcctg gaggtttatc tcacacatgc tggatcatgc tggagttcag cacatcttgg 120
 ggaatctttg tatgcagctt gttttgggta ttcccttggg aatgggtccac aaaggcctcc 180
 gtgtgtgggct ggtgtacctg gcaggagtga ttgcagggtc ccttgccagc tccatctttg 240
 acccactcag atatcttgtg ggagcttcag gaggagtcta tgctctgatg ggaggctatt 300
 ttatgaatgt tctgggtgaat tttcaagaaa tgattcctgc ctttgggaatt ttca 354

<210> 23
 <211> 356
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (356)..(356)
 <223> n is a, c, g, or t

<400> 23
 tttaaaacaa ttacagagaa aactttatct tgggccatct aggaggttta gatcattttg 60
 atcatcttca gctgtcttct cttcacatagc aggaaggcc ttggaaagca gtcgttgccg 120
 cagacagccc aggaagagc ggcacgctga ggacctaggc ccacctgctg ttccctggga 180
 ttcatgtcct tctggggagg agggaggacc caggacaatg gctgctgttc atgatctgga 240
 gatggagagc atgaatctga atatggggag agagatgaaa gaagagctgg aggaagagga 300
 gaaaatgaga gaggatgggg gaggtaaaga tcgggccaag agtaaaaagg tccacn 356

<210> 24
 <211> 524
 <212> DNA

<213> Homo sapiens

<400> 24

```

gggtctccgg tgtcttttgc agctcacatt gcagggtggat ttgctggaat gtccattggc      60
tacacgggtgt ttagctgctt tgataaagca ctgatgaaag atccaagggt ttggatagca      120
attgctgcat atttagcttg tgtcttattt gctgtgtttt tcaacatttt cctatctcca      180
gcaaactgac ctgcccctat tgtaagtcaa ttaataaaaa gagccatctg gaggaaataa      240
aaaaaaaaagg aagactctat gaagaaacag agaagtctca gcaaaggcta acaattttat      300
atagaggaca aaacagcatt aaactcatca gttgcaaaga ttgcctataa aaggacctta      360
ggatttaagg aaggggcttc ttaatgtaga aagggaagaa gaagagagaa agaagggtta      420
gtaaaaacta gagattgggg ccaggcgcag tggctcacgc ctgtaatccc agcactttgg      480
gaggctgagg cgggtgaaat cacctgaggt catgagttca agac                        524

```

<210> 25

<211> 428

<212> DNA

<213> Homo sapiens

<400> 25

```

ttttttttcc ctttctacat taagaagccc cttccttaaa tcctaaggtc cttttatagg      60
caatctttgc aactgatgag ttaatgctg ttttgtctc tatataaaat tgtagcctt      120
ttctgagact tctctgtttc ttcataagat cttccttttt tttattattt cctccagatg      180
gctcttttta ttaattgact tacaataggg gcaggtcagt ttgctggaga taggaaaatg      240
ttgaaaaaca cagcaaataa gacacaagct aaatatgcag caattgctat ccaaacctt      300
ggatctttca gcagtgcctt atcaaagcag ctaaaccaccg tgtagccaat ggacattcca      360
gcaaatccac ctgcaatgtg agctgcaaaa gacaccttgg gagaaaagga gggaaaatgg      420
aaataagt                                         428

```

<210> 26

<211> 271

<212> DNA

<213> Homo sapiens

<400> 26

```

ctcgaaactg tgaaaacaca gcaaataaga cacaagctaa atatgcagca attgctatcc      60
aaaaccttgg atctttcagc agtgctttat caaagcagct aaacaccgtg tagccaatgg      120
acattccagc aaatccacct gcaatgtgag ctgcaaaaga cacctgggag aaaaggaggg      180
aaaatggaaa tcagtgaaga caatgctaatt tgtgtatttc agttgcatct ctcttattca      240
aacacaaata ccgtcacagt taaacaagcg t                                         271

```

<210> 27
 <211> 262
 <212> DNA
 <213> Homo sapiens

<400> 27
 tgaaatacac agcaaataga cacaagctaa atatgcagca attgctatcc aaaagcttgg 60
 atcttttcagc agtgctttat caaagcagct aaacaccgtg tagcgcatgg acattccagc 120
 aaatccacct gcaatgtgag ctgcaaaaga caccttggga gaaaaggagg gaaaatggaa 180
 ataagtgaag acaatgctaa ttgtgtatct cagttgcac tctcttattc aaacacaaat 240
 accgtcacag ttaaacaagc gt 262

<210> 28
 <211> 120
 <212> DNA
 <213> Homo sapiens

<400> 28
 gctcgggtccg aaactgttga aaaacacagc aaataagaca caacgctaaa tatgcagcaa 60
 ttgctatcca aaaccttggga tctttcagca gtgctttatc aaagcagcta aacaccgtgt 120

<210> 29
 <211> 894
 <212> DNA
 <213> Homo sapiens

<400> 29
 ctcatccacg ttacttgaag aactccctgg tttaccaccc acagctgca gcacaggttt 60
 ggcgctacct gacatacatc ttcatgcatg cagggtatag aacacctgga ctcaatgtgg 120
 tgctgcagct gtggtggggg tgcccctgga gatggtgcat ggagccaccc gaattgggct 180
 tgtctacgtg gccgggtgtg tggcaggggc cttggcagtg tctagtggct gacatgaccg 240
 ctccagtcgt gggctcttct ggaggggtgt atgtctcgt ctctgcccac ctggccaaca 300
 ttgtcatgaa ctggtcaggc atgaagtgcc agttcaagct gctgcggatg gctgtggccc 360
 ttatctgtat gagcatggag cttgggcggg ccgtgtggct ccgcttcac ccgtcggcta 420
 tccccgtgc cctcaccaca gctttgtggc gcaattgggt ggcgtggccg tgtgcatcac 480
 cctggccgtg gtggtcctga ggaactacga gcagaggctc caggaccagt cactgtgggtg 540
 gatttttgtg gccatgtaca ccgtcttcgg gctgttgcgt gtctctggaa catctttggc 600
 tacacctgtt ggacttaaag ctgcgctccc ccctgaggct tgaggcccca gtccgcgagg 660
 ggagggaaag caggacccca gggagcgctt ggaagggtct tcctcatcac aggttcagta 720
 agcgggcaaa cacgcacaaa cactgcgcct gtattgtgtt atttggccac gggggcacct 780
 ctgcgaaggg ctgcgggcga aacacaagac aaacggacta acacaagaac aaggggcgaa 840

tcatcgcgcg ggagcacttg gaaagcaggt taaaacggga cacaaggaca cttt 894

<210> 30
 <211> 344
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (19)..(19)
 <223> n is a, c, g, or t

<220>
 <221> misc_feature
 <222> (49)..(49)
 <223> n is a, c, g, or t

<220>
 <221> misc_feature
 <222> (92)..(92)
 <223> n is a, c, g, or t

<220>
 <221> misc_feature
 <222> (130)..(130)
 <223> n is a, c, g, or t

<400> 30
 cctccgccag atgcctttna gaaaaggggc tccactgtgt gtccaaatnt aaacacaaac 60
 attacagtgg cccagagtct tctgtccttg tntgcccggc ctagctgagc tgatgatgag 120
 aagaaacctn gcaggcgctc cctgtgggtg ctgcttttcc ctccccctccc cgaccttggg 180
 cctccagccc tcagggggga ggccgcagct ttaagtccag caggggtgtag gcaaagatgt 240
 tccagaagac agcgaacagc acgaagacgg tgtacatggc cacaaaaatc caccacagtg 300
 actggctctg gagcctctgc tcgtagttcc tcaggaccac cacg 344

<210> 31
 <211> 1140
 <212> DNA
 <213> Homo sapiens

<400> 31
 atggcggtggc gaggctgggc gcagagaggc tggggctgcg gccaggcggtg ggggtgcgtcg 60
 gtggggcgcc gcagctgcga ggagctcact gcggtcctaa ccccgccgca gtcctcggga 120
 cgcaggttta acttctttat tcaacaaaaa tgcggattca gaaaagcacc caggaagggtt 180
 gaacctcgaa gatcagaccc agggacaagt ggtgaagcat acaagagaag tgctttgatt 240
 cctcctgtgg aagaaacagt cttttatcct tctccctatc ctataaggag tctcataaaa 300
 cctttatattt ttactgttgg gtttacaggc tgtgcatttg gatcagctgc tatttggcaa 360
 tatgaatcac tgaaatccag ggtccagagt tattttgatg gtataaaagc tgattgggtg 420

gatagcataa gaccacaaaa agaaggagac ttcagaaagg agattaacaa gtgggtggaat 480
aacctaagtg atggccagcg gactgtgaca ggtattatag ctgcaaagt ccttgtattc 540
tgtttatgga gagtaccttc tctgcagcgg acaatgatca gatatttcac atcgaatcca 600
gcctcaaagg tcctttgttc tccaatgttg ctgtcaacat tcagtcactt ctccttattt 660
cacatggcag caaatatgta tgttttgtgg agcttctctt ccagcatagt gaacattctg 720
ggatcaagagc agttcatggc agtgtaccta tctgcaggtg ttatttccaa ttttgtcagt 780
tacctgggta aagttgccac aggaagatat ggaccatcac ttggtgcac tggtgccac 840
atgacagtcc tcgcagctgt ctgcactaag atcccagaag ggaggcttgc cattattttc 900
cttccgatgt tcacgttcac agcagggaaat gccctgaaag ccattatcgc catggataca 960
gcaggaatga tcctgggatg gaaatttttt gatcatgcgg cacatcttgg gggagctctt 1020
tttggaatat ggtatgttac ttacggtcat gaactgattt ggaagaacag ggagccgcta 1080
gtgaaaatct ggcatgaaat aaggactaat ggccccaaaa aaggaggtgg ctctaagtaa 1140

<210> 32
<211> 2964
<212> DNA
<213> Homo sapiens

<400> 32
gactcggcgc gggcgccctc cgggagcgc ggggagccc ctctctcccgc gcgcccctcag 60
gacccccag agacccccgc cggcggcagc ctgccttgct ctgccaggaa ccatgagtga 120
ggcccgagc gacagcacga gcagcctgca gcgcaagaag ccacctggc taaagctgga 180
cattccctct gcggtgcccc tgacggcaga agagcccagc ttcttgagc ccctgaggcg 240
acaggctttc ctgaggagtg tgagtatgcc agccgagaca gcccacatct cttcacccca 300
ccatgagctc cggcgggccgc tgctgcaacg ccagacgtcc atcacacaga ccatccgcag 360
ggggaccgcc gactggtttg gactgagcaa ggacagtgc agcaccaga aatggcagcg 420
caagagcatc cgtcactgca gccagcgcta cgggaagctg aagccccagg tcctccggga 480
gctggacctg ccagccagg acaacgtgtc gctgaccagc accgagacgc cccccact 540
ctacgtgggg ccatgccagc tgggcatgca gaagatcata gacccctgg cccgtggccg 600
tgcttccgt gtggcagatg aactgcgga aggcctgagt gcccacaca ctccgtcac 660
gccgggtgct gcctccctct gctccttctc cagctcccgc tcagggttcc accggtccc 720
gcggcggcgc aagcgagagt cgggtggcaa gatgagcttc cggcgggccg cagcgctgat 780
gaaaggccgc tccgttaggg atggcacctt tcgcccggca cagcgctgaa gcttcaactc 840
agctagcttt ctggaggagg acacaactga tttccccgat gagctggaca catccttctt 900

tgccccgggaa ggtatcctcc atgaagagct gtccacatac ccggatgaag ttttcgagtc	960
cccatcggag gcagcgctaa aggactggga gaaggcaccg gagcaggcgg acctcaccgg	1020
cggggccctg gaccgcagcg agcttgagcg cagccacctg atgctgccct tggagcgagg	1080
ctggcggaag cagaaggagg gcgccgcagc cccgcagccc aagggtgcggc tccgacagga	1140
ggtggtgagc accgcggggc cgcgacgggg ccagcgtatc gcggtgccgg tgcgcaagct	1200
cttcgcccgg gagaagcggc cgtatgggct gggcatggtg ggacggctca ccaaccgcac	1260
ctaccgcaag cgcatcgaca gcttcgtcaa gcgccagatc gaggacatgg acgaccacag	1320
gcccttcttc acctactggc ttaccttctg gcactcgctc gtcgccatcc tagccgtgtg	1380
catctatggc atcgcgcccg tgggcttctc gcagcatgag acggtggact cgggtgctgcg	1440
gaaccgcggg gtctacgaga acgtcaagta cgtgcagcag gagaacttct ggatcgggcc	1500
cagctcggag gccctcatcc acctggggcg caagttttcg ccctgcatgc gccaggaccc	1560
gcagggtcac agcttcattc gctcggcgcg cgagcgcgag aagcactccg cctgctgcgt	1620
gcgcaacgac aggtcgggct gcgtgcagac ctcggaggag gagtgtctgt ccacgctggc	1680
agtgtgggtg aagtggccca tccatcccag cgcccagag cttgcgggcc acaagagaca	1740
gtttggctct gtctgccacc aggatcccag ggtgtgtgat gagccctcct ccgaagaccc	1800
tcatgagtgg ccagaagaca tcaccaagtg gccgatctgc accaaaaaca gcgctgggaa	1860
ccacaccaac catccccaca tggactgtgt catcacagga cggccctgct gcattggcac	1920
caagggcagg tgtgagatca cctcccggga gtactgtgac ttcatgaggg gctacttcca	1980
tgaggaggcc acgctctgct ctcagggtgca ctgcatggat gatgtgtgtg ggctcctgcc	2040
ttttctcaac cccgaggtgc ctgaccagtt ctaccgcctg tggctatccc tcttcctgca	2100
cgcggggtc ttgcaactgc tgggtgtccat ctgcttccag atgactgtcc tgcgggacct	2160
ggagaagctg gcaggctggc accgcatagc catcatctac ctgctgagtg gtgtcaccgg	2220
caacctggcc agtgccatct tcctgccata ccgagcagag gtgggtcctg ctggctccca	2280
gttcggcatc ctggcctgcc tcttcgtgga gctcttccag agctggcaga tcctggcgcg	2340
gccctggcgt gccttcttca agctgctggc tgtggtgctc ttcctcttca cctttgggct	2400
gctgccgtgg attgacaact ttgccacat ctcggggttc atcagtggcc tcttcctctc	2460
cttcgccttc ttgccctaca tcagcttttg caagtctgac ctgtaccgga aacgctgcca	2520
gatcatcatc tttcagggtg tcttctctgg cctcctggct ggctggtgg tcctcttcta	2580
cgtctatcct gtccgctgtg agtggtgtga gttcctcacc tgcacccct tactgacaa	2640
gttctgtgag aagtacgaac tggacgctca gctccactga gctggctgcg ggctccagcg	2700
gccgtgtgct ccagcaggcc agagccagac acgacctccc tgagcctcac aggccttacg	2760

gagtcacctg ctccatgtgg ggactggcct gtttcctgaa cacagacctc tttcttgtgc 2820
 cttgttcact tctgttgaac ccctcgact gccgggcatt tattatacta cttcctgtca 2880
 taaccttcta acttgtttct tgacgaccac ctcatgtggc caataaatga actgggagcg 2940
 ttttaaaaaa aaaaaaaaaa aaaa 2964

<210> 33
 <211> 2643
 <212> DNA
 <213> Homo sapiens

<400> 33
 gctactccca cctgccacgc cgcaagagaa tgtctgtggc ccacatgagc ttgcaagctg 60
 ccgctgccct cctcaagggg cgctcgggtgc tggatgccac cggacagcgg tgccgggtgg 120
 tcaagcgcag ctttgccctc ccgagcttcc tggaggagga tgtggtcgat ggggcagaca 180
 cgtttgactc ctcccttttt agtaaggaag aaatgagctc catgcctgat gatgtctttg 240
 agtccccccc actctctgcc agctacttcc gagggatccc aactcagcc tcccctgtct 300
 cccccgatgg ggtgcaaata cctctgaagg agtatggcgg agccccagtc cccgggcccc 360
 ggcgcgga caa ggcacatgcc tccaagggtga agcactttgc ctttgatcgg aagaagcggc 420
 actacggcct cggcgtggtg ggcaactggc tgaaccgcag ctaccgccc agcatcagca 480
 gcactgtgca gcgcagctg gagagcttcg acagccaccg gccctacttc acctactggc 540
 tgaccttcgt ccatgtcatc atcacgctgc tgggtgattg cacgtatggc atcgcacccg 600
 tgggctttgc ccagcacgtc accaccagc tgggtgctgg gaacaaaggt gtgtacgaga 660
 gcgtgaagta catccagcag gagaacttct ggggtggccc cagctcgatt gacctgatcc 720
 acctgggggc caagttctca ccctgcatcc ggaaggacgg gcagatcgag cagctggtgc 780
 tgccgcgagc agacctggag cgggactcag gctgctgtgt ccagaatgac cactccggat 840
 gcatccagac ccagcgggaag gactgctcgg agactttggc cacttttgtc aagtggcagg 900
 atgacactgg gcccccatg gacaagtctg atctgggcca gaagcggact tcgggggctg 960
 tctgccacca ggaccccagg acctgcgagg agccagcctc cagcggtgcc cacatctggc 1020
 ccgatgacat cactaagtgg ccgatctgca cagagcaggc caggagcaac cacacaggct 1080
 tcctgcacat ggactgcgag atcaagggcc gccctgctg catcggcacc aagggcagct 1140
 gtgagatcac caccgggaa tactgtgagt tcatgcacgg ctatttccat gaggaagcaa 1200
 cactctgtc ccaggtgcac tgcttgga ca aggtgtgtgg gctgctgccc ttcctcaacc 1260
 ctgagggtccc agatcagttc tacaggctct ggctgtctct cttcctacat gctggcgtgg 1320
 tgcactgcct cgtgtctgtg gtctttcaaa tgaccatcct gagggacctg gagaagctgg 1380
 ccggctggca ccgtatcgcc atcatcttca tcctcagtgg catcacaggc aacctcgcca 1440


```

gtgccatctt tctcccatat cgggcagagg tgggcccggc cggctcacag ttccggcctcc 1500
tcgectgcct ctctcgtggag ctcttcacaga gctggccgct gctggagagg ccctggaagg 1560
ccttcctcaa cctctcggcc atcgtgctct tctgttcat ctgtggcctc ttgccttgga 1620
tcgacaacat cggccacatc ttccggttcc tcagtggcct gctgctggcc ttgccttcc 1680
tgccctacat caccttcggc accagcgaca agtaccgaa gcgggcactc atcctggtgt 1740
cactgctggc ctttgccggc ctcttcgccc ccctcgtgct gtggctgtac atctaccca 1800
ttaactggcc ctggatcgag cacctcacct gcttccccct caccagccgc ttctgcgaga 1860
agtatgagct ggaccagggt ctgcactgac cgctgggcca cagggtgcc cctcagccct 1920
gctggaacag ggtctgcctg cgagggtgc cctctgcaga gcgctctctg tgtgccagag 1980
agccagagac ccaagacagg gcccgggctc tggacctggg tgccccctg ccaggcgagg 2040
ctgactccgc gtgagatggg tgggtaaggc ggggttttcc tggggcgta ggctgtgag 2100
atcctgacct aagctcaggc acaccaagg cacctgcctc tctgagtctt gggctctcagt 2160
tcctaataat ccgctccttg ctgagaccat ctctggggc agggctcctt tcttcccagg 2220
tcctcagcgc tgcctctgct ggtgccttct ccccaactac tactggagcg tgcccttgct 2280
ggggacgtgg ctgtgccctc agttgcccc agggctgggt gccaccatg ccccttctc 2340
tttctcctcc tacctctgcc ctgtgagccc atccataagg ctctcagatg ggacattgta 2400
ggaaaggctt tggccatggg ctgggggcag agaacaaggg gggagacaca agtagacctc 2460
aggtagaacg acaccgggcg gagccacccc agggctgct cccaggaggt gctcgaggcg 2520
catcaggccc gttttttacc agtttatatc acggctctca tttttaaaag taacgctaac 2580
tttgtacgga cgatgtctca tggattaaat aatattcttt atggcaaaaa aaaaaaaaaa 2640
aaa 2643

```

```

<210> 34
<211> 1305
<212> DNA
<213> Homo sapiens

```

```

<400> 34
atggggcggg gcctctggga ggcgtggcct ccggccggct cctctgctgt tgccaaggga 60
aactgccgcg aggaggcgga aggagcagag gaccggcagc cggcgctcag gcggggcgcg 120
ggaacgacgg cggccatggc ggcctcgggg cccgggtgtc gcagctgggt cttgtgtccc 180
gaggtgccat ccgccacctt cttcactgcg ctgctctcgc tgctggtttc cgggcctcgc 240
ctgttctctg tcgagcagcc cctggcgccc tcgggcctca cgctgaagtc cgaggccctt 300
cgcaactggc aagtttacag gctggttaacc tacatctttg tctacgagaa tcccatctcc 360

```

ctgctctgcg ggcctatcat catctggcgc tttgctggca atttcgagag aaccgtgggc	420
accgtccgcc actgcttctt caccgtgac ttcgccatct tctccgctat catcttcttg	480
tcattcgagg ctgtgtcatc actgtcaaag ctgggggaag tggaggatgc cagaggtttc	540
accccgatgg cctttgccat gctgggagtc accaccgtcc gttctcggat gaggcgggcc	600
ctgggtgtttg gcatgggtgt gccctcagtc ctgggtccgt ggctcctgct gggtgccctg	660
tggctcattc cccagacctc tttcctcagt aatgtctgcg ggctgtccat cgggctggcc	720
tatgctcacc tactgctatt ccacgcacct ctcagagcga gtggcgctga agctcgatca	780
gaccttcccc ttcagcctga tgaggaggat atccgtgttc aagtacgtct cagggtcttc	840
agccgagagg agggcagccc agagccgga actgaacccg gtgcctggct cctacccac	900
acagagctgc caccctcacc tgtccccaag ccaccctgtg tcccagacgc agcacgccag	960
tggtcagaag ctggcctcct ggctcctgc accccgggc acatgccac cttgcctcgg	1020
taccagcctg cctccggcct gtgctatgtg cagaaccact ttggtccaaa cccacctcc	1080
tccagtgtct acccagcttc tgccggcacc tccctgggca tccagcccc cagcctgtg	1140
aacagccctg gcacggtgta ttctggggcc ttgggacacc aggggctgca ggctccaagg	1200
agtctccag ggtcccatg ccctgagaga atttctaggg aagtcattc acttggcctt	1260
ctgaaggctc tccctaagag tctcctgaca aaagttactt attga	1305

<210> 35

<211> 1738

<212> DNA

<213> Homo sapiens

<400> 35

tgctgactaa tccgcgggcc gcggggaatg ggtggcgcg cagccggctg gcaggggtca	60
cgcgccggg tgtgcgcgga atggattcag ggttctcggg tcgcgcccgg gagctagagt	120
cctgactccc agccggggtc cccgaccggc ctttcggggg tgccggggcg gctctgcaaa	180
ggggaaactg agggccaagg aaattgagta tctctgcaaa gtcaccagct gagttcgaac	240
cagacagatc ctgggacccc ctactctatc cgccaccgga agaattctcg tccttgtctt	300
tccattctgc cctccgggct tcccagtaag cacacagccc caggaccacc cttgaccgtc	360
ctcaaccaag cgatgcatgc caggggcccc catggccaac tgtcccagc actgcctctg	420
gcctcctcag tctgatgct gctgatgagc accctgtggc tgggtggggc cggccccggc	480
ctggctcctg ccccgagct gttgctggac ccctggcagg tgcaccggct gctgacccat	540
gccctgggcc acacggccct gccaggcctg ctcctgagcc tgctgtcct gccactgtg	600
ggctggcagc aggagtcca cctgggcacg ctgagattcc tgcatgcctc agccctgtc	660
gccctggctt ctgggctgct ggcagtgtg ctggcaggcc ttgggctgtc cagtgcagcc	720

ggcagctgtg gatacatgcc tgtccacctg gccatgctgg ctggggaggg acaccgccct 780
 agacggcccc gtggggcact gccaccgtgg ctgtcgccgt ggctgctgct tgccctgacc 840
 ccactgctca gctctgagcc acccttcctg cagctccttt gcggcctcct tgccggcctg 900
 gcctatgcag ctggggcctt cgggtggctg gaaccctcag agcgacggct gcaggtgctg 960
 caggagggcg tcttgtgcag gaccttggcg ggggtgctggc ccctgaggct ccttgccacc 1020
 ccgggtagcc tggcggagct gcctgtcacc catcctgccg gagtgaggcc tcccatccct 1080
 ggaccgcctt atgtggcctc ccctgacctc tgggtccact gggaagactc agccctgcc 1140
 ccaccaagcc tgaggcctgt gcagcccacc tgggagggct cctcagaggc aggcctggac 1200
 tgggctgggg ccagcttctc ccagggact ccgatgtggg cggccttga tgagcagatg 1260
 ctgcaggagg gcatccaggc ctcgcttctt gacgggccag cccaggaacc ccagagcgca 1320
 ccatggctgt ccaagtcctc tgtctcctct ctggggctgc agcagctgga gcgcatgggc 1380
 ttccctacgg agcaggcggg ggtggcactg gcagccacag gccgtgtgga gggtgccgtg 1440
 tcactgttgg ttggaggaca agtgggcact gagaccctgg tgacccatgg aaagggtggg 1500
 cctgcccact ccgaggggtc tgggcctccc tagcccaggc agagagtggg gcacaggcag 1560
 gcccttgggt gctaagggct gggtgcatg tgggtagccc gagctcctac tctgtctaaa 1620
 gagggccaca gtggggagca ggggcacctc tggaggcagg agaggccccc cagcatgctg 1680
 ccctagtacg tgtttagaat aaaaaccagt ttgtttttca acctggacct ccttgagg 1738

<210> 36
 <211> 438
 <212> PRT
 <213> Homo sapiens

<400> 36

Met Gly Arg Val Glu Asp Gly Gly Thr Thr Glu Glu Leu Glu Asp Trp
 1 5 10 15

Asp Pro Gly Thr Ser Ala Leu Pro Ala Pro Gly Ile Lys Gln Gly Pro
 20 25 30

Arg Glu Gln Thr Gly Thr Gly Pro Leu Ser Gln Lys Cys Trp Glu Pro
 35 40 45

Glu Pro Asp Ala Pro Ser Gln Pro Gly Pro Ala Leu Trp Ser Arg Gly
 50 55 60

Arg Ala Arg Thr Gln Ala Leu Ala Gly Gly Ser Ser Leu Gln Gln Leu
 65 70 75 80

Asp Pro Glu Asn Thr Gly Phe Ile Gly Ala Asp Thr Phe Thr Gly Leu
 85 90 95

Val His Ser His Glu Leu Pro Leu Asp Pro Ala Lys Leu Asp Met Leu
 100 105 110

Val Ala Leu Ala Gln Ser Asn Glu Gln Gly Gln Val Cys Tyr Gln Glu
 115 120 125

Leu Val Asp Leu Ile Ser Ser Lys Arg Ser Ser Ser Phe Lys Arg Ala
 130 135 140

Ile Ala Asn Gly Gln Arg Ala Leu Pro Arg Asp Gly Pro Leu Asp Glu
 145 150 155 160

Pro Gly Leu Gly Val Tyr Lys Arg Phe Val Arg Tyr Val Ala Tyr Glu
 165 170 175

Ile Leu Pro Cys Glu Val Asp Arg Arg Trp Tyr Phe Tyr Arg His Arg
 180 185 190

Ser Cys Pro Pro Pro Val Phe Met Ala Ser Val Thr Leu Ala Gln Ile
 195 200 205

Ile Val Phe Leu Cys Tyr Gly Ala Arg Leu Asn Lys Trp Val Leu Gln
 210 215 220

Thr Tyr His Pro Glu Tyr Met Lys Ser Pro Leu Val Tyr His Pro Gly
 225 230 235 240

His Arg Ala Arg Ala Trp Arg Phe Leu Thr Tyr Met Phe Met His Val
 245 250 255

Gly Leu Glu Gln Leu Gly Phe Asn Ala Leu Leu Gln Leu Met Ile Gly
 260 265 270

Val Pro Leu Glu Met Val His Gly Leu Leu Arg Ile Ser Leu Leu Tyr
 275 280 285

Leu Ala Gly Val Leu Ala Gly Ser Leu Thr Val Ser Ile Thr Asp Met
 290 295 300

Arg Ala Pro Val Val Gly Gly Ser Gly Gly Val Tyr Ala Leu Cys Ser
 305 310 315 320

Ala His Leu Ala Asn Val Val Met Asn Trp Ala Gly Met Arg Cys Pro

325 330 335

Tyr Lys Leu Leu Arg Met Val Leu Ala Leu Val Cys Met Ser Ser Glu
340 345 350

Val Gly Arg Ala Val Trp Leu Arg Phe Ser Pro Pro Leu Pro Ala Ser
355 360 365

Gly Pro Gln Pro Ser Phe Met Ala His Leu Ala Gly Ala Val Val Gly
370 375 380

Val Ser Met Gly Leu Thr Ile Leu Arg Ser Tyr Glu Glu Arg Leu Arg
385 390 395 400

Asp Gln Cys Gly Trp Trp Val Val Leu Leu Ala Tyr Gly Thr Phe Leu
405 410 415

Leu Phe Ala Val Phe Trp Asn Val Phe Ala Tyr Asp Leu Leu Gly Ala
420 425 430

His Ile Pro Pro Pro Pro
435

<210> 37
<211> 292
<212> PRT
<213> Homo sapiens

<400> 37

Met Asn Leu Asn Met Gly Arg Glu Met Lys Glu Glu Leu Glu Glu Glu
1 5 10 15

Glu Lys Met Arg Glu Asp Gly Gly Gly Lys Asp Arg Ala Lys Ser Lys
20 25 30

Lys Val His Arg Ile Val Ser Lys Trp Met Leu Pro Glu Lys Ser Arg
35 40 45

Gly Thr Tyr Leu Glu Arg Ala Asn Cys Phe Pro Pro Pro Val Phe Ile
50 55 60

Ile Ser Ile Ser Leu Ala Glu Leu Ala Val Phe Ile Tyr Tyr Ala Val
65 70 75 80

Trp Lys Pro Gln Lys Gln Trp Ile Thr Leu Asp Thr Gly Ile Leu Glu
85 90 95

Ser Pro Phe Ile Tyr Ser Pro Glu Lys Arg Glu Glu Ala Trp Arg Phe
 100 105 110

Ile Ser Tyr Met Leu Val His Ala Gly Val Gln His Ile Leu Gly Asn
 115 120 125

Leu Cys Met Gln Leu Val Leu Gly Ile Pro Leu Glu Met Val His Lys
 130 135 140

Gly Leu Arg Val Gly Leu Val Tyr Leu Ala Gly Val Ile Ala Gly Ser
 145 150 155 160

Leu Ala Ser Ser Ile Phe Asp Pro Leu Arg Tyr Leu Val Gly Ala Ser
 165 170 175

Gly Gly Val Tyr Ala Leu Met Gly Gly Tyr Phe Met Asn Val Leu Val
 180 185 190

Asn Phe Gln Glu Met Ile Pro Ala Phe Gly Ile Phe Arg Leu Leu Ile
 195 200 205

Ile Ile Leu Ile Ile Val Leu Asp Met Gly Phe Ala Leu Tyr Arg Arg
 210 215 220

Phe Phe Val Pro Glu Asp Gly Ser Pro Val Ser Phe Ala Ala His Ile
 225 230 235 240

Ala Gly Gly Phe Ala Gly Met Ser Ile Gly Tyr Thr Val Phe Ser Cys
 245 250 255

Phe Asp Lys Ala Leu Leu Lys Asp Pro Arg Phe Trp Ile Ala Ile Ala
 260 265 270

Ala Tyr Leu Ala Cys Val Leu Phe Ala Val Phe Phe Asn Ile Phe Leu
 275 280 285

Ser Pro Ala Asn
 290

<210> 38
 <211> 174
 <212> PRT
 <213> Homo sapiens

<400> 38

Leu Cys Arg Val Gln His Ile Leu Gly Asn Leu Cys Met Gln Leu Val
 1 5 10 15

Leu Gly Ile Pro Leu Glu Met Val His Lys Gly Leu Arg Val Gly Leu
 20 25 30

Val Tyr Leu Ala Gly Val Ile Ala Gly Ser Leu Ala Ser Ser Ile Phe
 35 40 45

Asp Pro Leu Arg Tyr Leu Val Gly Ala Ser Gly Gly Val Tyr Ala Leu
 50 55 60

Met Gly Gly Tyr Phe Met Asn Val Leu Val Asn Phe Gln Glu Met Ile
 65 70 75 80

Pro Ala Phe Gly Ile Phe Arg Leu Leu Ile Ile Ile Leu Ile Ile Val
 85 90 95

Leu Asp Met Gly Phe Ala Leu Tyr Arg Arg Phe Phe Val Pro Glu Asp
 100 105 110

Gly Ser Pro Val Ser Phe Ala Ala His Ile Ala Gly Gly Phe Ala Gly
 115 120 125

Met Ser Ile Gly Tyr Thr Val Phe Ser Cys Phe Asp Lys Ala Leu Leu
 130 135 140

Lys Asp Pro Arg Phe Trp Ile Ala Ile Ala Ala Tyr Leu Ala Cys Val
 145 150 155 160

Leu Phe Ala Val Phe Phe Asn Ile Phe Leu Ser Pro Ala Asn
 165 170

<210> 39
 <211> 162
 <212> PRT
 <213> Homo sapiens

<400> 39

Met Gln Leu Val Leu Gly Ile Pro Leu Glu Met Val His Lys Gly Leu
 1 5 10 15

Arg Val Gly Leu Val Tyr Leu Ala Gly Val Ile Ala Gly Ser Leu Ala
 20 25 30

Ser Ser Ile Phe Asp Pro Leu Arg Tyr Leu Val Gly Ala Ser Gly Gly
 35 40 45

Val Tyr Ala Leu Met Gly Gly Tyr Phe Met Asn Val Leu Val Asn Phe
 50 55 60

Gln Glu Met Ile Pro Ala Phe Gly Ile Phe Arg Leu Leu Ile Ile Ile
65 70 75 80

Leu Ile Ile Val Leu Asp Met Gly Phe Ala Leu Tyr Arg Arg Phe Phe
85 90 95

Val Pro Glu Asp Gly Ser Pro Val Ser Phe Ala Ala His Ile Ala Gly
100 105 110

Gly Phe Ala Gly Met Ser Ile Gly Tyr Thr Val Phe Ser Cys Phe Asp
115 120 125

Lys Ala Leu Leu Lys Asp Pro Arg Phe Trp Ile Ala Ile Ala Ala Tyr
130 135 140

Leu Ala Cys Val Leu Phe Ala Val Phe Phe Asn Ile Phe Leu Ser Pro
145 150 155 160

Ala Asn

<210> 40
<211> 379
<212> PRT
<213> Homo sapiens

<400> 40

Met Ala Trp Arg Gly Trp Ala Gln Arg Gly Trp Gly Cys Gly Gln Ala
1 5 10 15

Trp Gly Ala Ser Val Gly Gly Arg Ser Cys Glu Glu Leu Thr Ala Val
20 25 30

Leu Thr Pro Pro Gln Leu Leu Gly Arg Arg Phe Asn Phe Phe Ile Gln
35 40 45

Gln Lys Cys Gly Phe Arg Lys Ala Pro Arg Lys Val Glu Pro Arg Arg
50 55 60

Ser Asp Pro Gly Thr Ser Gly Glu Ala Tyr Lys Arg Ser Ala Leu Ile
65 70 75 80

Pro Pro Val Glu Glu Thr Val Phe Tyr Pro Ser Pro Tyr Pro Ile Arg
85 90 95

Ser Leu Ile Lys Pro Leu Phe Phe Thr Val Gly Phe Thr Gly Cys Ala

100	105	110
Phe Gly Ser Ala Ala Ile Trp Gln Tyr Glu Ser Leu Lys Ser Arg Val		
115	120	125
Gln Ser Tyr Phe Asp Gly Ile Lys Ala Asp Trp Leu Asp Ser Ile Arg		
130	135	140
Pro Gln Lys Glu Gly Asp Phe Arg Lys Glu Ile Asn Lys Trp Trp Asn		
145	150	155
Asn Leu Ser Asp Gly Gln Arg Thr Val Thr Gly Ile Ile Ala Ala Asn		
165	170	175
Val Leu Val Phe Cys Leu Trp Arg Val Pro Ser Leu Gln Arg Thr Met		
180	185	190
Ile Arg Tyr Phe Thr Ser Asn Pro Ala Ser Lys Val Leu Cys Ser Pro		
195	200	205
Met Leu Leu Ser Thr Phe Ser His Phe Ser Leu Phe His Met Ala Ala		
210	215	220
Asn Met Tyr Val Leu Trp Ser Phe Ser Ser Ser Ile Val Asn Ile Leu		
225	230	235
Gly Gln Glu Gln Phe Met Ala Val Tyr Leu Ser Ala Gly Val Ile Ser		
245	250	255
Asn Phe Val Ser Tyr Leu Gly Lys Val Ala Thr Gly Arg Tyr Gly Pro		
260	265	270
Ser Leu Gly Ala Ser Gly Ala Ile Met Thr Val Leu Ala Ala Val Cys		
275	280	285
Thr Lys Ile Pro Glu Gly Arg Leu Ala Ile Ile Phe Leu Pro Met Phe		
290	295	300
Thr Phe Thr Ala Gly Asn Ala Leu Lys Ala Ile Ile Ala Met Asp Thr		
305	310	315
Ala Gly Met Ile Leu Gly Trp Lys Phe Phe Asp His Ala Ala His Leu		
325	330	335
Gly Gly Ala Leu Phe Gly Ile Trp Tyr Val Thr Tyr Gly His Glu Leu		
340	345	350

Ile Trp Lys Asn Arg Glu Pro Leu Val Lys Ile Trp His Glu Ile Arg
 355 360 365

Thr Asn Gly Pro Lys Lys Gly Gly Gly Ser Lys
 370 375

<210> 41
 <211> 855
 <212> PRT
 <213> Homo sapiens

<400> 41

Met Ser Glu Ala Arg Arg Asp Ser Thr Ser Ser Leu Gln Arg Lys Lys
 1 5 10 15

Pro Pro Trp Leu Lys Leu Asp Ile Pro Ser Ala Val Pro Leu Thr Ala
 20 25 30

Glu Glu Pro Ser Phe Leu Gln Pro Leu Arg Arg Gln Ala Phe Leu Arg
 35 40 45

Ser Val Ser Met Pro Ala Glu Thr Ala His Ile Ser Ser Pro His His
 50 55 60

Glu Leu Arg Arg Pro Val Leu Gln Arg Gln Thr Ser Ile Thr Gln Thr
 65 70 75 80

Ile Arg Arg Gly Thr Ala Asp Trp Phe Gly Val Ser Lys Asp Ser Asp
 85 90 95

Ser Thr Gln Lys Trp Gln Arg Lys Ser Ile Arg His Cys Ser Gln Arg
 100 105 110

Tyr Gly Lys Leu Lys Pro Gln Val Leu Arg Glu Leu Asp Leu Pro Ser
 115 120 125

Gln Asp Asn Val Ser Leu Thr Ser Thr Glu Thr Pro Pro Pro Leu Tyr
 130 135 140

Val Gly Pro Cys Gln Leu Gly Met Gln Lys Ile Ile Asp Pro Leu Ala
 145 150 155 160

Arg Gly Arg Ala Phe Arg Val Ala Asp Asp Thr Ala Glu Gly Leu Ser
 165 170 175

Ala Pro His Thr Pro Val Thr Pro Gly Ala Ala Ser Leu Cys Ser Phe
 180 185 190

Ser Ser Ser Arg Ser Gly Phe His Arg Leu Pro Arg Arg Arg Lys Arg
 195 200 205
 Glu Ser Val Ala Lys Met Ser Phe Arg Ala Ala Ala Ala Leu Met Lys
 210 215 220
 Gly Arg Ser Val Arg Asp Gly Thr Phe Arg Arg Ala Gln Arg Arg Ser
 225 230 235 240
 Phe Thr Pro Ala Ser Phe Leu Glu Glu Asp Thr Thr Asp Phe Pro Asp
 245 250 255
 Glu Leu Asp Thr Ser Phe Phe Ala Arg Glu Gly Ile Leu His Glu Glu
 260 265 270
 Leu Ser Thr Tyr Pro Asp Glu Val Phe Glu Ser Pro Ser Glu Ala Ala
 275 280 285
 Leu Lys Asp Trp Glu Lys Ala Pro Glu Gln Ala Asp Leu Thr Gly Gly
 290 295 300
 Ala Leu Asp Arg Ser Glu Leu Glu Arg Ser His Leu Met Leu Pro Leu
 305 310 315 320
 Glu Arg Gly Trp Arg Lys Gln Lys Glu Gly Ala Ala Ala Pro Gln Pro
 325 330 335
 Lys Val Arg Leu Arg Gln Glu Val Val Ser Thr Ala Gly Pro Arg Arg
 340 345 350
 Gly Gln Arg Ile Ala Val Pro Val Arg Lys Leu Phe Ala Arg Glu Lys
 355 360 365
 Arg Pro Tyr Gly Leu Gly Met Val Gly Arg Leu Thr Asn Arg Thr Tyr
 370 375 380
 Arg Lys Arg Ile Asp Ser Phe Val Lys Arg Gln Ile Glu Asp Met Asp
 385 390 395 400
 Asp His Arg Pro Phe Phe Thr Tyr Trp Leu Thr Phe Val His Ser Leu
 405 410 415
 Val Ala Ile Leu Ala Val Cys Ile Tyr Gly Ile Ala Pro Val Gly Phe
 420 425 430

Ser Gln His Glu Thr Val Asp Ser Val Leu Arg Asn Arg Gly Val Tyr
 435 440 445
 Glu Asn Val Lys Tyr Val Gln Gln Glu Asn Phe Trp Ile Gly Pro Ser
 450 455 460
 Ser Glu Ala Leu Ile His Leu Gly Ala Lys Phe Ser Pro Cys Met Arg
 465 470 475 480
 Gln Asp Pro Gln Val His Ser Phe Ile Arg Ser Ala Arg Glu Arg Glu
 485 490 495
 Lys His Ser Ala Cys Cys Val Arg Asn Asp Arg Ser Gly Cys Val Gln
 500 505 510
 Thr Ser Glu Glu Glu Cys Ser Ser Thr Leu Ala Val Trp Val Lys Trp
 515 520 525
 Pro Ile His Pro Ser Ala Pro Glu Leu Ala Gly His Lys Arg Gln Phe
 530 535 540
 Gly Ser Val Cys His Gln Asp Pro Arg Val Cys Asp Glu Pro Ser Ser
 545 550 555 560
 Glu Asp Pro His Glu Trp Pro Glu Asp Ile Thr Lys Trp Pro Ile Cys
 565 570 575
 Thr Lys Asn Ser Ala Gly Asn His Thr Asn His Pro His Met Asp Cys
 580 585 590
 Val Ile Thr Gly Arg Pro Cys Cys Ile Gly Thr Lys Gly Arg Cys Glu
 595 600 605
 Ile Thr Ser Arg Glu Tyr Cys Asp Phe Met Arg Gly Tyr Phe His Glu
 610 615 620
 Glu Ala Thr Leu Cys Ser Gln Val His Cys Met Asp Asp Val Cys Gly
 625 630 635 640
 Leu Leu Pro Phe Leu Asn Pro Glu Val Pro Asp Gln Phe Tyr Arg Leu
 645 650 655
 Trp Leu Ser Leu Phe Leu His Ala Gly Ile Leu His Cys Leu Val Ser
 660 665 670
 Ile Cys Phe Gln Met Thr Val Leu Arg Asp Leu Glu Lys Leu Ala Gly
 675 680 685

Trp His Arg Ile Ala Ile Ile Tyr Leu Leu Ser Gly Val Thr Gly Asn
690 695 700

Leu Ala Ser Ala Ile Phe Leu Pro Tyr Arg Ala Glu Val Gly Pro Ala
705 710 715 720

Gly Ser Gln Phe Gly Ile Leu Ala Cys Leu Phe Val Glu Leu Phe Gln
725 730 735

Ser Trp Gln Ile Leu Ala Arg Pro Trp Arg Ala Phe Phe Lys Leu Leu
740 745 750

Ala Val Val Leu Phe Leu Phe Thr Phe Gly Leu Leu Pro Trp Ile Asp
755 760 765

Asn Phe Ala His Ile Ser Gly Phe Ile Ser Gly Leu Phe Leu Ser Phe
770 775 780

Ala Phe Leu Pro Tyr Ile Ser Phe Gly Lys Phe Asp Leu Tyr Arg Lys
785 790 795 800

Arg Cys Gln Ile Ile Ile Phe Gln Val Val Phe Leu Gly Leu Leu Ala
805 810 815

Gly Leu Val Val Leu Phe Tyr Val Tyr Pro Val Arg Cys Glu Trp Cys
820 825 830

Glu Phe Leu Thr Cys Ile Pro Phe Thr Asp Lys Phe Cys Glu Lys Tyr
835 840 845

Glu Leu Asp Ala Gln Leu His
850 855

<210> 42
<211> 619
<212> PRT
<213> Homo sapiens

<400> 42

Met Ser Val Ala His Met Ser Leu Gln Ala Ala Ala Ala Leu Leu Lys
1 5 10 15

Gly Arg Ser Val Leu Asp Ala Thr Gly Gln Arg Cys Arg Val Val Lys
20 25 30

Arg Ser Phe Ala Phe Pro Ser Phe Leu Glu Glu Asp Val Val Asp Gly

35	40	45													
Ala	Asp	Thr	Phe	Asp	Ser	Ser	Phe	Phe	Ser	Lys	Glu	Glu	Met	Ser	Ser
50					55						60				
Met	Pro	Asp	Asp	Val	Phe	Glu	Ser	Pro	Pro	Leu	Ser	Ala	Ser	Tyr	Phe
65					70					75					80
Arg	Gly	Ile	Pro	His	Ser	Ala	Ser	Pro	Val	Ser	Pro	Asp	Gly	Val	Gln
				85					90					95	
Ile	Pro	Leu	Lys	Glu	Tyr	Gly	Arg	Ala	Pro	Val	Pro	Gly	Pro	Arg	Arg
			100					105					110		
Gly	Lys	Arg	Ile	Ala	Ser	Lys	Val	Lys	His	Phe	Ala	Phe	Asp	Arg	Lys
		115					120					125			
Lys	Arg	His	Tyr	Gly	Leu	Gly	Val	Val	Gly	Asn	Trp	Leu	Asn	Arg	Ser
	130					135				140					
Tyr	Arg	Arg	Ser	Ile	Ser	Ser	Thr	Val	Gln	Arg	Gln	Leu	Glu	Ser	Phe
145					150				155						160
Asp	Ser	His	Arg	Pro	Tyr	Phe	Thr	Tyr	Trp	Leu	Thr	Phe	Val	His	Val
				165					170					175	
Ile	Ile	Thr	Leu	Leu	Val	Ile	Cys	Thr	Tyr	Gly	Ile	Ala	Pro	Val	Gly
			180					185					190		
Phe	Ala	Gln	His	Val	Thr	Thr	Gln	Leu	Val	Leu	Arg	Asn	Lys	Gly	Val
		195					200					205			
Tyr	Glu	Ser	Val	Lys	Tyr	Ile	Gln	Gln	Glu	Asn	Phe	Trp	Val	Gly	Pro
	210					215					220				
Ser	Ser	Ile	Asp	Leu	Ile	His	Leu	Gly	Ala	Lys	Phe	Ser	Pro	Cys	Ile
225					230					235					240
Arg	Lys	Asp	Gly	Gln	Ile	Glu	Gln	Leu	Val	Leu	Arg	Glu	Arg	Asp	Leu
				245					250					255	
Glu	Arg	Asp	Ser	Gly	Cys	Cys	Val	Gln	Asn	Asp	His	Ser	Gly	Cys	Ile
			260					265					270		
Gln	Thr	Gln	Arg	Lys	Asp	Cys	Ser	Glu	Thr	Leu	Ala	Thr	Phe	Val	Lys
		275					280					285			

Trp Gln Asp Asp Thr Gly Pro Pro Met Asp Lys Ser Asp Leu Gly Gln
 290 295 300

Lys Arg Thr Ser Gly Ala Val Cys His Gln Asp Pro Arg Thr Cys Glu
 305 310 315 320

Glu Pro Ala Ser Ser Gly Ala His Ile Trp Pro Asp Asp Ile Thr Lys
 325 330 335

Trp Pro Ile Cys Thr Glu Gln Ala Arg Ser Asn His Thr Gly Phe Leu
 340 345 350

His Met Asp Cys Glu Ile Lys Gly Arg Pro Cys Cys Ile Gly Thr Lys
 355 360 365

Gly Ser Cys Glu Ile Thr Thr Arg Glu Tyr Cys Glu Phe Met His Gly
 370 375 380

Tyr Phe His Glu Glu Ala Thr Leu Cys Ser Gln Val His Cys Leu Asp
 385 390 395 400

Lys Val Cys Gly Leu Leu Pro Phe Leu Asn Pro Glu Val Pro Asp Gln
 405 410 415

Phe Tyr Arg Leu Trp Leu Ser Leu Phe Leu His Ala Gly Val Val His
 420 425 430

Cys Leu Val Ser Val Val Phe Gln Met Thr Ile Leu Arg Asp Leu Glu
 435 440 445

Lys Leu Ala Gly Trp His Arg Ile Ala Ile Ile Phe Ile Leu Ser Gly
 450 455 460

Ile Thr Gly Asn Leu Ala Ser Ala Ile Phe Leu Pro Tyr Arg Ala Glu
 465 470 475 480

Val Gly Pro Ala Gly Ser Gln Phe Gly Leu Leu Ala Cys Leu Phe Val
 485 490 495

Glu Leu Phe Gln Ser Trp Pro Leu Leu Glu Arg Pro Trp Lys Ala Phe
 500 505 510

Leu Asn Leu Ser Ala Ile Val Leu Phe Leu Phe Ile Cys Gly Leu Leu
 515 520 525

Pro Trp Ile Asp Asn Ile Ala His Ile Phe Gly Phe Leu Ser Gly Leu

530 535 540
 Leu Leu Ala Phe Ala Phe Leu Pro Tyr Ile Thr Phe Gly Thr Ser Asp
 545 550 555 560
 Lys Tyr Arg Lys Arg Ala Leu Ile Leu Val Ser Leu Leu Ala Phe Ala
 565 570 575
 Gly Leu Phe Ala Ala Leu Val Leu Trp Leu Tyr Ile Tyr Pro Ile Asn
 580 585 590
 Trp Pro Trp Ile Glu His Leu Thr Cys Phe Pro Phe Thr Ser Arg Phe
 595 600 605
 Cys Glu Lys Tyr Glu Leu Asp Gln Val Leu His
 610 615
 <210> 43
 <211> 434
 <212> PRT
 <213> Homo sapiens
 <400> 43
 Met Gly Arg Gly Leu Trp Glu Ala Trp Pro Pro Ala Gly Ser Ser Ala
 1 5 10 15
 Val Ala Lys Gly Asn Cys Arg Glu Glu Ala Glu Gly Ala Glu Asp Arg
 20 25 30
 Gln Pro Ala Ser Arg Arg Gly Ala Gly Thr Thr Ala Ala Met Ala Ala
 35 40 45
 Ser Gly Pro Gly Cys Arg Ser Trp Cys Leu Cys Pro Glu Val Pro Ser
 50 55 60
 Ala Thr Phe Phe Thr Ala Leu Leu Ser Leu Leu Val Ser Gly Pro Arg
 65 70 75 80
 Leu Phe Leu Leu Gln Gln Pro Leu Ala Pro Ser Gly Leu Thr Leu Lys
 85 90 95
 Ser Glu Ala Leu Arg Asn Trp Gln Val Tyr Arg Leu Val Thr Tyr Ile
 100 105 110
 Phe Val Tyr Glu Asn Pro Ile Ser Leu Leu Cys Gly Ala Ile Ile Ile
 115 120 125

Trp Arg Phe Ala Gly Asn Phe Glu Arg Thr Val Gly Thr Val Arg His
 130 135 140

Cys Phe Phe Thr Val Ile Phe Ala Ile Phe Ser Ala Ile Ile Phe Leu
 145 150 155 160

Ser Phe Glu Ala Val Ser Ser Leu Ser Lys Leu Gly Glu Val Glu Asp
 165 170 175

Ala Arg Gly Phe Thr Pro Val Ala Phe Ala Met Leu Gly Val Thr Thr
 180 185 190

Val Arg Ser Arg Met Arg Arg Ala Leu Val Phe Gly Met Val Val Pro
 195 200 205

Ser Val Leu Val Pro Trp Leu Leu Leu Gly Ala Ser Trp Leu Ile Pro
 210 215 220

Gln Thr Ser Phe Leu Ser Asn Val Cys Gly Leu Ser Ile Gly Leu Ala
 225 230 235 240

Tyr Ala His Leu Leu Leu Phe His Arg Pro Leu Arg Ala Ser Gly Ala
 245 250 255

Glu Ala Arg Ser Asp Leu Pro Leu Gln Pro Asp Glu Glu Asp Ile Arg
 260 265 270

Val Gln Val Arg Leu Arg Val Phe Ser Arg Glu Glu Gly Ser Pro Glu
 275 280 285

Pro Glu Thr Glu Pro Gly Ala Trp Leu Leu Pro His Thr Glu Leu Pro
 290 295 300

Pro Ser Pro Val Pro Lys Pro Pro Cys Val Pro Asp Ala Ala Arg Gln
 305 310 315 320

Trp Ser Glu Ala Gly Leu Leu Ala Ser Cys Thr Pro Gly His Met Pro
 325 330 335

Thr Leu Pro Pro Tyr Gln Pro Ala Ser Gly Leu Cys Tyr Val Gln Asn
 340 345 350

His Phe Gly Pro Asn Pro Thr Ser Ser Ser Val Tyr Pro Ala Ser Ala
 355 360 365

Gly Thr Ser Leu Gly Ile Gln Pro Pro Thr Pro Val Asn Ser Pro Gly
 370 375 380

Thr Val Tyr Ser Gly Ala Leu Gly His Gln Gly Leu Gln Ala Pro Arg
 385 390 395 400

Ser Pro Pro Gly Ser Pro Cys Pro Glu Arg Ile Ser Arg Glu Val Ile
 405 410 415

Ser Leu Gly Leu Leu Lys Val Leu Pro Lys Ser Leu Leu Thr Lys Val
 420 425 430

Thr Tyr

<210> 44
 <211> 386
 <212> PRT
 <213> Homo sapiens

<400> 44

Met His Ala Arg Gly Pro His Gly Gln Leu Ser Pro Ala Leu Pro Leu
 1 5 10 15

Ala Ser Ser Val Leu Met Leu Leu Met Ser Thr Leu Trp Leu Val Gly
 20 25 30

Ala Gly Pro Gly Leu Val Leu Ala Pro Glu Leu Leu Leu Asp Pro Trp
 35 40 45

Gln Val His Arg Leu Leu Thr His Ala Leu Gly His Thr Ala Leu Pro
 50 55 60

Gly Leu Leu Leu Ser Leu Leu Leu Leu Pro Thr Val Gly Trp Gln Gln
 65 70 75 80

Glu Cys His Leu Gly Thr Leu Arg Phe Leu His Ala Ser Ala Leu Leu
 85 90 95

Ala Leu Ala Ser Gly Leu Leu Ala Val Leu Leu Ala Gly Leu Gly Leu
 100 105 110

Ser Ser Ala Ala Gly Ser Cys Gly Tyr Met Pro Val His Leu Ala Met
 115 120 125

Leu Ala Gly Glu Gly His Arg Pro Arg Arg Pro Arg Gly Ala Leu Pro
 130 135 140

Pro Trp Leu Ser Pro Trp Leu Leu Leu Ala Leu Thr Pro Leu Leu Ser

145	150	155	160
Ser Glu Pro Pro Phe Leu Gln Leu Leu Cys Gly Leu Leu Ala Gly Leu	165	170	175
Ala Tyr Ala Ala Gly Ala Phe Arg Trp Leu Glu Pro Ser Glu Arg Arg	180	185	190
Leu Gln Val Leu Gln Glu Gly Val Leu Cys Arg Thr Leu Ala Gly Cys	195	200	205
Trp Pro Leu Arg Leu Leu Ala Thr Pro Gly Ser Leu Ala Glu Leu Pro	210	215	220
Val Thr His Pro Ala Gly Val Arg Pro Pro Ile Pro Gly Pro Pro Tyr	225	230	235
Val Ala Ser Pro Asp Leu Trp Ser His Trp Glu Asp Ser Ala Leu Pro	245	250	255
Pro Pro Ser Leu Arg Pro Val Gln Pro Thr Trp Glu Gly Ser Ser Glu	260	265	270
Ala Gly Leu Asp Trp Ala Gly Ala Ser Phe Ser Pro Gly Thr Pro Met	275	280	285
Trp Ala Ala Leu Asp Glu Gln Met Leu Gln Glu Gly Ile Gln Ala Ser	290	295	300
Leu Leu Asp Gly Pro Ala Gln Glu Pro Gln Ser Ala Pro Trp Leu Ser	305	310	315
Lys Ser Ser Val Ser Ser Leu Arg Leu Gln Gln Leu Glu Arg Met Gly	325	330	335
Phe Pro Thr Glu Gln Ala Val Val Ala Leu Ala Ala Thr Gly Arg Val	340	345	350
Glu Gly Ala Val Ser Leu Leu Val Gly Gly Gln Val Gly Thr Glu Thr	355	360	365
Leu Val Thr His Gly Lys Gly Gly Pro Ala His Ser Glu Gly Pro Gly	370	375	380
Pro Pro	385		

<210> 45
 <211> 1413
 <212> DNA
 <213> Mus musculus

<400> 45
 atggcctact gggagaagct tctgccgcac gtggtgcaag agctggacgc aggcagtggc 60
 tctgggtcga gctgttcggc gggggacgat ctaccacaggc agtgcgtcga gttggggccg 120
 cgcgctgcgcg tgcaggatat gcgcgtactc tggctgtgtg cgcgctgggc tggcatctcg 180
 gtgacgctgt ctgagctcgg ggagcgcaga gcagagcagg cggagcgggc cggccggggc 240
 ggagcggagc ggtccgcaga gcagccctc cgggcctcgg ccgaccccg ccctcggccc 300
 ggctctatgg acaggagctc gctgctgcag ctcatcagg agcagcagct ggatcctgag 360
 aacacaggct tcatcgggtc ggacacctc gctggctctg tacacagcca tgagctgccc 420
 ctggacccca ccaagttgga catgttggtg gctctggctc agagcaacga gcggggccag 480
 gtctgctacc aggagctggg ggacctggtc agtgccatga tcagcagcaa gcgttcacg 540
 agcttcaaga gagcattgc taacggacag cgggcactgc cccgagacgg actgctagat 600
 gagccaggcc tgagtgtcta caagcggttt gtgcgctatg tggcctacga gatcctgccc 660
 tgcgagggtg accgccgctg gtacttctac aggcaccgca cctgcccacc ccctgtgttc 720
 atggcctcgg tcaactcttc ccagtgtccc cagatcatcg tgttcctgtg ctacggggca 780
 cgtctcaaca agtgggtgct ccagacctac caccctgaat atatgaagag ccctctggtg 840
 tatcaccag gacaccgtgc tcgggcatgg cgcttctca cctacatgtt catgcatgtc 900
 gggctggagc agctaggggt caatgccctc ctgcagctga tgatcgggtg gccctggag 960
 atggtacatg gcgtgcttcg catcagcctg ctctacctgg cgggcgtgct ggcaggctcc 1020
 ctgactgtct ctatcacaga tatgcgtgcc ccggtggtag ggggctctgg aggggtctat 1080
 gccctgtgct cagcacacct ggccaatgtt gtcatggtaa tgggactgcc tttctgggga 1140
 gtgagttccg aagtgggccc ggctgtgtgg ctacgcttct cccaccact gcctgcctca 1200
 ggcccacagc ccagcttcat ggcacacctg gctggtgcag tggtaggtgt aagcatgggc 1260
 cttaccatcc ttggagcta tgaggaacgc ctgagggacc agtgcggtg gtgggtggtg 1320
 ctacttgct atggcacctt cctgcttttc gccatcttct ggaacgtctt tgcctatgac 1380
 ctgctgggtg ccgatatccc ccctccacct tga 1413

<210> 46
 <211> 470
 <212> PRT
 <213> Mus musculus

<400> 46

Met Ala Tyr Trp Glu Lys Leu Leu Pro His Val Val Gln Glu Leu Asp
 1 5 10 15
 Ala Gly Ser Gly Ser Gly Ser Ser Cys Ser Val Gly Asp Asp Leu Pro
 20 25 30
 Arg Gln Cys Val Glu Leu Gly Pro Arg Val Arg Val Gln Asp Met Arg
 35 40 45
 Val Leu Trp Leu Cys Ala Arg Trp Ala Gly Ile Ser Val Thr Leu Ser
 50 55 60
 Glu Leu Gly Glu Arg Arg Ala Glu Gln Ala Glu Arg Ala Gly Arg Gly
 65 70 75 80
 Gly Ala Glu Arg Ser Ala Glu Gln Pro Leu Pro Ala Ser Ala Asp Pro
 85 90 95
 Gly Pro Arg Pro Gly Ser Met Asp Arg Ser Ser Leu Leu Gln Leu Ile
 100 105 110
 Gln Glu Gln Gln Leu Asp Pro Glu Asn Thr Gly Phe Ile Gly Ala Asp
 115 120 125
 Thr Phe Ala Gly Leu Val His Ser His Glu Leu Pro Leu Asp Pro Thr
 130 135 140
 Lys Leu Asp Met Leu Val Ala Leu Ala Gln Ser Asn Glu Arg Gly Gln
 145 150 155 160
 Val Cys Tyr Gln Glu Leu Val Asp Leu Val Ser Ala Met Ile Ser Ser
 165 170 175
 Lys Arg Ser Ser Ser Phe Lys Arg Ala Ile Ala Asn Gly Gln Arg Ala
 180 185 190
 Leu Pro Arg Asp Gly Leu Leu Asp Glu Pro Gly Leu Ser Val Tyr Lys
 195 200 205
 Arg Phe Val Arg Tyr Val Ala Tyr Glu Ile Leu Pro Cys Glu Val Asp
 210 215 220
 Arg Arg Trp Tyr Phe Tyr Arg His Arg Thr Cys Pro Pro Pro Val Phe
 225 230 235 240
 Met Ala Ser Val Thr Leu Ala Gln Cys Pro Gln Ile Ile Val Phe Leu

	245		250		255
Cys Tyr Gly Ala Arg Leu Asn Lys Trp Val Leu Gln Thr Tyr His Pro	260		265		270
Glu Tyr Met Lys Ser Pro Leu Val Tyr His Pro Gly His Arg Ala Arg	275		280		285
Ala Trp Arg Phe Leu Thr Tyr Met Phe Met His Val Gly Leu Glu Gln	290		295		300
Leu Gly Phe Asn Ala Leu Leu Gln Leu Met Ile Gly Val Pro Leu Glu	305		310		315
Met Val His Gly Val Leu Arg Ile Ser Leu Leu Tyr Leu Ala Gly Val		325		330	335
Leu Ala Gly Ser Leu Thr Val Ser Ile Thr Asp Met Arg Ala Pro Val		340		345	350
Val Gly Gly Ser Gly Gly Val Tyr Ala Leu Cys Ser Ala His Leu Ala		355		360	365
Asn Val Val Met Val Met Gly Leu Pro Phe Trp Gly Val Ser Ser Glu		370		375	380
Val Gly Arg Ala Val Trp Leu Arg Phe Ser Pro Pro Leu Pro Ala Ser		385		390	395
Gly Pro Gln Pro Ser Phe Met Ala His Leu Ala Gly Ala Val Val Gly		405		410	415
Val Ser Met Gly Leu Thr Ile Leu Arg Ser Tyr Glu Glu Arg Leu Arg		420		425	430
Asp Gln Cys Gly Trp Trp Val Val Leu Leu Ala Tyr Gly Thr Phe Leu		435		440	445
Leu Phe Ala Ile Phe Trp Asn Val Phe Ala Tyr Asp Leu Leu Gly Ala		450		455	460
Asp Ile Pro Pro Pro Pro	465		470		

<210> 47
 <211> 34319
 <212> DNA

<213> Mus musculus

<220>

<221> misc_feature

<222> (34274)..(34275)

<223> n is a, c, g, or t

<220>

<221> misc_feature

<222> (34285)..(34285)

<223> n is a, c, g, or t

<220>

<221> misc_feature

<222> (34297)..(34298)

<223> n is a, c, g, or t

<220>

<221> misc_feature

<222> (34301)..(34301)

<223> n is a, c, g, or t

<400> 47

actagtcctg caggtttaaa cgaattcgcc ctttcctcc tgagcctgcc ctcacccttg	60
tccagtgttc agccaagcac ttgaggaata tctcagagct gttctactat gcccagaagg	120
ctgttctgca tcccacagcc cccctttatg accctgaggc caaacaggta agtagtggat	180
gagggccaca cacctatttc tcgaaggccc atccatgtgg gccttgata aggagtacaa	240
gtaaatacgc gtaccgacag ggtggccagg aaaaggcca cttggtgttg tgtccctgca	300
gctcaggcct gcattgcgcc aggcgctcac acgcattctc aggctgtcgg accaggaccg	360
agaccacggg ctgagcgacg aggaactcaa tgccttccag gtgtggctcct gccctgacac	420
cctagctgcc ctgtgagag ggtcatcagg atagctgact gatctgaagc actctcttac	480
cggaggcaga agtcctgctt tggccatccc ctggccccac aggccctgga ggatgtgaag	540
agggtggtgt gcaagaacgt gtcagggtgt gtgcagaatg accggctgac cctggaaggt	600
gaggcagaca tcacctata tcctgatgtg tggggagtgg caaggcgggtg tactaacctt	660
ccaccttccc acaggcttcc tcttctgaa cacactcttc atccagcgtg gccggcacga	720
gaccacgtgg accatcctgc ggcgctttgg ttacagtgc tcaactggagc taacgcccga	780
ctacctctat ccagcgtgag ttgtaggctg gtgggcagtg gtgcttatcc tcttggggtc	840
aatccttggt gcctggggac tttagcttgg tctctgagca cacactcatt agctcccccc	900
ttgcttagcc ccaaccgtgt aaccagaggt tctgttggga gtgcctgact gccaggcac	960
ttttaactgt cacagggtgc agtaaccctg ccttgtagca gttaaagtgc atccccacag	1020
gctccatgtc cccctgggt gcagcacaga gctcaacct cgtggctacc agtttgtgca	1080
gcggatgttc gagaagcatg accaggtaag caggtcacct cagccttga caccctccc	1140

cgcaaagaaa cttgtcacca tagccccctt gcctgtagga ccatgatggg gtcctctcgc 1200
 ccacggaact acagaatctc ttcagtggtg tctcaggtgc tccatggggc cctgaactct 1260
 tacacacagt ccccaactcag gctgggtgtc tgcccttgca tggctatctc tgccagtggg 1320
 cgtaagtaca gtccctgtct acctgtctgt ccctacccat cccatcttcc actctcctta 1380
 gccctgtacc ctccctctact cccagcctca tgacctactt ggatgtccag caatgccttg 1440
 cacaccttgg ctacctgggt taccctaccc tctgtgaaca ggactctcag gcacaggcta 1500
 tcacaggtag gtgaccactt ttttggtccc tggggccagc caaaccttc ttgatggcca 1560
 acaccctcac gtgagcgcac cttttcagtt acccgcgaga agaagctgga ccaggagaaa 1620
 gggcagacac agcgagcgt tctcatgtgt aagggtgctgg gagcccgagg agtgggcaag 1680
 tcagccttct tgcaagcctt ccttggaac agccttgggg tgagaattat ggaggcccca 1740
 gccctacct ggggaactcaa gggggcagag ggggtgccaca gagaactctg gggccaaaaa 1800
 ctcttcctgc tgtccacaca ggaagccagg gacccccctg agaaattccc cctgcatacc 1860
 atcaatacag tgcgggtcaa cgggcaggag aagtacctga ttgtgagtga gggcctgtgc 1920
 accatgtcct gaagggtata cctgtgacac ccatataccc atatattctt cctcctaattg 1980
 gagctgtctc aggggtggcc ttgagggtac cagctgtggg gaggccacta aagctactgt 2040
 ttgtcccagc tgtgtgaagt gaatgctgac agcctgctgg acacctctct agacactacc 2100
 tgtgatgtcg cctgcttaat gtttgacagc agtgatccca agacctttgt aactgtgct 2160
 accatataca aggcaagccc tgacttgagg tcctgtgggg ggatcccagc catagcccag 2220
 ttagccagca agctcagctc tcgaatgttt cttttttcct ggcagcgta ttacatggat 2280
 gggcagaccc cctgcctctt tattgcctcc aaagctgac tccctgaagg tgttgcccc 2340
 ccagggtgt caccgctga gttctgccgg aggcaccggc tgctgcccc agcttcatte 2400
 tcctgcttgg gacccgcaat gccagcaca gatgtcttta cccaacttg taccatggcc 2460
 actttcccggt gagtataccg cagcctcaat gtgggtgggg cttagtctgt ctgggtacaa 2520
 aggtgtcaga gctgtagacc cgtggcctgt gggccagaga gctatgctgg atggcctctg 2580
 gcttgccctc cgctgcctg ggacgggttt tcagcagtga gcttttcgca agcacacgtc 2640
 ttgagtatta ttctttccac agacacctgg tgcacacaga gctgcacccc acctccttct 2700
 ggctccgagg agtgctggtg gctgtcggca cggctgtggc tgctgtcctt agcttctcac 2760
 tatacaggg cctagtgaag agccgatgac aaaggaccca tgagcctctc ctgccccag 2820
 gcacagcaca ctgtggtgcc ttatgtgggc ccacacggca aagaagtatg ggctttcggc 2880
 ctgttgaggt aatatctgtg tggctggcta cagggtccct gggactgatg tgggttcctg 2940
 ctcaggaact tggatattag atcttggtt ccaaaagtct gttgccctat tccaggcctg 3000

tcctgctgag	cctgagagta	ctgggtgccag	gtgtgaggcc	ttagccttgg	agggagggag	3060
gattgtgtgt	aaggggctgg	gctgtgggta	aggaatccag	tgaccaggac	ccgaagttct	3120
caggggtgtgt	tccccccoct	tcctggttgt	gcatggccag	tggacatgta	gagggcaaga	3180
ggtagtggtc	aaatcagggg	gggacccgat	agtcacagag	gtcccacctt	tttgtcaggt	3240
gtggagcctt	atccacactg	cgccagctgc	agagcaagcg	gaactgactt	tcctgattaa	3300
atttcacttg	tcttttccat	ctcttggtat	ttgatctctc	aatgcaactt	actttctctc	3360
tttctctccc	tcccttccct	ccttcctttc	ttccttcttc	ctttcttctt	tctttctttc	3420
tttctttctt	tctttctttc	tttctttctt	tcaaaaatat	gcttattttt	atgagtttga	3480
gtgttttgcc	tgcccatata	tgtaagtgcc	ccgtgtgtgt	gcctgggtacc	gacagaagcc	3540
aaatgagtat	gtcagagttc	cctggaactg	gagttaacta	atgattgtga	gccatcccgt	3600
gggtgttgga	aacaaaccca	ggtcctctgc	agaggcagca	agtgtctctga	accaccaagt	3660
cctctctcca	gccccataaa	aaatggcttt	tcaatcgcaa	tgtgactgct	gtttttgtgt	3720
gagttggaaa	tgccacctga	cccttctgag	acgtctcccc	tgtgtcctgg	ttcccattct	3780
gtggtctcac	tagtcatgct	gattaatggc	tacggaaggt	ccagttaagg	ggtctgctgt	3840
ctagtaatgc	cagcatccat	cagtgtcatg	gggggtggggg	tgtgtgtgag	agggctcctg	3900
ctgcttctct	gggaacattg	cttctgctgt	tggttcttca	ttgccccaca	ctccacacag	3960
cccaggcagc	ctgcctggct	ctgctgagaa	cctgggcttt	atgtgtagcc	tggctgttac	4020
aagcaagggc	cagccataga	tctcaacctg	gccagaagct	gagtgaatga	tggactcgga	4080
gttgatcctg	tgacatgacc	tcctaccacc	aaagacaccc	aaacaggaat	ccatacactc	4140
aacgagtcac	ttgggttggg	tcaggtgggt	tctctctgcc	ccagaatgtc	cacagaacag	4200
ggatccctga	aaaatgaggc	ctttgcctcc	tgctccagtc	ccaacctact	ggcctgccgt	4260
cccactgagc	atctaggcag	gctgcttatt	gtaggcagct	gtattccctg	tagaggccca	4320
tcgggcaaga	ataactgcca	tcccatcttt	gcagtagcag	attgttttga	atgatacttg	4380
gggggggtgc	catggttaca	catcctattg	gtctgtcacc	ttcactgtcc	tcaggagcac	4440
acccccctccc	ttatccaagc	tgcagtcagt	tctaccttca	cgtggctatt	tgtggtcttt	4500
cctggctatc	ccctgtgcct	tcttgactgg	tgtttttgca	ctggaggccc	tgctatctca	4560
aatccaattc	agggatgttt	ggttagagct	tccagccttg	gccatgaata	gggaacctgg	4620
ggcttcgttg	ctggaacagt	gcaaataagg	acaagtgtgt	ctactgaaag	cagctccagc	4680
tcaggagaaa	gtatgatggg	agcaactgga	ccttgacatt	ctggggatgg	gtcacagacc	4740
tatttttagt	agaaagaaag	tatcctaggg	aaggtaataa	aacaggctgc	ctaagagggg	4800
attgggtgct	gcaggctgcc	agctatggcc	tactgggaga	agcttctgcc	gcacgtggtg	4860

caagagctgg	acgcaggcag	tggttctggg	tcgagctgtt	cggtggggga	cgatctaccc	4920
aggcagtgcg	tcgagttggg	gccgcgcgtg	cgcgtgcagg	atatgcgcgt	actctggctg	4980
tgtgcgcgct	gggctggcat	ctcgggtgacg	ctgtctgagc	tcggggagcg	cagagcagag	5040
caggcggagc	gggccggccg	gggcggagcg	gagcgggtccg	cagagcagcc	cctcccggcc	5100
tcggccgacc	ccggccctcg	gcccggctct	atggacagga	gctcgtctgt	gcagctcatt	5160
caggagcagg	tgctgtgtgg	gtgggagctg	gcactgcctt	ggcgcccttc	cagccgggtga	5220
ggggttcggg	ggagggacca	tgctcgcaga	agggccaggc	agccgcctcc	caaaccccc	5280
tccctgcacg	accttggctt	ctgagactcc	cgaccccaag	aaaggctctg	cttcatcggg	5340
tttccacgga	aagctgggga	tggaggggtg	tggctccttg	acctaggtcc	ttcccccttc	5400
taggactggt	agagcgaata	tgaggtggta	gaacagccaa	accaggcacc	ttgccctgcc	5460
agggcctggg	aacaagctgg	ttctagcaaa	caggcactga	ctggcacaaa	ggtccagcct	5520
tcctagccag	cctagctcag	ccctgtggct	gggagttggc	ccaccctcta	ctccccacc	5580
acagcagctg	gacctgaga	acacaggctt	catcggtgcg	gacaccttcg	ctggtctggt	5640
acacagccat	gagctgcccc	tggaccccac	caagttggac	atgttggtgg	ctctggtcca	5700
gagcaacgag	cggggccagg	tctgctacca	ggagctggtg	gacctggtca	tgcccatggt	5760
gtgtgggcag	caggggaggg	cccaggcctg	gtcaagcgca	gcaccctgac	caaagctggg	5820
tgggcacaca	gctttttact	gggtagggct	tacagacctg	ggcagtggtg	aagaccctca	5880
gaccattagc	agccacacag	ccaagtaagg	tagggttcca	caggaacaat	actgtcacct	5940
acactaaggt	tcctgccctt	cagatcagca	gcaagcggtc	cagcagcttc	aagagagcca	6000
ttgctaacgg	acagcgggca	ctgccccgag	acggactgct	agatgagcca	ggcctgagtg	6060
tctacaagcg	gtttgtgcgc	tatgtggcct	acgagatcct	gccctgcgag	gtggaccgcc	6120
gctggtactt	ctacaggcac	cgcacctgcc	cacccctgt	gttcatggcc	tcggtcactc	6180
ttgccagggt	gggcctgcct	acacattgcc	ctggggcctc	ctgagtcctt	agtcttgacc	6240
agccctaaca	ctagtgtccc	cagatcatcg	tgttcctgtg	ctacggggca	cgtctcaaca	6300
agtgggtgct	ccagacctac	caccctgaat	atatgaagag	ccctctggtg	tatcaccag	6360
gacaccgtgc	tcgggcatgg	cgttcctca	cctacatgtt	catgcatgtc	gggtaagtga	6420
gccttgccct	ccagagcctc	ttctccactc	ctacccccag	cacctgcca	ctgactctga	6480
aggcttatct	cccagctaag	ttacactgcc	tgaggtggtc	tggagtaggc	agggtaacca	6540
tctgaccacc	tctgccatgc	tcttctgtgg	ccaggctgga	gcagctaggg	ttcaatgcc	6600
tcctgcagct	gatgatcggt	gtgcccctgg	agatggtaca	tggcgtgctt	cgcacagcc	6660
tgctctacct	ggcgggcgtg	ctggcaggtg	agactggcat	ataggtgggt	gggtcccctg	6720

cctacctcac	ctgctgaggg	gagggccagc	tgtgccctca	cctctccctt	tgcccacagg	6780
ctccctgact	gtctctatca	cagatatgcg	tgccccctg	gtagggggct	ctggaggggt	6840
ctatgccctg	tgctcagcac	acctggccaa	tgttgctcatg	gtaatgggac	tgccctttctg	6900
gggaggtggg	aaggggcca	ttggacatgc	ttcacagcct	gtctgcctgc	ccccatcaga	6960
actgggctgg	gatgcgggtg	ccatacaagc	tgctgaggat	ggtgctggct	ctggtgtgca	7020
gtgagtagtg	gggcaggtgg	ggcgctggga	ggccctttac	ctgcagacag	ggcccctccc	7080
acctgttgct	ccctctgcag	tgagttccga	agtggggcgg	gctgtgtggc	tacgcttctc	7140
cccaccactg	cctgcctcag	gcccacagcc	cagcttcatg	gcacacctgg	ctggtgcagt	7200
ggtaggtgta	agcatgggcc	ttaccatcct	tcggagctat	gaggaacgcc	tgagggacca	7260
gtgcggctgg	tgggtggtgc	tacttgcccta	tggcaccttc	ctgcttttcg	ccatcttctg	7320
gaacgtcttt	gcctatgacc	tgctgggtgc	cgatatcccc	cctccacctt	gacctgctcc	7380
ctagggccat	gtggctagag	ccaggcacca	ggtgggcctg	catgtctgcc	ctatatgaat	7440
ggacctctcg	gctgcttttc	cccacagggg	cagggggggc	ctgtgatccc	ccacccccag	7500
ctgaagccaa	gtcttactgc	agcccttgct	gcaccccccc	ccccaggct	ttccaggggg	7560
aggggtgtta	ctggggccagg	gctgggtgga	ggaccttgag	cgtagggccta	gaggagaccc	7620
cttccctccc	cactgacccc	aggacttgca	gtctgagcct	ttttggagaa	tgaataaata	7680
ttttacacag	caccaggtag	ctgtcctggg	gctctgcagg	cttgcttgcc	ccctcaccca	7740
agctgtatga	gacacctccc	ctccaggacg	gtctgtgctc	cctctctccc	ttcaagctct	7800
gcacctgact	tgagggtgta	gctttactcc	tggccatggc	cggaatccac	aggaggtggg	7860
gcccctggga	agaaactgtg	gacttccagg	tcagtccaag	ttcagcaatc	gtttattgtg	7920
ctacaagagc	ttccaatctc	ctgttctttc	agcctccctt	ccaaacgcaa	gcccacaaca	7980
gagctaggtt	cacttggcct	gggtcctgta	attcttggca	ggtccttggtg	ggaggcccag	8040
tcagggtgaa	ggtcaggcag	caggctctggg	catacatctg	ccccccaagt	ttgcacaaga	8100
cgaagagtct	acaactctaa	gacactcact	agtgaggtcc	acaaggggtc	tggggtcaag	8160
ttactaacca	ccagaaccat	gatggggccc	tggatgacct	ctcaggcccc	gctcagcccc	8220
gctcagaaag	gcagtgaact	cttgcacctg	ggctagaaac	agcaagtagt	gcttgagtgt	8280
gtactggagg	gttagcaagc	taggaactgt	agccaccgcc	tcttcgtata	tgaaagctgc	8340
aaccccagaa	acctctgcct	ccaggaccac	tgcaaagagt	gacaaggagg	ttctggggat	8400
agcttttttt	gagctcctag	agccccacc	caacttgggc	aaagatgcta	cctactcact	8460
cgaggcagtg	gcagaaacca	cgcagcgaag	tttgactgcg	gcgccagat	cagcagaaag	8520
ccgagttctg	cttctccaag	tccgggggtg	ggggtgttag	cggccaacag	ctgatgcagc	8580

agcgggctaga aatgcctcaa cgacagacca tgggggtcaaa ggggtcttggg gcaggcgctc	8640
tccaaggaaa ccagccgggt cggcccttga agaagctcct ccccatgacc gcgagaaggg	8700
caagagtcca gccgtgcacc cgcagctact tctatagcaa ccgcagaagg tccttcgggtg	8760
tcagcgtgtc acgtgcttgt ttctcagcca ttaatggccg caggcagctc ctcccgattc	8820
ccgaggaagc ccctcgagag cgggtctgatc tgcgggcctc tcttctaccc tcaattccgc	8880
cttctaccg gcgggtccaga cgttcccagg ctgtcgcgtg agcccggtg tgccctgggtg	8940
gtgcgctagg acccgggcct agtgaccggc cctcggaagg ccacttctca atgaccgctg	9000
ccacgttcgc ccagcttggg gccggagggg ggggcttgag cgccgatggg cggagccttg	9060
gtctgagcag ggtaggtctc gggttactgta gggcgggggc tgtagagcgc ggcggaagct	9120
tctcagagct aagggaactg gctcaatcca cgaggctccg cctttgctca cgtgtcccca	9180
tgccagctcc gccccacacc ggaagttccg gtggcggtat gccgaccggg cggagctgat	9240
cgctgcgcgg gctgcgagat ctaggtggcc gggcgcgagg cccaagccgt gccgcgcggc	9300
gccatgaagg gcaaggagga aaaggagggc ggcgcgcggc tgggcactgg tggcgcgggc	9360
agccctgata agagcccgag tgcgcaagag ctcaaggagc agggaaaccg gctcttcgtg	9420
ggccgcaagt acccggaggc ggcggcctgc tacggccgcg ccacgtgag tgcgggaggg	9480
gcgggggcgg ggcccgatgt gcctcccagt tttgtgatcc cggacagggt ttctggcagc	9540
ctggactcga tggctcgtc gctcccctgg tgcccgatt gctgttcccg agtctagggc	9600
tgatttgact gggaaactggc gggcggggatg cttgattaag taagactaag gatgacacca	9660
ggagctcttg ccccttgaga tttccaagcc cagagcttgt agaggggttg tgggatttct	9720
ctcaaaagca gttggaggca agaaagcatg gccagaatcc agagctcttg ataattcgga	9780
aatctccagt tcctctattc cagccatggg cccccaagc tggagcattg accagatcat	9840
tggctttcag cttgctaaat gagaggggat caactgagcc cacactcttt gacgcctggt	9900
cttggtcccc agactcggaa ccacttgtg gcagtgtact aactaaccg ggccctgtgc	9960
tatctgaaga tgcagcagcc tgaacaggca cttgctgact gccggcgagc cctggagctg	10020
gacgggcagt ctgtgaaggc gcacttcttc ctggggcagt gccagctgga gatggagagt	10080
tatgatgagg ccattgcaa tctgcagcga ggtgggctga caagatgcaa ggttgagggc	10140
accttggggg accaggattg ttgacaagt taacaaaag tgtccactcc cctagcctat	10200
agtttgcca aggagcagcg actcaacttt ggggatgata ttcctagtgc ctttcgcatt	10260
gctaagaaga agcgtggaa cagtatcgag gaacggcgca tccaccagga gagtgagctg	10320
cattcatatc tcaccaggct cattgctgct gagcgagaga ggtgggtccc tccccagca	10380
gacactgagt caccagcaca agtgttttta ttgaaggaca acaaccggt catgggaagc	10440

ccaggaggaa agaggcctga gatggggcgg gctctgtgat ttcctggtca gggccatcac 10500
 caactagctt ctgggtaacc ctcagggaac tggaggagtg tcagcggaac cacgagggtc 10560
 atgaagatga tggccacatc cggggccagc aggcctgcat tgaggccaag cacgtgaggg 10620
 tgcccctagt ccacatgtgg gtcatttgtgt gtgtgtttgt gtgtggcatg aaaagtgtcc 10680
 cagccttttg tggatctgtt ggcccatgca tgacatgagt gccctggtgg acagagtgtg 10740
 tgtgtctgtg tggagacagg agttgccac acatggcact caactctgca ggataaatac 10800
 atggcagata tggatgagct cttctctcag gtggacgaga aaagaaaggt gaggaccccc 10860
 ggaagctcat cggctgctgc tggatatgtg acccccacag cccctccag ttctagacat 10920
 tttccccttg tgctatagaa gcgagatata cctgactact tgtgtggcaa gattagcttt 10980
 gagctgatgc gggaaccctg cattacaccc agtggtatca cctatgaccg caaggacatt 11040
 gaggagcacc tgcaggtaag gagggcccat ggagaccggg ccatggagag ccttgggtacc 11100
 cggatcggtg tactcaggcc attgaacact cttgactctg ttgcagcgtg tgggccactt 11160
 tgaccctgtg acccggagcc ctctgacca ggaacagctc atccccaaact tggccatgaa 11220
 ggaagtcatt gacgctttca tctctgagaa cggtgggtg gaggactatt gaggcccat 11280
 gtcctgcctg gcaccctggc ccaggaggat ctggagacgg aagctccagt ccctgtatag 11340
 tttgtgtccc tgggcctgcc cccatcggcc ctgctgatgg gttctgaact gctccccttc 11400
 tcagcatacc ccttgctgga ccatgagcct cccttgtccc ccttctgggc tggagagtgg 11460
 gtgaggggtg gctgaggttg ctgctgctgc cactgtcctg taataaagtc tgtgagcact 11520
 acattggcat gtgctggtgc agtgggcttg ccagttgcct gtctggctag ccaaggaagg 11580
 tagaaatgaa gacactggtg tocagattga gtgtggcatg ccaccaccga tcaggaaaat 11640
 acagtacctg ggagaaaaga aaaaaaaaaat tttgattcat ggctttttga gttagcccaa 11700
 cccccacctg ccagctactg acttcaccag cctggatggg acattctaga ggccgtgctg 11760
 acagggttag agatgggtat atttccagca gccaggccaa tgttgtcttg ttaggggtgga 11820
 actcaggtgt cttctcagga gggtagagga accagcgtg gaggcaggag tgtaagggtca 11880
 ctatgtgtgc aatggactat agacaccaga ttacacaca cacataacac acacataaca 11940
 ctgtactgac cttccgacca tagataacct ctgagaaacc aggcccatgc cagtgaagg 12000
 gtaccccaga tccagctcct gtgggggtcag ttatgaatag ctgatgaaca aaatatatat 12060
 gggctcccc aagtttgttt tgaagtagcc cctacctgca attccaaagc tgtaagcagg 12120
 ggtggttccc aggagacgga atggtggcgg agagtagtgc tggaagaggg atgcccactc 12180
 agtgaagttg ttgtctccaa aaaagtacag ggtgtctgcc agcagagagg gtgataggca 12240
 ggagtactgg caacagtggg ctcccaagta cccactccc accgctgcct ggggtctggct 12300

caccattgcc tagggatgca ggatcctggg gctgcagcag ctgttccaca tattcctgga 12360
 agggcaggtc cactacaaga agggggagcc agtcgggact ggtcagttct gcctttgaga 12420
 tgtgaaggca aaccttgca ggtgtgtagc tcacctttct ggtaggagta ggtgttggt 12480
 gtactcaagc gaacaatgtt gtccccgaac gaggctagca ggttttcccg ggaacacagg 12540
 gcccggaaact tctggggatg aaggataagg taactgagca cttggccaaa cagaagcagt 12600
 tgcccacagg gtacgaatac tcaccgagtt gtccgtgagt ccttgcaaga tgacgggctt 12660
 gaggaaggca tagctggggg gaaagaggag aatgttgca tgagtttgaa ctctgcaggg 12720
 gcctctccta cacaccaggc agtgagacct ccgcagccaa agaccatgat tggtagtatg 12780
 cctagaatca tgctgagact tggggcaggc acacatactg ctgcatgaat tcggagtacg 12840
 tgatgtgtgc ccgacgtcc actgtgcaac gtcctctc cttactgtct gctgcaattc 12900
 ccagcccatt ctgtttcctg cagggacaac aggtcaggcc agtcaaagct ttaagttcag 12960
 gtcctctccc gccacccatt tcctatggct caccatcccc ctccgccaga ggcaggcaga 13020
 atcacagtca ccatcatcca tagtaaaaag agcaaaagca gaccgcgccg cccagccgcc 13080
 gccatgaaca tacgggtgca ggcgcgcagc acgctgggaa ttctgtcgac acttgcgtaa 13140
 ccacaagcca ttggcttatc atccggactg ctgtagctac ctgcaagcag tagccagccg 13200
 gccttggaag tctctctctt cttcatacaa gcgcattgta cacacacaca cacacacaca 13260
 cacacatcca cacaacatcc ctatttaatc atttcctcaa ggcggaggga aaattgtatc 13320
 gagcagccat tgactactgt gcgttaccat ggctacagcg cccctaccgg aagttcccat 13380
 gtgctagaaa gttcccaccc aaggcgcaat gaaagtgaca agtggccaaa cagcagctct 13440
 ggatttttat tgggttaaaa agttgtagcc accttccatc ctggtacaca ggatgggtaa 13500
 ggggctgtga gcctagtccg gtggacatcg ggatcgagct gaggacgctg catcagcctt 13560
 agggaaggcc aggtcaggag tattcacaga ggtggccgca gcctgcaggg cagtgggagc 13620
 tgctttccag ccatttcatg atgtgtgaa gatggccacc atgactgcaa ccctgacacc 13680
 acacgaacag gcctttaacc acatgatggc agacagcaca catgctggca cagcgggtggc 13740
 acctgggaat gggagggtg ttcagctggg gagctctggg ctgcctccct tccaacctcc 13800
 ccaactgcac catcgacact gtcacagacc cagccgcggc tgctcatggg ccgcttgca 13860
 tgactacagt tgacatgcag cgtgggtggag gcttggttga ggcagctaac tgccctgctg 13920
 gtgctcagct tcaccacctc attggacaca ttccagaggc agaagcgctg cagcagatct 13980
 atgtaggatg tgtaccagtg ctcttggtgg ggagggacag gctcagggtta gtgtttccag 14040
 gaacactagg gtgtgtgtgt gggggggggg gggggtacag tggaaaagaa acaggctcag 14100
 atgagggaac cagcagccag tctggggaga ccctccccct gcccgcgcc gatccatcac 14160

acacctgggt ctgctcatcg atgtcctttc gcactctctc acccagcacg atgagaacag 14220
 acacagccat ctgcacgtcg ccttgctcag cgtagaagcg caacatgtcc cgtaccaata 14280
 cactgaagaa gtcagggtggc aggcgggtgt catagagtgc atgtgagacg gagaggaggg 14340
 aggaggagga gtccaccggg gctaataaag ctgtgtctgc ctcatcgccg ctacagtgcg 14400
 gagagtccgc cttgtcctgc agatgctcgg gccagatgg tgtgtctaca atttcatggc 14460
 gcagtgggaa ggctcttgtt ggcagcacat attcaggctc ctcggtggg aggcaggggt 14520
 gaggaaagat ggacctgagt gtaggcttcc cgctocactc agctcctgct cacagagcca 14580
 tgttctgggt tggaccctc tatagctgac ccagcccta ctacagtgac acatgttctg 14640
 tgtcaagtgg gtacagctca tcgtcctctc cttctacgtc acccagcagg tagtcagcag 14700
 gcacatcact cccttctgtt tcctcattat ctgccagca ggaagggag gtcattcagg 14760
 gagtggcacc ccagcctttt cccatcagca tccctgccta cctcattgg tgacgagtgt 14820
 ggctgaagag tcaagcagcg ccgcatcact ccgtgtgtcc ccttgctgc gatccagccg 14880
 tgtctcactg ccagccccg ggccatata cttcagattg aagctatagg aagggaaggc 14940
 ttgtgggtgg ccagccccg gcagtcact ctggggaggg ggcaggaagt agccaaggat 15000
 ctacctgttc atgagcggta ggccgcagga gctgcctttg ccgacgctgt ggttgagatt 15060
 ggcagaggat accaagccag ggctacagta gataatccgt aacatgggtcc atgtctgtgc 15120
 cacctagggg gagggaccaa ccactcaggg gtctccatct cctgagcaga gtcataacc 15180
 aacctatggc ctgatctact accaccagg aaactggccc ctggaaagag cagtgtccac 15240
 tctacagtag cacatctgtg aggtgtctgg tgctaagatt agcaatggaa cagatatggg 15300
 gaagagagag agcagaaggc agctccaaca ccaagaacga ggtagaagag cgggctgtag 15360
 gcacgggact ccttggcctc caccctgagt ttgggctagt ctggttccca tgatctttac 15420
 actgtcaaact actgtgacag aggtgtccac tgtgactgag ctcagggggg ttgtgggcat 15480
 gatgccccaa ctacagcagc agtctcggac cctgggtctc cagcttcaca tacctgattg 15540
 cggccaagct ccgagccac ttttgcattg tgatcacaga gctcggctag tggccggcca 15600
 gccagaacat aacgttcagc tgtatccacg aaccagctca tgctgccacc gctagactcc 15660
 gtctcaaaga cgctcagtg actggaagcg aggcagaaa aaggttctgc aggtccaac 15720
 ttgcgcttga agaagatggg gtggcgccgg tcaccagcat agggtttacg tccagattca 15780
 gcagccacca ggctttcctt aacagcaaag gccaggtccc caaagaggcc atagcagaga 15840
 ccctcagggt tggcacgtc aacaggctga ctggcatccc gaaacaggtg ctggcataac 15900
 gtgtgtctc tggagccgga gagcaggaaa gaggggtcat gcggatgacg ccaggcaatg 15960
 cctgttgtga catcacgatg ctcttcaaac atagcagctg gcacgaaggg gcggcgctaca 16020

tcccatacat agatgttgtg gtccaccatc atggaacacg tggccagggtg gtggcgccat 16080
 tcaggccgcc acttgactcg ggccactgaa gcgattgtct gcacacagtg tatctctttg 16140
 gcacgatgtg tggatcatgtc ccagaccttt accatcttgt cacgtccacc cgtggccagc 16200
 cagccccctgc ggagaatgtc ccactcaagc tgtcagtgac ctctcccatg ggaaaggaga 16260
 ccctgccaca cccaaaggca tcgttttagcc tccagcctta aggcccatct tctccccaca 16320
 ccacacctgt cctctgggtg ccaatcacag cagaagacag gccatttgtg ggctgtgaac 16380
 atcctctcac agcggtcggg tcggcggatg tcccagagct gtacgttgcc attctcaaag 16440
 gtggaagcaa aggtaaagta gtcacggata ctgaactgca catcccgac gctctcagac 16500
 tggcctgtgg gcacacagtg gtacagacat aagtgagggg ctggcccca gggcgtgggg 16560
 gagcccttgg agtggctcca accatgttat tcaccagcct atagcccagc ccagaaacac 16620
 aggactggag aaacaccacc aaaatctgcc taggccagag tatgccaggt agtttcagtg 16680
 cagactccca gcctcagttc cctagattct gcacaggcaa agctgcagac ctgcctcaga 16740
 aggctatgtg tgagcactgg aggcacagga gtcacccctc gttggctgtc tatatgtccc 16800
 aggccaaaggc ttgtggaaga cccaacgaat ggaagaggag gctcctcgtg ctctgctcaa 16860
 tgccctgcca gctgcctagt tagcctcgaa ggaccgaaga gctcagagac catacccacc 16920
 ttaaaagatc gccaaacaag gagcctgcgg gtacagtgtc cagtgccttt ccctgccaca 16980
 gacagctcaa ggtctgttgt aacccccaaa ttctccaggc cccaaggctc tgcctgcgca 17040
 tcatttggtc tcattaaaga tctgagagag gggaaatgagg gctgaatgca gacagcttgg 17100
 cctgccatga aaggggcgct gtgtactccc cacctcactc tgagtcccg tctgagtcct 17160
 gaagacacaa ggctccgcca ctactacgg tgtgtctgaa ggctccgcct tccctattag 17220
 ccacaccac caccagtct caatttcac tggcttgacc tcctgcctcg ccccccccc 17280
 caccccaaca gcagtgttgg gcttatcctt ctccagggggc caaccgttcc catttcacac 17340
 actctttctg ggtgttccg cctctgaggg tccatctgag ggctctgat ggacacacat 17400
 ttcagcctgc ctgtcaggtc taacacctac agagatccag atgtaccccc aacttcctct 17460
 gtgggcctca ccagagaagg tgctgacaga gtccttcctt cggagggtcaa agcacttcat 17520
 gaagccatcc tgagagccac tgagcagcac atgggcctcg gtgggatgga aacagacttt 17580
 gttcactgtg cgcttgtgtt ctgtgaacag ctgatcctgc ttgttgctgg atggccgacc 17640
 taggttccat gtaaccacca caccattggg ggctgctgtg gccagcagat tctcatccat 17700
 ctggtgccag accacatcag cacagctcag gttgagcgag ggcttcctac ccacacgtaa 17760
 attcagcttc totacgaact gtcctcctc aatggcataa atcttgaaga tgcttcggcc 17820
 tgccacgaca acctgggcag cgtcggcgca cacgctgatg gcattagctg gcgcgtccag 17880

atggcagtg c atggttcggc ccgtcagagc actgccgctc agggctgtgc taactcggga 17940
 catcttttcc atgtctgcac aggtatcaga ggcaggagga ggtcagtgaa gtgtgctgac 18000
 ctggtcagcc atggtgtcct aacaattcag gttcagtcct ctctcgaacc ttcggcctga 18060
 gatggtccat cagccagtcg ttatcttagg aagggtcaca ttcattggag ttctccagaa 18120
 ggctccagag ctaacctgag gttcgtcagc accatatctg acgtctcccg attctatgtg 18180
 gtttaattctc catcagtgag atcctggggg cggggagtggt ctcttgactt cacagcccca 18240
 tgctccgag gattgcgggg cttccaagag gaaaaacaac cgagtggaaa gggcagagaa 18300
 gctagatggg tatggactga gttcatcac cttggaatgc agaagagagg ctacagcaat 18360
 aagtgcaggt gtctaattgc accgtgcagc ggcaactgag ggaccagaac cagggtgctag 18420
 aaaagcctct gcccgcgcg acgccagtc tttaccgaaa gcacagagtg gtcttcagct 18480
 acagtaccgg aagagactca cctactgcc gccattacc ggaagtagtc ctttcttatt 18540
 ctcaactcgc cggaagtga tgacaactcg tctttcacac cataagagtt ccgtgtctgc 18600
 gaacagcgct tctgcagagc aaacggacca gaattgggct ccctggtttt aacttggtta 18660
 tatatgtccg ctattgatcg tttggtagg gtttctgttt tgttttgttt tgttttgttt 18720
 tgttttatgg ttctggaaat cgagcccgag agacaagttc tctaccaact aggctatgtt 18780
 tattttaatc tcaatatctt aataccagta cttaaacctt tcctccggtc ttggagaagt 18840
 aattttcttt ttcttttttt ctttctttct tttctttctt ttttctttt tctttttttt 18900
 ttcttttttg gttttttgtt ttttgtttgt ttgtttttgt tgttttgttt tgttttttgt 18960
 ttttcgagac agggtttctt tgtacatccc tggctgtact ggaattcact ctgtagacca 19020
 ggctggcctc gaactcagaa atccgtactg ggattaaagg cgtgcaccac cactgcccgg 19080
 cgaggagaaa ttttctaagg agaccttgag ttccaccact ctgccacagt gccacttccc 19140
 ctacggaagc actagcagcc tgggtgccttt tctactctag gtcattattt cactgtgtat 19200
 aggtctcaga gacctgacc tgacctttct ggctcatctg cagctctacc tttcttggtt 19260
 cctcctttct cagcttggtt ctatcccagc ttccttcctt ctgagtattt tccttcttct 19320
 ttcacagttc atcagctgtg ttataaaacc atggcttcta cctatggctt ctaccctgc 19380
 agaaggtgtc tcctttacac ataggtaggc acttagctca aatgcctacc atcttaggcc 19440
 atccttgcca cccagtccta atgtctcctt tgtctactgt ggtttgatg attacacctg 19500
 ttttggtact ttcagagcct gacaaccagt catcagtggt ttagtagatt tagtctccat 19560
 tgtaagaaat cagtgttggc tgaggagcct caagaccaag ttctcagcat aaagaaagat 19620
 tgtttgcccc aaagggacaa aggctaggca atagggacag acagcagata gagggtcagg 19680
 gaggagaggg agtagggata tttgaccaga ggggataaag aactgcctct ggatagaaag 19740

gagacagaca tggcctatag acaaagacaca ctttatgact ttataagtgt aagagggaga 19800
 aacaccatgt tagaatgagg tatttaattt tgattagaca ggttaattaa gtgaacccat 19860
 gggggctttt gattgctgga cctcaatact ttgatagctg ggccttcac gccagcctca 19920
 ggagtagaag tggccaaata aggggaacata ccatgggtggc tagctctagg aatataatct 19980
 aagggttttta gcaagacaga aggcattgca gagaaggaca aagctggcta gcagtcctcc 20040
 cttcagcact tgccctttct cttgaggtgg acacaatagt cttgatcttc tagcagctctg 20100
 tgattattta tttatttaaat ctttatttga gtgtgtgcac atgtaaatgc aaaaatgtgc 20160
 ctgtgtgagt ttttatttgt accaatgtgt gttcaggttc cctaggcggc cagagaacat 20220
 catatcccta aagctagagt tactgggcag cttaaaggca tctgacgtga gtgctgggaa 20280
 actgacccca ggtcttctgt aggaacagta gattctttgc actaccgagc catctatcca 20340
 gcatcactgg attatttttag aacatcacaa tccctcactt agtgaccccg tgcagaagat 20400
 aaggctcatt tctgctcttg gatagcaaga tcctagatcc ttgttttctt tttctttttt 20460
 tttctttttt tctttttttt tcttttttct tttctttttt ctttttcttt ttttttctga 20520
 gacaggggtt ctctgtgtag ttctggctgt cctggaactc actctgtaga ccaggatgac 20580
 cttgaactca gaaatccgcc tgcctctgtc tcccaagtgc tgggattaaa ggtgtgcgcc 20640
 accaccgccc ctagatcctt tggaagacct acattttgac atcaaacaca ttgttccctt 20700
 ttgtgcattc aggacattag agaacttatg ttcgtcagtt gctaagggtc cagtccagcc 20760
 ctgagcctcc aacctctcct gtttatggag tacctctaata gatttaggag gcaggaagac 20820
 ccagagacct gggccatcca gggcagagtt atcccttcca acttggtcag gtgacaaaaa 20880
 aataataacc cacttgtgtt ttacaactt ttttaataata tatttttata aacaggtcac 20940
 gtgataaaat agcacaagaa acacttacca aatataaggt tatactctcc gcatatacag 21000
 gagaatgagg tcgttatgta caataagaaa atgatttttag gggttggttg gttttgtttg 21060
 tgttcctctc tccccttaaa tttcctcct acagtcattg gaaatatcac agcttcagtt 21120
 gcattaatac tttgggcaaa tggacagctt cccctccct gttgggggac tgtgggaaga 21180
 ggggctgaca aatctggctc ctgaccactt cagcctgggg cctccctggc caacactgca 21240
 ggggcaggga actctttggg ctgttgaatc ttacttttc gtcaacagca tctctctctc 21300
 tctctctctc ctctctctct ctctctctct ctctctctct ctctctctcc ctctggctct 21360
 ctgggaatta gtttactctc ttccaaccag ataggggtgt cccaagatta gttgtgggtg 21420
 ctgtgtctcc tgggggtcaa ggttgggagg gaaggttccc agtcccttcc tcggtattgt 21480
 cacctatatg ccaggatccc ctaggacttc caaatttctt gagacctgac agaacatggt 21540
 cacaaccct ttgggcaggc aaagggtttg gaagctggga gaactaaagg aggactggga 21600

aggagccctc cttccctggg gtgtgaaggg ggtgctggcc actgggagat cagaatcgct 21660
 ggtcagtact gaccctgact catggccagg ccctctgcag attccacctg aggagaccta 21720
 tccagggaga agcacagtgc tgagaggcct gagctcagct ggggtagggg ctcaaatggg 21780
 tgggctgata ccaaggcctg atagccctgt ggggtggctcc agattggggg ctgccagggtg 21840
 cctattccac ccttgcaactt aggtggatgg gcaccgggta ggctaattgtt gattgttcca 21900
 aggtctgcgt cctcctaaag ctgtcccaac ctggccaggg cccaaggatc tgtaggccag 21960
 ggggaaagaa ggagatacgg gggtaggggc gcaggataca gaggacaaga actgacctat 22020
 atggcaacaa ggaagaaggg aagccactag gtcagggtggg catggggagt tgggctcctc 22080
 agggggccat ggctttggca gtggcctgtg gcgcctcata ccctttggag ggctccaggc 22140
 gcggtgcagt ggaagatgtg ggaggccctt gggggctctt ccaggccccct aagtcctggg 22200
 ggaccaatcc ctccatccc ttgggccttg cccgggatgg ggggacagcg aggggaggca 22260
 gagagcagaa tgaagtgggg aagccagtga gggcatatgc actcaaagca tccccaaaag 22320
 gacctagacc ccagaaggc gggcgagtga tggcgggcggc tgtggcatgg tggagggcgg 22380
 gcgtgcagtg cgggcacgag gggcatgcac aaagtccccg agtgtgcgtg cgtgcgtggg 22440
 gctggccccct ctgcctcttg cccagacag ggaaggagca gttgagcaga ggagcaaggg 22500
 aggcacgggg cgggatttcc ctcagcttgc cccgcctcca gcgcagggcg gagaagagat 22560
 gctccggcag ggatggaaga agggctactg cccagggcgt ggccggttcc cgcggccggg 22620
 tcgggaagcc tgcgctattc aatgacgagg cagcgcggca ggtgctgcga gaagtacttg 22680
 aagagctcgg gtgtggctcc agggcagttg gtcagctcca gtcctccag ctcttgcaagt 22740
 tgcacgaggc cagatagccc ggtggtagtt aggagcgggc agcctgcaag catgaggcac 22800
 ggggacagcc gctcagcagt tcagttccgc ctggcgggag gactggactg ggctgggcag 22860
 gtgggaaagg ctgaagagga caatgacctg ctaggggtgg attctactgg gcttgctgaa 22920
 tgaggcggga cctgtaccgg gcgtggaggg gtgggaatct gacaggatgg gcgggacctc 22980
 caccagggaa ggatgtgttc agacttggcc tgatgcaggg tagatctgag cagggcgtca 23040
 ccaacagggc ttggattcac cctctaacc gctacctggg atgcgatttg ccaaacctg 23100
 aatgaatct atctgcacaa aggacctgat gggggtgggg gtcagagtga gtgggcatcg 23160
 ttgccatga ggcattgcggg ggtggggggg gtctcacctg ctagagacaa gagtcgcaaa 23220
 tttctcatgg ctaagaggtg cttcaacccg aagtcctgca cctgggcggg agtgagagga 23280
 ggtcgtgtta acttatttcc acttgcttgg gacctggagc cactagaggg gcaggaggtg 23340
 ggggttcaaa atggagattt ctgagccca cccccaccc ccgtagactg gaagcaccaa 23400
 gagggccttg gggaatctgg aaccaatga ttctgcttag tttggctgct ggttctagag 23460

gactctgaat gaggcggggt gaagttcaag gataatctgt actggtcctc caccctgtcc 23520
tgtcccgcca ccaaccacgc totaatttcc agatccctgg gggcatggag gtacctggca 23580
gcaccatcgc aggtagaggc tgcggagaga cgacatggtg gacaaatagc tgagaccagt 23640
gtccgtgatg cgtacacacc tgtggttgca ccagaagggg gactgcccac ctttaaataca 23700
ctttacccca tccccaccta ggtccttccc cccttttcag aggtgtccct agagcaagtc 23760
tctgcccac acaccaatgc cctattcttc ccctggtcce acccgtggc ctcccccttt 23820
ggaatatggc ttttctcca gttttgggca ctctattcta tgtggttttc cccatccccg 23880
aagcctgtga accgggcacc acccactcta ggcttcaggt tcctggagat cccaccagta 23940
cccgctgga gtgtagggat gtgtgcacct gtccagcacg agctcctcca gtcggtgcaa 24000
gtcgcaggcc acatactcca gtgcatatc ggtgatccga gggcaccagg aaaggtcaag 24060
gctgcgcaat ttgcgcaggc tctcagctac aagctccact ccatcatcag tgacctaga 24120
gcagcctgag aggtgagtg aggtgaggtt gggcagactg tgaccacat tgaccacccc 24180
atggttggtt atctcccagc aggaaagcag gcgcagcgtg tgggtgctgt ggcctggcg 24240
tgctgtgaag taggccaggc ccgtgtcagt cacgtgatag gcctgaaggc ttagctctgc 24300
caggttgggt agaagttgtg agatggcagc gattgcacat tctgccacat tgatgcagtc 24360
actgacgctc agggaggtga tacgggcact gaggctggac cacaggccag cctcggtgaa 24420
atcgttgagc cctgacagct ccagacgcac cactccctgc atctgctcca gcacaccta 24480
caggcaggta caaatcagac taagtcagct ccctggctca cgcctgcccg aagaagtttc 24540
tgggattctg gaagcgggtg tctctcaggc gtgggcttgt gcaagcatgc atcaggctga 24600
actccaagga atgcacttaa actgcttgtt atactttccc aatattccaa ggggggaggc 24660
agggcacagg ggacacaagc cagggtgtgg ccagtgcagg tgtttagaac agatgatttg 24720
aatattaaaa ttacaatcca ggtccagggt tgtggggagg cagggcaggg tgcttctagg 24780
agactcttag ctacccctcc acacaccagc ccagcagccc attcaccaag gcagcttcac 24840
aagcaagcag acaccacat gtaattgact actctcttgc accaaccttg acatcttaat 24900
cactttttta caagcccaca ttttacttt gcattaggcc ccgaaatacc atccttgctg 24960
ctttttatcc tgtatacacc tgagtatctt ccaggcccct caacctccca tgtccataag 25020
gttgcttctg catgttcccc acccccctcc ccagacccag acccatctgg tcatagccaa 25080
ttttgggaca cacctctaga ccagcgtcag tgatggtgga gcgcttgaga ctcatggcct 25140
tgactccctt cttagagaga gaatagttgt cgatgaactc gcagatgtct aagtcggaga 25200
caccaactag gcagaagccc tcaaaacctc tagcagcaaa gccctgcagg ttcacaaact 25260
ccttctcgcc tccaggcagc acgttgtaga gctccttggc atgcagcaca ggcgtgaggc 25320

ccgcccagaa cttgggctgg tagagcacac gccgccaagc cttgcacacc tgagctagta 25380
 tgcacttctc acatgctgag aaataccaga aaagcccatt aaggatcttc tcgtctgtgg 25440
 ccagtgggtgg ccgctccact ggggggtccag aactgggacc cgaggctggg cactgggtg 25500
 gggggcatgg cccccggcc agagcaaccc gggatagtgg tgtagccagg cttggagggtg 25560
 ggagggtcgg tgggggtggg ggctggcaag gacggttctt ggcagcagggt gtgcccttgg 25620
 tgatgctggc tgcacctagg ccgttgggct ggccaggcag cttcaccaga ccgtttcgag 25680
 gcaagcatgg aggcttgggg tcaccatcga taccggggct cgacattttt cttgcacgct 25740
 ctgtggatga gacaggagga acaaataagt gtgccaggct ctggagggaac tggagagcta 25800
 accttttctc atcctcgagg ctctcacc aaagcatttc caggaccact taacagaaca 25860
 gatcctagta tgcgcctttc tctgtctgca caagaggcaa gctggccaca ttcacagcta 25920
 gctcaagtgt ccacatctc tctctgtcct attcactggc tcagcctgcc ttgctcaaag 25980
 gtgctttctg atgctctgat tcccatcgag gcttggggag atggaatcct tttttttttt 26040
 tttttttttt cgtttttcaa gaagacaggg tttctctgtg tagccctggg tgcctgttag 26100
 accaggctgg ccttgaactc agaaatctgt ctgcctctgc ttcccgagtg ctgggattaa 26160
 aggcattgct caccatgccc ggtggtagat ggaatcctta actgttctt tcacctcaa 26220
 tcctctgatg ccccttctg gagtttatct tgaggactag gatcggagaa ccagagaag 26280
 aaagggtgct gaatagccct gagtcacaga ccagctctcc aaaatctgtg cacctagtca 26340
 gtggtggcac atgccagtgc ttgggaggca gaggcaggca gatctctgag ttcaagagca 26400
 gccttgtaca gagagagttc caggacagcc aggactacac agaggagccc aatctcaaaa 26460
 gagaaaaaaa atctgtatcc ctataaaagg ctctgaggct cagtcgcac cctgtatag 26520
 ctgactgtgc taccagcaca gtgatacttg gacacatgct tttcatatt taaaaattt 26580
 aaaattttct gtgtacgggt gttctcttgc atgtgtacca catacatatc tgtggaattc 26640
 agaagaggac atagatctcc tggagctgga gttacaggcc tctaggctct ctgtaaaagc 26700
 agcaggcacc caagcactga ggcaattctc cagcctgctg gacacatgct ctaatcccc 26760
 acccgagtca acgcaaacag tgctgtgctg tctctgtaca accaaggaag tagaggccaa 26820
 gaccaagaga actatctgac taagctcatg gccgatagtt aagtggtaga gctaaaattc 26880
 ggatcagaac tggaggagga aggtcatatg gaaggccagc atgaccaaca tgtctgaatc 26940
 actccctccc tgttcccctc agactcccag gcctgtgact gggttgcaga ggacactgtg 27000
 agtggacagc acggagggaa agagcagtga gactctcata atgccatgat ctctgtctgc 27060
 tctgcattcc agacatcaca cccaccacca cccagcatcc agaaggatgt gatttttcca 27120
 tggatttcac ctgcagatga aatgctcaat gcaacactta aaacattcaa aggcaatact 27180

ctggactcca gatggcccct tctataaatg agtgctcttg gctagcctct ggtgagtcca 27240
 tttcctcaac atgaaatctg ggaggtgaca ccacaaactc cttcctgttg gagggatttg 27300
 cagtctatgg tagtgtaaatt cctgagtgccc agagctgacc caagggcaag gcagggtttc 27360
 cctttctgtg tgtataggtg atcccaggat gagcaggatg cacacatgaa ttgtcagttt 27420
 ttgccctaca atctccaggg agtcaacaaa agcctctacc aaaccttcag tcataccttc 27480
 atacctggct ttggtggctc ataaccttgc caggagacct ggagcaatac caagggctac 27540
 atgaagaagc ttcccaagtg gtgggcactg aagtagacag accctcagca gagaaagact 27600
 catgttacct tttaggttga gacccctttc tggagctaca gggccaagga aattggctac 27660
 tctttcttat actctggctt gacacaagag aagtagacaa gaactacctt ctgcctttat 27720
 tttgcaattc tgtctctatg gttccctata ttgcacatgc aatctgtgct agtttccac 27780
 ctctgggaag ggaaagtccc gggttaagct ctcccagaa catggtcctc atgatgaacc 27840
 tagttctgtg tctaaggatg gccattggca gtccacccta aggccctcag tactgtctcc 27900
 tgattactcc ctttctcaat atgactctta agacctggg agtaggccag ctattcccta 27960
 aaggtttcag accttggtg ctctttgtgc tcaagaacag gcaatagtag ctacaaaaaa 28020
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa agccagtata gcaagtagca atgctaacac 28080
 ttgcaaatg cacactggtt caatcacata agaaccgag gctctgaaga gttatagttt 28140
 gcatgctacg gtaagggaag tctaaagtgt ggttacccca attccctgga gcaatatagg 28200
 gaagctggcc tgttgcgga cagcagcatc catgatggga aaggatgaga ttcttcctta 28260
 tgtgctccaa ttcttatcca gtaccacccg cattcctggg aactgtgctg aggggtggaga 28320
 gacagcccca gcctgagtac cagccatata ctaagtcttg gtctacctct agactgcccc 28380
 tggaacctca gccactaggc gcagaatcag cgctctcgga accccgcaag cagcaccacg 28440
 gacagctccc gcgctcaatg tggtcagccg ggctttctgc accagagaca gtggagcgga 28500
 ggaggaatga tcggtggctc cggcaaacc agctccacag gaagaggcag agggagaagc 28560
 cgcccgcttg gctaggatct ggaggaggct gcggaggcgg tgccagggga gaaataatcc 28620
 aagagctgtg ccgatttagg ctgaaaatgc tcccagccgc ggggtgtgcac gtggagcgcc 28680
 tgagttgctg ggggttgagg gggagatgca cgtggatgcc ccgggcatct gcagggtcc 28740
 tttctgcccc tctctgcagc tctggagatt cacgtttcca gaagcacacg atcgaccccc 28800
 ttcccacctt ccagacttgg gtcacatagg tgaatccagg ccagattcag tgtctaattc 28860
 ccccatccag gccctgtcc ttggtggaag cctcatctcc acaaagtcaa cctcgctttg 28920
 tgattagaga ggttctcact tattctttct ctttgacatt aaccaagcct gagaacccca 28980
 caggcgcca aaaatggaaa cagttacacc tgaactcaag ggctgaggct ttcagagaaa 29040

gctaagcaca cactggaaga ctggaacttt cttgccaggt gagcaagagg ttattgaggg 29100
 gttctaagca aggatcactc cacctggccc tgccatctca ccagatgatg acataaggtc 29160
 tgctcccagg tatcgttcta tagcccaggc tggccttgta ttacagcca tcctacctct 29220
 ctgggattac aaatatgtac tgccatgcct agctagagct ccttaatgat gctgcaactc 29280
 ctgcttccta ggaggggggt ggggggatgg atcagtggcc aagtgacca ggttgtctct 29340
 cccttcacgg acaagcacct caggagctaa tactgaggag aggccttctg gaatgatgtc 29400
 cgaactggca cagcactcag ccccaggccc agtatgtcct gtgcttggtt tggagtagaa 29460
 ttctgggtca tctctagact gcaggggagc ttgagatggg actccctagg tccatcctgg 29520
 gctaggaagg gacagtgaca acttcctcag ctgctgtgtc ttcacagcc cgtctcatca 29580
 aggtgctctc cgggagcact gactgtagtc tttgccctct ctagaggact cattgtggac 29640
 ataaggaaca tggaggggtac aatgggtctc ctgggaaagt tccagtactt ggtgatttcc 29700
 agattggcat tccacagagt ctgatgctga accacgagga cttgcctctt ttgtatatcc 29760
 agtcatcaaa agagtctcct ggcctcatct ccatccattc ctcttccct agccagacct 29820
 ctgctcttca ttctcccacc cccctcacc tggacaagtg gcggtgccct gcttgtctca 29880
 taggctttct cagtgccgag cttgctgcct gcctccatag ataggatccc ctagtactct 29940
 gaaaaagcac ctccaggtgt cacctggaga tgcatcact gtttctaccc gtggcctggg 30000
 aactccttga agacaaatag caccatgagt gtgtggcttc tgggtactgc tgaaggaatg 30060
 gacaagaaat ggtcacctga accattcagg ttcaggggtg cagctgcccg aggctgagag 30120
 ccttggtctg ttggcggggc aggagaagca caagaattta actccgggta tgcttgtggg 30180
 cagcatttgt tcaaaagaaa tagattgggtg ccagcaccta gctcccaga cccgctgatc 30240
 ttgaggtgca ggtactcagg gagaggagat cttttggcta agtccctctt ctgggggttc 30300
 atggctaagg aatctgactg gctgaccccc aaaatctgac ctgatcctag gaaaaagaga 30360
 caggagtcac caaggagatg ctttcctcca acatccctgc ctcatgactt cctagagcac 30420
 caccgttcaa aacaagcctc agaaagggat ggatctttct gatgctgcag agaactgcca 30480
 cttgtcccaa gaccccaaag cactcagact aggaacacat ctgttgtaac ttgtgtcttt 30540
 ttcttctgcc tcaactgggtc ctgtgttttc ctatgatttg actagtggcc tctgcacaca 30600
 gctatcttcc agcgggggtca ctgctgttac cctgaaatct taaagaccgc caactacaca 30660
 ctaccaatg tctcaccctg catctctctc agtctagga ctctttccct gagtctgtgc 30720
 acaggtcagg atgaaggcca ccttgagggc tggcttctgg cctgcatagc acctgccctg 30780
 atcagctcct caggtttgaa aaggacaacc aaaacacca tctccagttc agtctgtagt 30840
 ggcagccccg tggcctgggg tgacgaccag ggtatctgag attgggggca cagagcaggc 30900

aacaatgttct gtgacagtgc tgtgactcag gcacattcac atccctgccca ctttgacagt 30960
gtcaaccaca gcacttttct tccaccccct gctggagctg cctctggggt gtgtgtgtgc 31020
cagggacttg gggataggga gacccttggc ccaggaggta aacagaccca ggttgaacct 31080
aggtcaatgc tccagaagat caaactaatg ctctgtgacc aaactaaggg caaatagtca 31140
cgacaggctt catcttatca cgccccccct cccgctgttc ttagaatat gagaccttca 31200
aaaccaagg ctcttttaga cacctgccaa gtgaggccca cctgcaactt ctgcagccag 31260
tttggccatc tgaccactaa tggggacagt tcagggtgc tcactgacct cctgaccttg 31320
ctttgctgcc ttccccccc cccccagtg ccagtcctgg aaggttcatc agtgactctc 31380
agaggtgaag gcttatggtc aggaagggtg agtgatgcc accagaagcc ccttctagt 31440
cctctgaaat agagagccca atctctcagg gctacctaga tgcaactgtt ttaacttcat 31500
cagtgcgaga cgggacctgt tcagggtgat ttgggagAAC agtgagtgc atactgttc 31560
tggaacttaca gtatgggtga ctggggaggc aaatagttag ggtggcagga aatgaggctg 31620
ctgaatcttg ggctatctca acatcaccca cgttttcaac cttgactttc cctctcttct 31680
cccaatggct ccagaccgg ccctcaggcc acgtctaggc aggaatcga ggctccacca 31740
tgcccttgac ctgatctagg cagcccctaa tcatccaggc tcctgtcca acaccggccc 31800
ccccccccc cccctcccc ctgctgcgat gaggtgctg tccagcactt catcaaagt 31860
ttgggctgga acagagggtc ccgtgtccc aaggggcagc agggaggcct gagaagaccc 31920
tggtcagtgg gccctgggac aagatggcag tggaaagtcg gcctccagga gggagcagg 31980
gaagtggcat ctgccatctg ctctgggag gtcagctgct ggctcgggct gggaggggct 32040
agcaggagg ctgctaggcc agatccaggc ctcaacagt cagccttctc tcctgggatg 32100
acgggtgggg aggatcccc actgactaac atcctagaag gacatggaag ccctgctcag 32160
caatctcagg atgcccattc cccattttga agccttccaa tcttggccaa ggcccatttt 32220
ctttttctag aaccaagac atgccagccc actccctgg atagagctgt gggcggcagc 32280
tctataccag ctcatgtcca gaaggcgtcc aggacccta ggccacatat ccacatggct 32340
cctccctttt aattgtatca acacctgtca cgtcctccag tccgccggag gctcatagg 32400
ctggagtggg gagagcccag agccaagcca gaaggagtgg gggctgacgg cctgcaattc 32460
cgctcctcg cagccaccgt gcttcctctc agcaattccg tggtagcgag atacagcctg 32520
accgcgctgc cggtagcgag cggggcgcaa tcaggccggc cactaggggt aggggctgcg 32580
gaaccgctgg ctacagcgcc gaagggtgag gtggggaagg ggctgcgatt acgtaagccc 32640
tgctcaggcg gcggcgccg cggcgagtc ccaagtcccc accccgcccc ctctcattg 32700
tagccttcgc ggaggcagt gctcgacctg gaccctcccc tccccgcagc ctggccgcgc 32760

gcgcacacac accttacctc ggccagggct tctggctcat gccacgctgt ccgccgggcg 32820
 acctcgcgtc atggggagct cagcaccgga gccgcctagc gcggaggcg cagaggtgct 32880
 ggactggcta gcagggcccg actgcgcccc gcgccggcg ccccgcgcca gggccctctc 32940
 cgcagagttc cgagcgggagc tccccgggtgg gagaggatct gccgccctga gcctcccacc 33000
 ggggagggtg aggtgtgtcc cagataaata aggcgcttt gcagggaacc cggcggggcg 33060
 ctctccctc cccttccct cctccctgcg ctctctctcc gcgtctctcc ccgcgctcct 33120
 ccgcggccgc cgccggcgcc gcgcgtgccc cccgccggc cctaagagcc gcgccctcgc 33180
 gcgtcccg cgtccggacc aatccgggtt ctagagccag gctgaggtgt cgagaatccc 33240
 caaccccgcc ccgtctctct ccgcgcgctc ccggtgcggt accgacgggc acgcgccagg 33300
 cggcagcacc cgctcagcgg ccggtgttcg cacaataaca aaccccgtag gccgcgcggc 33360
 cgaagagcag cttcgagtgt cccatctgca caaagtgggt gcgaccaga atcttgccag 33420
 ctgaggagaa ttgcggtcct tggcgcgagc ggttggcagt ggggttcgg atcctgcgca 33480
 gatgcaactg gccctatgtg cccttgcttg gattgccacc tgtgagggcg aggaaagaga 33540
 agccgagtgg agacatcgac atcgtcgac ctccacacac acacacacac accctcctta 33600
 tggcccctag gattcatatc ccttccgac ctgttgcgag tcaggcagaa gaatccctgt 33660
 tctgggtgac atagtacac acttgcaagc aagtgactaa agcaagcgtc aatgaaaaca 33720
 caagacagtg gagtgatagt ggggaggggt tacttctgac aatgtcaca atgatctgag 33780
 acgcctctc tgtcccata gcacttggtg ggaatgggtc gagaatcgac ctgccttatg 33840
 gagttccaca ggttcgctct gagatgaagt agaggtcagg tgggtggcagc ctgtatggac 33900
 cagggaggggt ggtttgtgta aaaggaccca ttaagggctg ctggagagcc agacctctca 33960
 gcaggctaaa gggaccccaa tctgggtctc tcttgagacc agaaacagca gatgttgga 34020
 cagcagagta gcgtcttgga agagaaggag cttcaggggtg gaatgtcctc agatgagacc 34080
 aactgctgtg gagatgggag gtctgtctcc tgggaatgat gtgaggccta ggggcaataa 34140
 gaaaggaag taagctgtat cctagctggt gtgtgtgtgt gtgtgtgtgt gtgtgtgtgt 34200
 gtgtgtgtgt gtgtgtgtgt ggtgtatgat catatatgtg aagggcgaat tcgtttaaac 34260
 ctgcaggact agtnnccctt agtnagggg ttaattnntg naggcctggc gtataccta 34319

<210> 48

<211> 21

<212> DNA

<213> Artificial

<220>

<223> primer, human and rat origin

<400> 48

cacccaccag ccgcactggc c 21

<210> 49
<211> 19
<212> DNA
<213> Artificial

<220>
<223> primer, human and rat origin

<400> 49
gagcagctag ggttcaatg 19

<210> 50
<211> 32
<212> DNA
<213> Artificial

<220>
<223> primer, mouse origin

<400> 50
ggttaattaa ccaccctcca tccccagctt tc 32

<210> 51
<211> 37
<212> DNA
<213> Artificial

<220>
<223> primer, mouse origin

<400> 51
agctttggcc ggcctgggtcc ttggacctag gtccttc 37

<210> 52
<211> 32
<212> DNA
<213> Artificial

<220>
<223> primer, mouse origin

<400> 52
aaggcgcgcc cctggaagtc cacagtttct tc 32

<210> 53
<211> 35
<212> DNA
<213> Artificial

<220>
<223> primer, mouse origin

<400> 53
agctttgcga tcgctcagtc caagttcagc aatcg 35

<210> 54
<211> 38
<212> DNA
<213> Artificial

<220>
<223> primer, mouse origin

<400> 54
atagtttagc ggccgcccgg taggaaggcg gaattgag 38

<210> 55
<211> 35
<212> DNA
<213> Artificial

<220>
<223> primer, mouse origin

<400> 55
atagtttagc ggccgccagt tcccttagct ctgag 35

<210> 56
<211> 20
<212> DNA
<213> Artificial

<220>
<223> primer, mouse origin

<400> 56
tatcacaggt aggtgaccac 20

<210> 57
<211> 21
<212> DNA
<213> Artificial

<220>
<223> primer, mouse origin

<400> 57
acagccctgg gggggcaaca c 21

<210> 58
<211> 20
<212> DNA
<213> Artificial

<220>
<223> primer, mouse origin

<400> 58
gagccattgc taacggacag 20

<210> 59

<211> 20
<212> DNA
<213> Artificial

<220>
<223> primer, mouse origin

<400> 59
cctcatagct ccgaaggatg 20

<210> 60
<211> 20
<212> DNA
<213> Artificial

<220>
<223> primer, mouse origin

<400> 60
cagacgttcc caggctgtcg 20

<210> 61
<211> 20
<212> DNA
<213> Artificial

<220>
<223> primer, mouse origin

<400> 61
ctacagtaac cgagacctac 20

<210> 62
<211> 34
<212> DNA
<213> Artificial

<220>
<223> primer, mouse origin

<400> 62
agctttgttt aaactcaata cagtgcgggt caac 34

<210> 63
<211> 20
<212> DNA
<213> Artificial

<220>
<223> primer, mouse origin

<400> 63
cgacagcctg ggaacgtctg 20

<210> 64
<211> 22
<212> DNA

<213> Artificial

<220>

<223> primer, mouse origin

<400> 64

cctcgaatcg tggatccact ag

22

<210> 65

<211> 19

<212> DNA

<213> Artificial

<220>

<223> primer, mouse origin

<400> 65

tccctgcacg accttggtc

19

<210> 66

<211> 20

<212> DNA

<213> Artificial

<220>

<223> primer, mouse origin

<400> 66

gttctaccac ctcattatcg

20

<210> 67

<211> 20

<212> DNA

<213> Artificial

<220>

<223> primer, mouse origin

<400> 67

tttggaggagg aggctgaaag

20

<210> 68

<211> 20

<212> DNA

<213> Artificial

<220>

<223> primer, mouse origin

<400> 68

ggaagaaact gtggacttcc

20

<210> 69

<211> 20

<212> DNA

<213> Artificial

<220>

<223> primer, mouse origin

<400> 69

gtgaacctag ctctgttg

20

<210> 70

<211> 20

<212> DNA

<213> Artificial

<220>

<223> primer, mouse origin

<400> 70

accttggtct ctgagactcc

20

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/01858

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07K 16/00; A61K 39/395

US CL : 530/387.1, 387.3, 388.8, 388.85, 350, 391.7; 424/130/1, 133.1, 156.1, 181.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/387.1, 387.3, 388.8, 388.85, 350, 391.7; 424/130/1, 133.1, 156.1, 181.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 02/05843 (EXELIXIS, INC.) 24 January 2002 (24.01.02), see entire document, especially abstract.	1-7

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"B" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

13 May 2003 (13.05.2003)

Date of mailing of the international search report

23 JUN 2003

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Authorized officer

Larry R. Helms

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/01858

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claim Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
Please See Continuation Sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: 1-7

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

PCT/US03/01858

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, claim(s) 1-7, drawn to a method of inhibiting growth of tumor cells by administering an antibody.

Group II, claim(s) 8-12, drawn to a method of screening for agents.

Group III, claim(s) 13, drawn to a method of diagnosing a tumor cell by measuring the RRP expression.

Group IV, claim(s) 14-25, 29-32, drawn to a method of identifying a p53 or p21 modulating agent.

Group V, claim(s) 26-28, drawn to a method of modulating a p53 or p21 cell wherein the modulator binds to a RRP polypeptide and function is restored.

Group VI, claim(s) 33-35, drawn to a method of modulating a p53 or p21 cell wherein the modulator binds to a RRP polypeptide or nucleic acid.

Group VII, claim(s) 36-38, drawn to a method of diagnosing a disease.

Group VIII, claim(s) 39-45, drawn to a nucleic acid, host cell and expression system.

Group IX, claim(s) 46-49, drawn to a transgenic mouse.

Group X, claim(s) 50, drawn to a method of selecting an agent.

Group XI, claim(s) 51, drawn to a method of making an antibody.

The inventions listed as Groups I-XI do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: Pursuant to 37 C.F.R. 1.475 (d), the ISA/US considers that where multiple products and processes are claimed, the main invention shall consist of the first invention of the category first mentioned in the claims and the first recited invention of each of the other categories related thereto. Accordingly, the main invention of Group I, comprises the first method of inhibiting the growth of tumor cells by contacting the cells with an antibody. Further pursuant to 37 C.F.R. 1.475 (d), the ISA/US considers that any features which the subsequently recited products and methods share with the main invention does not constitute a special technical feature within the meaning of PCT Rule 13.2 and that each of such products and methods accordingly defines a separate invention.

INTERNATIONAL SEARCH REPORT

PCT/US03/01858

Continuation of B. FIELDS SEARCHED Item 3:

CAPLUS, MEDLINE WEST, USPATFULL, BIOSIS, PIR, SWISSPROT, GENESEQ, GENEMBLE

Search terms: SEQ ID NO:2, 4, 6, 8, 10, 12, 14, 16, RRP, antibody, rhomboid related protein, humanized, cytotoxic label.